arXiv:hep-th/0307169v1 18 Jul 2003

University of California
Los Angeles

Operators in the $d = 4$, $\mathcal{N} = 4$ SYM and the AdS/CFT correspondence

A dissertation submitted in partial satisfaction
of the requirements for the degree
Doctor of Philosophy in Physics

by

Anton Vladimirovich Ryzhov

2003

© Copyright by
Anton Vladimirovich Ryzhov
2003

The dissertation of Anton Vladimirovich Ryzhov is approved.

University of California, Los Angeles

2003

to my teachers

Table of Contents

1 Introduction **1**

1.1 Protected operators in SYM and SUGRA 2

1.1.1 Chiral primaries 2

1.1.2 Quarter BPS operators 3

1.1.3 Comparison with supergravity 4

1.2 Near the plane wave limit of AdS 5

2 Quarter BPS Operators in $\mathcal{N}$=4 SYM **7**

2.1 $SU(2,2|4)$ group theory 10

2.2 Gauge invariant scalar composite operators 13

2.3 Contributing diagrams 15

2.4 The simplest cases 18

2.4.1 Scalar composites with weight [2,0,2] 18

2.4.2 Scalar composites with weight [2,1,2] 22

2.5 A Gauge Invariance Argument 23

2.6 Operators of dimension 6 and higher 24

2.7 Large N analysis 29

2.7.1 Operators $\mathcal{O}_{[p,q,p]}$ and $\mathcal{K}_{[p,q,p]}$ 30

2.7.2 General correlators $\langle \mathcal{O}_{[p,q,p]} \bar{\mathcal{O}}_{[p,q,p]} \rangle$ to order g^2 31

2.7.3 General correlators $\langle \mathcal{K}_{[p,q,p]} \bar{\mathcal{O}}_{[p,q,p]} \rangle$ 37

2.7.4 General correlators $\langle \mathcal{K}_{[p,q,p]} \bar{\mathcal{K}}_{[p,q,p]} \rangle$ 39

2.7.5 Quarter BPS operators 40

2.8 Conclusion 42

2.9 Appendix 43

2.9.1 $\mathcal{N}$= 4 SUSY in various forms 43

2.9.2 Miscellaneous identities for $SU(N)$ 46

3 Systematics of Quarter BPS operators in $\mathcal{N} = 4$ SYM 49

3.1 $\mathcal{N} = 4$ SYM in harmonic superspace 50

3.1.1 (4,1,1) Harmonic Superspace 51

3.1.2 Quarter BPS Operators 53

3.1.3 Examples of systematic description 55

3.1.4 Multiplicity of quarter BPS operators 62

3.1.5 Relationship between $\mathcal{N} = 4$ and $\mathcal{N} = 1$ superfields 64

3.2 Explicit Computations 65

4 Three-Point Functions of Quarter BPS Operators in $\mathcal{N}$=4 SYM 77

4.1 Contributing diagrams 78

4.2 Restrictions from $\mathcal{N}$=4 SUSY and gauge invariance 80

4.3 Position dependence of $\tilde{B}$ and $\tilde{C}$ 81

4.4 Structure of the three-point functions 82

4.4.1 Space-time coordinate dependence 83

4.4.2 Group theory simplifications 84

4.5 Three-point functions of BPS operators 87

4.5.1 Correlators $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ 88

4.5.2 Three-point functions of $\frac{1}{4}$-BPS operators 97

4.6 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators in the large N limit 99

4.6.1 Large N operators 99

4.6.2 $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}}$, $\langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}}$, $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}}$, and $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}}$ . . 100

4.6.3 $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, $\langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, and $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$ 102

4.6.4 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators are protected 103

4.7 Supergravity considerations 104

4.7.1 OPE definition of $\frac{1}{4}$-BPS chiral primaries 105

4.7.2 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$ and $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}'_{\frac{1}{2}} \mathcal{O}_{\frac{1}{4}} \rangle$ correlators 106

4.7.3 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}'_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators 108

4.7.4 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}'_{\frac{1}{4}} \mathcal{O}''_{\frac{1}{4}} \rangle$ correlators 108

4.7.5 Detailed agreement between SYM and AdS 109

4.8 Conjectures 109

4.9 Appendix 111

4.9.1 $[p,q,p] \otimes [r,s,r] \otimes [0,k,0]$ and BZ triangles 111

4.9.2 Partitioning a tableau into 2 flavors 114

4.9.3 Details of $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$ calculations 115

5 Strings in the near plane wave background and AdS/CFT . . . 119

5.1 Corrections to the plane wave string spectrum 121

5.2 Anomalous dimensions and AdS/CFT 126

5.2.1 Operators 127

5.2.2 Two-point functions and the light cone Hamiltonian 132

5.2.3 Equality of matrix elements 134

5.3 Summary and further developments 140

5.4 Appendix 141

5.4.1 An alternative worldsheet discussion 141

5.4.2 $\mathcal{N}$=4 SYM 150

5.4.3 Equality of matrix elements: generic states 156

References 162

List of Figures

2.1 Structures contributing to two-point functions of scalars at order g^2 through four-scalar blocks and the propagator. Thick lines correspond to exchanges of the gauge boson, and of the auxiliary fields F_i and D (in the $\mathcal{N}$=1 formulation; after integrating out F_i and D, the $zz\bar{z}\bar{z}$ vertex). The scalar propagator remains diagonal in both color and flavor indices at order g^2. At order g^2, there are corrections to the scalar propagator coming from a fermion loop (dashed line) and a gauge boson semi-loop (wiggly line). Also, blocks involving four scalars get contributions from a single gauge boson exchange, and from the four-scalar vertex. Gauge fixing and ghost terms in the Lagrangian do not contribute to (2.12) at $\mathcal{O}(g^2)$. 16

2.2 Diagrams contributing to two-point functions of scalars at order g^2. 17

4.1 Building blocks for g^2 corrections to three-point functions. The three space-time points are x (with two legs attached) and y and w (single leg each). 79

4.2 F-term contributions to $\langle \mathcal{O}_{\rm BPS}(x)\, \mathcal{O}'_{\rm BPS}(y)\, \mathcal{O}_{\frac{1}{2}}(w)\rangle_{g^2}$ in the case when the correlator can be "partitioned into two flavors:" (a) proportional to $\tilde{B}(x,y)$; (b) proportional to $\tilde{C}(x;y,w)$; (c) proportional to $\tilde{C}(y;x,w)$. 85

4.3 Order g^2 corrections to correlators of the form (4.15): (a) and (b) includes a gauge boson exchange; (c) and (d) F-terms. Self energy contributions (not shown) also include a gauge boson exchange. . 86

4.4 Order g^2 corrections to extremal correlators: (a) and (b) within a single peddle; (c) and (d) between the two peddles. Self energy contributions (not shown) and diagrams (a) and (c) are gauge dependent, while (b) and (d) diagrams arise from the F-terms. . . 87

4.5 Nonvanishing Born diagrams for non-extremal 3-point functions $\langle \mathcal{O}_{[p,q,p]} \bar{\mathcal{O}}_{[p,q,p]} (\mathrm{tr}\, \bar{z} t z) \rangle$ (a-b); $\langle \mathcal{O}_{[p,q,p]} \bar{\mathcal{O}}_{[p-1,q+2,p-1]} (\mathrm{tr}\, \bar{z}_2 z_1) \rangle$ (c). . . 90

4.6 Order g^2 corrections to $\langle \mathcal{O}_{[p,q,p]}(x) \bar{\mathcal{O}}_{[p,q,p]}(y) (\mathrm{tr}\, \bar{z} t z)(w) \rangle$ with $t_{22} = 0$: (a) within the rainbow ($i, j = 1, 2$ in the second diagram); (b) from the rainbow to X^2 (there are similar ones with the other leg of X^2 uncorrected). 91

5.1 Order g^2_{YM} corrections to scalar propagators consist of a gauge boson exchange and a fermion loop. 152

5.2 Order g^2_{YM} corrections to two point functions of operators of the form $\mathrm{tr}\, z ... \phi^1 z ... \phi^2$: four-field irreducible blocks. When scalars ϕ^i are involved, the diagrams above represent the net contribution of all contributing Feynman diagrams, packaged in a way to mimic the $\mathcal{N}$=1 component fields Feynman diagrams. (Thick lines would correspond to exchanges of auxiliary fields F_i and D in the $\mathcal{N}$=1 formulation.) Diagrams with z_2 are given for comparison only. There are similar diagrams with one or both z-lines running in the opposite direction. 153

5.3 Order g^2 corrections to two point functions of operators of the form $\mathrm{tr}\, z ... \phi^1 z ... \phi^1$: four-field irreducible blocks. These diagrams represent the net contribution of all contributing Feynman diagrams. . 153

ACKNOWLEDGMENTS

First of all, I would like to express my gratitude to my thesis adviser Eric D'Hoker, for sharing with me his insights and physical intuition, and teaching me many things about physics — as well as about how to do physics. I am also in debt to my other advisers: I wish to thank Larry Yaffe, who taught me field theory; Nick Dorey, who introduced me to the supersymmetric world; and Per Kraus, from whom I learned a lot about gauge theories, supergravity, and other things. It is also my pleasure to thank Masaki Shigemori, Paul Heslop, Paul Howe, and especially Andrei Parnachev for wonderful collaborations. All these people have influenced my growth both as a physicist, and as a person.

Finally, I am grateful to my parents, my sister, and my grandmother, for their love and support, and for giving me the opportunity to carry out my Ph.D. studies.

Chapter 2 is a version of A. V. Ryzhov, "Quarter BPS operators in N = 4 SYM," JHEP **0111**, 046 (2001). Here I would like to thank Sergio Ferrara for useful discussions. I am especially grateful to Eric D'Hoker, for suggesting the problem and continuous involvement in this paper.

Chapter 3 is a version of E. D'Hoker, P. Heslop, P. Howe and A. V. Ryzhov, "Systematics of quarter BPS operators in N = 4 SYM," JHEP **0304**, 038 (2003).

Chapter 4 is a version of E. D'Hoker and A. V. Ryzhov, "Three-point functions of quarter BPS operators in N = 4 SYM," JHEP **0202**, 047 (2002). Here we wish to thank Daniel Z. Freedman and Sergio Ferrara for useful conversations during various stages of this project. We also thank the referee for suggesting clarifications on a number of issues.

Chapter 5 is a version of A. Parnachev and A. V. Ryzhov, "Strings in the

near plane wave background and AdS/CFT," JHEP **0210**, 066 (2002). Here we would like to thank Per Kraus, David Kutasov and David Sahakyan for many useful discussions. We are also grateful to Dan Freedman and Lubos Motl for sharing their insights on their construction of leading order SYM operators.

This research was supported in part by NSF Grants No. PHY-98-19686 and PHY-01-40151.

Abstract of the Dissertation

Operators in the $d = 4$, $\mathcal{N} = 4$ SYM and the AdS/CFT correspondence

by

Anton Vladimirovich Ryzhov
Doctor of Philosophy in Physics
University of California, Los Angeles, 2003

In this dissertation we explore various aspects of the AdS/CFT correspondence, which is a duality between $d = 4$, $\mathcal{N} = 4$ SYM, and IIB superstring theory on $AdS_5 \times S^5$ with selfdual RR field strength. String quantization on general backgrounds with fluxes is very difficult. So instead, one uses the duality at the level of canonical fields of supergravity and the corresponding $\frac{1}{2}$-BPS operators in SYM, since they both belong to the shortest multiplets of the superconformal group $SU(2, 2|4)$.

In addition to $\frac{1}{2}$-BPS operators, there are others with non-renormalization properties. One such class of operators is the $\frac{1}{4}$-BPS operators, which are dual to threshold bound states of elementary supergravity excitations. Their scaling dimension is also determined by their internal quantum numbers. In Chapter 2, we consider scalar composites with the right quantum numbers, and construct $\frac{1}{4}$-BPS operators as the ones with $\mathcal{O}(g^2_{\mathrm{YM}})$-protected two-point functions. Extended superspace methods (Chapter 3) make it simple to identify and remove descendant pieces from $\frac{1}{4}$-BPS candidates. In Chapter 4, we compute three-point functions involving $\frac{1}{4}$-BPS operators, and explain how their non-renormalization

translates into statements about the dual supergravity quantities.

But we can go beyond discussing supergravity modes and protected SYM operators. The GS superstring on $AdS_5 \times S^5$ can be quantized exactly in the limit where the AdS_5 radius $R \to \infty$ and the R-charge $J \sim R^2$. String states with finite energy and momentum (BMN states) are then dual to single trace operators with certain phases inserted (BMN operators). BMN operators are another natural generalization of $\frac{1}{2}$-BPS operators, to which they reduce in the zero-momentum limit.

The perturbative expansion of scaling dimensions of BMN operators is in powers of gN/J^2. Moreover, one can do perturbation theory around the $R \to \infty$, $J \sim R^2$ limit. Both expansions have the same regime of validity in string theory and in SYM. In Chapter 5, we calculate the first $1/R^2$ corrections to BMN states and their energies, and the $1/J$ corrections for the corresponding BMN operators. We find complete agreement between the dual quantities.

CHAPTER 1

Introduction

The celebrated AdS/CFT correspondence asserts that the dual description of $\mathcal{N}$=4 four dimensional super Yang Mills is type IIB string theory in $AdS_5 \times S^5$ with self-dual RR five-form field strength [1]. The radius of curvature of AdS_5 and S^5 scales like $R/l_s \sim (g^2_{\mathrm{YM}}N)^{1/4} \sim (gN)^{1/4}$. The spectrum of string states in this background corresponds to the spectrum of single trace operators in SYM. Part of the reason that the AdS/CFT conjecture has not been verified directly, is that string quantization in the presence of RR flux is notoriously difficult. Type IIB supergravity, which describes the dynamics of massless string modes, is only valid for the large values of R/l_s, while on the SYM side one can perform reliable computations only for small 't Hooft coupling gN: SYM and SUGRA calculations have complementary regimes of validity.

Thus on the one hand, the AdS/CFT conjecture still has not been proven directly, despite the abundance of circumstantial evidence. On the other hand, the AdS/CFT correspondence provides a powerful tool for deriving dynamical information in $\mathcal{N}$=4 superconformal YM theory outside the regime of weak coupling perturbation theory. Comparison of weak SYM coupling (small g^2_{YM}, perturbative gauge theory calculations) and strong SYM coupling (large AdS radius R, reliable supergravity calculations) behaviors has given rise to a number of surprising new conjectures [2, 3, 4, 5, 6, 7, 8, 9], which were later confirmed using extended superspace methods [10, 11, 12].

1.1 Protected operators in SYM and SUGRA

There are very few things one can calculate exactly in the four-dimensional superconformal Yang-Mills theories. One class of such quantities consists of a set of correlation functions of the Bogomolnyi Prasad Sommerfield (BPS) operators. In $\mathcal{N} = 4$ superconformal Yang-Mills, there are $\frac{1}{2}$-BPS, $\frac{1}{4}$-BPS and $\frac{1}{8}$-BPS operators, which are operators that are invariant under 8, 4 and 2 (out of 16) Poincaré supercharges respectively. Based on very general arguments involving only the supersymmetry algebra [13, 14, 15], the anomalous dimension of any of these operators vanishes identically in the full quantum theory.

1.1.1 Chiral primaries

In the AdS/CFT correspondence, the local gauge invariant operators of $\mathcal{N} = 4$ superconformal Yang-Mills are mapped to the physical states of the Type IIB superstring on $\mathrm{AdS}_5 \times \mathrm{S}^5$. (See [1] for the original papers and [16] for reviews.) The single trace $\frac{1}{2}$-BPS operators (also referred to as chiral primary operators or CPOs) play a special role as they are in one-to-one correspondence with the short multiplets of supergravity and Kaluza-Klein states with spins ≤ 2. Driven by the success of this correspondence, several authors have derived non-renormalization results for various correlation functions of these operators. Results on the perturbative non-renormalization of two- and three-point functions were derived in [2, 17, 3] in components and in superspace in [6, 11]; for further references on three-point computations see [18]. An argument for the complete non-renormalization of the three-point function of the supercurrent multiplet based on anomalies was given in [17] and a superspace version of this was presented in [11]. An argument for the (non-perturbative) non-renormalization of all two- and three-point functions of BPS operators based on an extra $U(1)_Y$

symmetry was given in [10] and this was verified using analytic superspace methods in [19] following on from the earlier work of [20]. Generalizations to n-point functions were obtained for extremal [8] and near-extremal correlators [9, 21, 22].

Other BPS operators are also important, both from the perspective of superconformal Yang-Mills theory, and from that of the AdS/CFT correspondence. The simplest generalization is to multi-trace 1/2 BPS operators [23, 12], for which non-renormalization results are the same as for single trace $\frac{1}{2}$-BPS operators; see also [24]. Indeed, the arguments of [10] and [19] apply in this case too.

1.1.2 Quarter BPS operators

A more delicate generalization is to the multi-trace scalar operators obeying a $\frac{1}{4}$-BPS shortening rule. A general group theoretic classification of such operators in free field theory was amongst the results derived in [25]. These $\frac{1}{4}$-BPS operators share many non-renormalization properties with $\frac{1}{2}$-BPS operators. However, they are much more involved, which renders their construction nontrivial in the fully interacting theory [26]. In the full quantum interacting theory, the true $\frac{1}{4}$-BPS operators involve admixtures of classical $\frac{1}{4}$-BPS operators of [25] with descendants of non-BPS operators that occur in long supersymmetry multiplets. In Chapter 2 we calculate $\mathcal{O}(g^2)$ two-point functions of local, polynomial, scalar composite operators within a given representation of the $SU(4)$ R-symmetry group. By studying these two-point functions, we identify the eigenstates of the dilatation operator, which turn out to be complicated mixtures of single and multiple trace operators.

However, this procedure for constructing the candidate operators is somewhat ad hoc and difficult to generalize. In Chapter 3, we use extended superspace methods to study $\frac{1}{4}$-BPS operators in a systematic fashion. In the framework of analytic superspace it turns out that they are described as tensor superfields carrying superindices [27]. In the construction, the operators of the classical theory annihilated by 4 out of 16 supercharges are arranged into two types. The first type consists of those operators that contain $\frac{1}{4}$-BPS operators in the full quantum theory. The second type consists of descendants of operators in long unprotected multiplets which develop anomalous dimensions in the quantum theory. The $\frac{1}{4}$-BPS operators of the quantum theory are defined to be orthogonal to all the descendant operators with the same classical quantum numbers. It is shown, to order $g^2_{\rm YM}$, that these $\frac{1}{4}$-BPS operators have protected dimensions. In fact, they are the same as the ones found in Chapter 2.

1.1.3 Comparison with supergravity

Using the operators thus constructed, in Chapter 4 we will compute three-point functions involving $\frac{1}{2}$-and $\frac{1}{4}$-BPS operators. The combinatorics of the problem is rather involved, and we concentrate on certain classes of three-point functions; some results are valid for general N, while others are large N approximations [28]. In all cases studied, these correlators[1] are shown to be non-renormalized to order $g^2_{\rm YM}$.

In the AdS/CFT correspondence, $\frac{1}{4}$-BPS chiral primaries are dual to threshold bound states of elementary supergravity excitations. We present a supergravity discussion of two- and three-point correlators involving these bound states, and show agreement of the SYM correlators with their large N, large $g^2_{\rm YM}N$ limit accessible through the AdS/CFT correspondence.

---

[1] Correlation functions with $n \geq 4$ operators are, in general, expected to receive quantum corrections, just as the multipoint functions of $\frac{1}{2}$-BPS operators do [29, 12]. See also [16] for further references.

1.2 Near the plane wave limit of AdS

So far, we have been studying the properties of supergravity modes, and the corresponding protected SYM operators, appealing to nonrenormalization theorems to compare their correlators in the dual descriptions [16].

An improvement to the $\frac{1}{2}$-BPS chiral primaries was constructed by Berenstein, Maldacena and Nastase (BMN) in [30]. The GS superstring can be quantized exactly in the plane wave background [31, 32], which can be viewed as a double scaling of the $AdS_5 \times S^5$ geometry [30, 33]. Remarkably, the parameter controlling perturbative expansion of scaling dimensions of such operators is $\lambda' = gN/J^2$, which can be made small to allow reliable gauge theory computations. String states with finite plane wave light cone energy and momentum correspond to single trace operators in the gauge theory with certain phases inserted [30, 34, 35, 36]. Later, string interactions were studied both in the plane wave string theory and in the gauge theory [36, 37, 38, 39, 40, 41, 42, 43, 44, 45].

The plane wave limit is an improvement over being able to handle only supergravity states and protected operators. But we would still like to get closer to the full AdS string theory. One way to gain insight is to do systematic perturbation theory around the plane wave limit, taking $1/R^2$ as a small parameter [46]. This approach has been tested in [47] on the $AdS_3 \times S^3$ background with NS-NS flux. In chapter 5, we study the AdS/CFT correspondence for string states which flow into plane wave states in the Penrose limit. Leading finite radius corrections to the string spectrum are compared with scaling dimensions of finite R-charge BMN-like operators. We find agreement between string and gauge theory results. This is a constructive step towards proving the AdS/CFT conjecture.[2]

---

[2] However, it should be mentioned that $1/R^2$ expansion around the BMN limit is still much closer to the plane wave geometry, than it is to doing string theory on the full AdS background.

The AdS/CFT correspondence is in turn just an example of the general phenomenon of holography, by which a CFT on the (conformal) boundary of some manifold is dual to a string theory in the bulk. The study of holography remains an open and exciting field, which keeps giving theorists valuable insights into both field theory phenomena, and into string theory. Hopefully, dualities will bring us closer to understanding confinement in gauge theories, or even to having a nonperturbative formulation of string theory.

CHAPTER 2

Quarter BPS Operators in $\mathcal{N}$=4 SYM

During the past years, there has been a renewed interest in the study of chiral operators in the $\mathcal{N}$=4 supersymmetric Yang-Mills theory in four dimensions. Forming short representations of the global $SU(2,2|4)$ superconformal symmetry group, chiral operators have tightly constrained quantum numbers. In particular, the scaling dimension of a chiral operator is not renormalized.[1]

Chiral primary operators have been classified in [25, 48]. They can be $\frac{1}{2}$-BPS, $\frac{1}{4}$-BPS, and $\frac{1}{8}$-BPS. The $\frac{1}{2}$-BPS operators provide the simplest example of chiral primaries. These are scalar composite operators in the $[0,q,0]$ representations of the R-symmetry group $SU(4) \sim SO(6)$; their scaling dimension is $\Delta = q$, see [25]. $\frac{1}{2}$-BPS operators are annihilated by eight out of the sixteen Poincaré supercharges of the theory. Similarly, $\frac{1}{4}$-BPS primaries belong to $[p,q,p]$ representations of the R-symmetry group, are annihilated by four supercharges, and have protected scaling dimension of $\Delta = 2p + q$. Finally, $\frac{1}{8}$-BPS primaries live in $[p,q,p+2k]$ of $SU(4)$, are killed by only two supercharges, and their $\Delta = 3k + 2p + q$. Quantum numbers of the descendant operators are related to those of their primaries by the $\mathcal{N}$=4 superconformal algebra.

$\frac{1}{2}$-BPS operators have been much studied. Using the conjectured AdS/CFT correspondence [1], it was shown, that for gauge groups $SU(N)$ with N large, two and three point functions of $\frac{1}{2}$-BPS chiral primaries are the same at weak and strong coupling [2].[2] It was then verified that these SYM correlators get no $\mathcal{O}(g^2)$ corrections, for all N [3]. Chiral descendant operators share these non-renormalization properties with their parent primaries [3]. $\mathcal{O}(g^4)$ and instanton contributions to two and three point functions of $\frac{1}{2}$-BPS primaries turn out to vanish as well [11, 7, 12, 6]. Non-renormalization of these correlators was further established on general grounds in [10, 11]. Besides $SU(N)$ theories and single trace chiral primaries, multiple trace operators with the same $SU(2,2|4)$ quantum numbers, as well as arbitrary gauge groups were considered [5]. In these cases, two and three point functions were also found to receive no $\mathcal{O}(g^2)$ corrections.

---

[1] The possibility that certain non-chiral operators may have vanishing anomalous dimension was raised in [12].

It is natural to ask whether other chiral operators, for example $\frac{1}{4}$-BPS primaries, have protected correlators. Here the situation is much less straightforward than for $[0,q,0]$ operators. In fact, except for the simplest operator found in [7], no other $\frac{1}{4}$-BPS chiral primaries were written down[3] in the fully interacting theory. The main difficulty is that unlike in the free theory, where a kinematical (group theoretical) treatment of [25] is sufficient, for nonzero coupling the problem of determining primary operators becomes a dynamical question.[4]

Apart from the double trace scalar composite operators in the $[p,q,p]$ of the R-symmetry (flavor) group $SU(4)$ (the free theory chiral primaries from the classification of [25]), there are other single and multiple trace scalar composites with the same $SU(4)$ quantum numbers and the same $\mathcal{O}(g^0)$ scaling dimension. Unlike in the $\frac{1}{2}$-BPS case where this phenomenon occurs [5], scalar composites in the $[p,q,p]$ generally do not have a well defined scaling dimension. Thus, one should first find their linear combinations which are eigenstates of the dilatation operator, which we call pure operators. To this end, we calculate two point functions of local, gauge invariant, polynomial, scalar composite operators in a given $[p, q, p]$ representation; diagonalize the dilatation operator within each representation of $SU(4)$; and find that some of the pure operators receive no $\mathcal{O}(g^2)$ corrections to their scaling dimension or normalization. These operators have the right $SU(4)$ quantum numbers and protected $\Delta = 2p + q$, and are the only candidates for being the $\frac{1}{4}$-BPS chiral primaries from the classification of [25].

---

[2] Higher n-point functions also agree with supergravity predictions in the large N limit [49].

[3] $\frac{1}{4}$-BPS operators have been studied indirectly through OPEs of $\frac{1}{2}$-BPS chiral primaries, see [7, 12, 50, 22].

[4] We would like to thank Sergio Ferrara for bringing this to our attention.

Calculating the symmetry factors for Feynman diagrams is a formidable combinatorial problem for general representation $[p, q, p]$ of $SU(4)$, and general N of the gauge group $SU(N)$. So to keep the formulas manageable, we concentrate on two special cases. For low dimensional operators ($2p+q < 8$), we perform explicit computations for arbitrary N; in particular, we recover the simplest $\frac{1}{4}$-BPS operator studied previously in [7]. Alternatively, we give a leading plus subleading large N argument (valid for general $[p, q, p]$ representations) for a class of $\frac{1}{4}$-BPS chiral primaries, which are linear combinations of double- and single-trace scalar composite operators.

The plan of this Chapter is as follows. First we review some aspects of $SU(2, 2|4)$ group theory, and describe the scalar composite operators we will be dealing with. Then we set the stage for $\mathcal{O}(g^2)$ calculations of two-point functions, and outline the main ingredients of these calculations. After that, we explicitly compute the simplest sets of correlators. In the course of these computations, it turns out that only one type of Feynman diagrams contributes to the correlators at order g^2, and we provide a simple explanation of this fact.[5] We present the full calculation for these two point functions. For higher Δ, calculations were done using *Mathematica* and only the results are shown. Several new features come into play, and we describe them as we go along. Finally, we switch gears and do a large N analysis of $\frac{1}{4}$-BPS operators with arbitrary scaling dimension.

---

[5] The argument we give applies more generally. In particular, it provides an alternative interpretation of the work in [3] and [5].

2.1 $SU(2,2|4)$ group theory

Four dimensional $\mathcal{N}$=4 superconformal Yang-Mills theory has been studied extensively for a long time, and we begin by reviewing some well known facts.

$\mathcal{N}$=4 SYM can be formulated in several (equivalent) ways; see Appendix 2.9.1 for some of the descriptions. None of them shows all the features of the theory explicitly. For example, working with six scalars $\phi^I = \phi^I_a t^a$ (where $a = 1, ..., N^2 - 1$ runs over the gauge group $SU(N)$, and $\phi^I_a(x)$, $I = 1, ..., 6$ are real scalar fields), and grouping the fermions as λ^i_a, $i = 1, ..., 4$, makes the full $SU(4)$ R-symmetry group manifest, but hides all the supersymmetries. On the other hand, formulating the theory in terms of $\mathcal{N}$=1 superfields shows some of the supersymmetry, but the Lagrangian looks invariant just under the $SU(3) \times U(1)$ subgroup of the full $SU(4)$. In practice, the more supersymmetries we use, the simpler it is to perform actual calculations.[6] For the purposes of $\mathcal{O}(g^2)$ computations, it suffices to use component fields of the $\mathcal{N}$=1 superfield formulation of the theory, with the (Euclidean signature) Lagrangian [3]

$$\begin{aligned}
\mathcal{L} \;=\; & \operatorname{tr}\left\{\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + \tfrac{1}{2}\bar{\lambda}\gamma^{\mu}D_{\mu}\lambda + \overline{D_{\mu}z_j}D^{\mu}z_j + \tfrac{1}{2}\bar{\psi}^j\gamma^{\mu}D_{\mu}\psi^j\right\} \\
& + i\sqrt{2}gf^{abc}\left(\bar{\lambda}_a\bar{z}^j_b L\psi^j_c - \bar{\psi}^j_a R z^j_b \lambda_c\right) - \tfrac{1}{2}Y f^{abc}\epsilon_{ijk}\left(\bar{\psi}^i_a z^j_b L\psi^k_c - \bar{\psi}^i_a R\bar{z}^j_b\psi^k_c\right) \\
& - \tfrac{1}{2}g^2(f^{abc}\bar{z}^j_b z^j_c)(f^{ade}\bar{z}^k_d z^k_e) + \tfrac{1}{4}Y^2 f^{abc}f^{ade}\epsilon_{ijk}\epsilon_{ilm}z^j_b z^k_c \bar{z}^l_d \bar{z}^m_e
\end{aligned} \tag{2.1}$$

(L and R are chirality projectors). The theory defined by (2.1) has $\mathcal{N}$=1 supersymmetry. We use separate coupling constants g and Y to distinguish the terms coming from the gauge and superpotential sectors. When $Y = g\sqrt{2}$, SUSY is enhanced to $\mathcal{N}$=4.

---

[6] E.g., the order g^4 calculations in [7] were done in the $\mathcal{N}$=2 harmonic superspace formalism.

Since the manifest symmetry group is now $SU(3) \times U(1)$, we first project onto it the representations of the full $SU(4)$. This can be done by mapping the quantum numbers as

$$[p, q, r] \mapsto [p, q]^{-\frac{1}{2}(p+2q+3r)} \tag{2.2}$$

Under this projection, the fermions in the theory are mapped as: $\lambda_{1,2,3} \mapsto \psi_{1,2,3} \in [1, 0]^{-\frac{1}{2}}$, $\lambda_4 \mapsto \lambda = [0, 0]^{\frac{3}{2}}$, so $\mathbf{4} = [1, 0, 0] \to [1, 0]^{-\frac{1}{2}} \oplus [0, 0]^{\frac{3}{2}}$. Similarly the scalars are projected as

$$\mathbf{6} \;=\; [0, 1, 0] \;\;\to\;\; [1, 0]^1 \oplus [0, 1]^{-1} \;=\; \{z_j\} \oplus \{\bar{z}^k\} \tag{2.3}$$

Put more simply, this amounts to rewriting the real scalars ϕ^I, and fermions λ^i as $\phi^i = \frac{1}{\sqrt{2}}(z_i + \bar{z}_i)$, $\phi^{i+3} = \frac{1}{i\sqrt{2}}(z_i - \bar{z}_i)$, and $\lambda^i = \psi_i$, $\lambda^4 = \lambda$. Index $i = 1, 2, 3$ labels the $\mathbf{3}$ or $\bar{\mathbf{3}}$ of the $SU(3)$ factor of the manifest symmetry group of (2.1).

The R-symmetry group of the theory is $SU(4) \sim SO(6)$, which is a part of the larger superconformal $SU(2, 2|4)$. Unitary representations of $\mathcal{N}$=4 SYM were classified in [48]. As in any conformal theory, operators are classified by their scaling dimension Δ. Each multiplet of $SU(2, 2|4)$ contains an operator of lowest dimension, which is called a primary operator. The action of generators of the conformal group[7] on a primary operator $\Phi(x)$ is given by

$$[P_\mu, \Phi(x)] \;=\; i\partial_\mu \Phi(x) \tag{2.4}$$

$$[M_{\mu\nu}, \Phi(x)] \;=\; [i(x_\mu\partial_\nu - x_\nu\partial_\mu) + \Sigma_{\mu\nu}]\, \Phi(x) \tag{2.5}$$

$$[D, \Phi(x)] \;=\; i\left(-\Delta + x^\mu\partial_\mu\right) \Phi(x) \tag{2.6}$$

$$[K_\mu, \Phi(x)] \;=\; \left[i(x^2\partial_\mu - 2x_\mu x^\nu\partial_\nu + 2x_\mu\Delta) - 2x^\nu\Sigma_{\mu\nu}\right] \Phi(x) \tag{2.7}$$

---

[7] See for example [51], or one of the big reviews [16].

Notice that $[M_{\mu\nu}, \Phi(0)] = \Sigma_{\mu\nu}\Phi(0)$, $[D, \Phi(0)] = -i\Delta\Phi(0)$, and $[K_\mu, \Phi(0)] = 0$. Together with the 16 Poincaré supersymmetry generators Q (and $\bar{Q}$), and 16 special conformal fermionic generators S (and $\bar{S}$), these close in a superconformal algebra of $SU(2,2|4)$. The additional (anti)commutation relations are schematically given by

$$[D, Q] = -\tfrac{i}{2}Q, \quad [D, S] = +\tfrac{i}{2}S, \quad [K, Q] \sim S \quad [P, S] \sim Q, \tag{2.8}$$

$$[Q, S] \sim M + D + R, \quad [S, S] \sim K, \quad [Q_i, Q_j] \sim P\,\delta_{ij}\ (i, j = 1, ..., 4) \tag{2.9}$$

where R stands for the quantum numbers of the R-symmetry group $SU(4)$. The Lagrangian of the theory, as well as the action of supersymmetry generators on the elementary fields, are listed in Appendix 2.9.1.

Primary operators of the superconformal group which are annihilated by at least some of the Q-s are called chiral primaries. Descendants of chiral primaries are then chiral operators, in the $\mathcal{N}$=4 sense. Chirality is a property of the whole $SU(2,2|4)$ multiplet; just being annihilated by say 8 Poincaré SUSY generators doesn't make an operator $\frac{1}{2}$-BPS. Since the supercharges anticommute, we can take a non-chiral operator and act on it with some of the Q-s. The resulting (non-chiral!) operator will be annihilated by the same Q-s.

For a chiral primary field Φ annihilated by a Poincaré supercharge Q, we can write $[Q, \Phi(x)] = 0$ and $[K, \Phi(0)] = 0$, and so $[S, \Phi(0)] \sim [[K, Q], \Phi(0)] = 0$ as well. Hence we can express the conformal dimension Δ of Φ entirely in terms of its spin Σ and $SU(4)$ quantum numbers R

$$0 = [[Q, S], \Phi(0)] \sim [M + D + R, \Phi(0)] = (\Sigma - i\Delta + R)\,\Phi(0) \tag{2.10}$$

by the superconformal algebra (2.4-2.9). Quantum numbers of descendants are related to those of their parent primaries by (2.4-2.9) as well. In particular, Δ of any chiral operator can not receive quantum corrections.

2.2 Gauge invariant scalar composite operators

A kinematic (group theoretic) classification of BPS operators was given in [25]. Chiral primaries[8] are Lorentz scalars, which are made by taking local gauge invariant polynomial combinations of the $\phi^I(x)$, $I = 1, ..., 6$. They fall into one of the three families [48]. The simplest one consists of $\frac{1}{2}$-BPS operators. These chiral primaries are annihilated by half of the Q-s, and live in short multiplets with spins ranging from zero to 2. $\frac{1}{2}$-BPS chiral primaries are totally symmetric traceless rank q tensors of the flavor $SO(6)$. $SU(4)$ labels of these representations are $[0, q, 0]$ with the corresponding $SO(6)$ Young tableau[9] $\begin{array}{|c|c|c|}\hline \ & \cdots & \ \\ \hline\end{array}$, one row of length q. Operators with the highest $SU(4)$ weight in the $[0, q, 0]$ have the form $\mathrm{tr}\,(\phi^1)^q$, modulo the $SO(6)$ traces.[10] Because the color group is $SU(N)$ rather than $U(N)$, $\mathrm{tr}\,\phi^I = 0$ so $q \geq 2$. Conformal dimension of a $\frac{1}{2}$-BPS chiral primary is related to its flavor quantum numbers as $\Delta = q$.

$\frac{1}{4}$-BPS operators form the next simplest family of chiral operators in the classification of [25]. Their multiplets have spins from zero to 3. The primaries belong to $[p, q, p]$ representations, and are annihilated by four out of sixteen Poincaré supercharges. There is a restriction $p \geq 2$: for $p = 0$ the operators are $\frac{1}{2}$-BPS; and in the case $p = 1$, they vanish after we take the $SU(N)$ traces. The highest weight state of $[p, q, p]$ corresponds to the

$$\begin{array}{|c|c|c|c|c|c|}\hline 1 & \cdots & 1 & 1 & \cdots & 1 \\ \hline 2 & \cdots & 2 & \multicolumn{3}{c}{q} \\ \cline{1-3} \multicolumn{3}{c}{p} \end{array} \tag{2.11}$$

$SO(6)$ Young tableau. In the free theory, $\frac{1}{4}$-BPS primaries corresponding to

---

[8] When referring to "primary" fields, we often have in mind the entire $SU(4)$ multiplet to which the actual primary belongs. This slight abuse of notation is common in the literature.

[9] See for example [52] for a general discussion on constructing irreducible tensors of $SO(n)$.

[10] For example, the highest weight state in the [2,0,2] is $\mathrm{tr}\,(\phi^1)^2 - \frac{1}{6}\sum_{I=1}^{6} \mathrm{tr}\,\phi^I\phi^I$. Operators in this representation are usually referred to as "$\mathrm{tr}\,X^2$" in the literature, and are special since their descendants include the $SU(4)$ flavor currents and the stress tensor.

(2.11) are of the form $\mathrm{tr}\,(\phi^1)^{p+q}\,\mathrm{tr}\,(\phi^2)^p$ (modulo (ϕ^1, ϕ^2) antisymmetrizations, and subtraction of the $SO(6)$ traces). However, there are many other ways to partition a given Young tableau, and each may result in a different operator after we take the $SU(N)$ traces. A priori, we do not know if any of them are pure (i.e. eigenstates of the dilatation operator D), or are mixtures of operators with different scaling dimensions. So these operators should be regarded just as a basis of gauge invariant, local, polynomial, scalar composite operators in the $[p, q, p]$ of $SU(4)$. By taking linear combinations of these, we will construct eigenstates of D in general, and $\frac{1}{4}$-BPS primaries in particular.

For completeness, let us mention the $\frac{1}{8}$-BPS operators, which form the last family of chiral operators in the classification of [25]. $\frac{1}{8}$-BPS multiplets are also short, with spins from zero to 7/2, and the chiral primaries are of the form $\mathrm{tr}\,(\phi^1)^{p+k+q}\,\mathrm{tr}\,(\phi^2)^{p+k}\,\mathrm{tr}\,(\phi^3)^k$ (modulo (ϕ^1, ϕ^2, ϕ^3) antisymmetrizations, and minus the $SO(6)$ traces), in the free theory. As before, there is a $k \geq 2$ restriction on the quantum numbers: $k \geq 1$ so the operators are annihilated by exactly two Poincaré supercharges; while operators with $k = 1$ necessarily contain commutators after we take the $SU(N)$ traces, as $\mathrm{tr}\,\phi^I = 0$. $\frac{1}{8}$-BPS chiral primaries have $SU(4)$ labels $[p, q, p + 2k]$, and their scaling dimensions have protected values of $\Delta = 3k + 2p + q$. Although these operators are also interesting, we will not study them here.

When calculating n-point functions, it suffices to consider one (nonzero) correlator for a given choice of representations; all others will be related to it by $SU(4)$ Clebsch-Gordon coefficients (by the Wigner-Eckart theorem). Therefore, we are free to take the most convenient representatives of the full $SU(4)$ representations, or of the smaller $SU(3) \times U(1)$ bits into which a given representation of

$SU(4)$ breaks down.[11] The combinatorics of the problem simplifies if we consider operators of the form $[(z_1)^{p+q}(z_2)^p]$ and their conjugates, which is what we will do in this Chapter.

Finally, suppose we have disentangled the mixtures of $[p,q,p]$ scalar composite operators annihilated by a quarter of the Poincaré supercharges, into linear combinations of operators with definite scaling dimension. Furthermore, assume we found an operator $\mathcal{Y}$ whose scaling dimension is protected. Since $\mathcal{Y}$ is a pure operator annihilated by four Poincaré supercharges, it can be either a $\frac{1}{4}$-BPS primary; or a level two descendant of a $\frac{1}{8}$-BPS primary, but this case is excluded[12] by group theory; or a level four descendant of a non-chiral primary. If $\mathcal{Y}$ were non-chiral, its primary would be a scalar composite operator of the form $[z^{2p+q-3}\bar{z}]$; and in all examples that we studied in this Chapter, such operators do receive $\mathcal{O}(g^2)$ corrections to their scaling dimension.[13] We conclude that a scalar composite operator in the $[p,q,p]$, which is annihilated by a quarter of the supercharges and has a protected scaling dimension $\Delta = 2p+q$, is a $\frac{1}{4}$-BPS chiral primary.

2.3 Contributing diagrams

The two point functions we will be calculating are of the form

$$\langle \left[{z_1}^{(p+q)}{z_2}^p\right](x)\ \left[\bar{z}_1^{(p+q)}\bar{z}_2^p\right](y)\rangle \tag{2.12}$$

where [...] stands for gauge invariant combinations. The free field part of such a correlator is given by a power of the free scalar propagator $[G(x,y)]^{(2p+q)}$, times

---

[11] All correlators in the resulting $SU(3)\times U(1)$ representations will have identical spatial dependence, since they come from the same $SU(4)$ representation.

[12] If $\mathcal{Y}$ came from a $\frac{1}{8}$-BPS primary, the parent primary would in the $[p',q',p'+2k]$ representation of $SU(4)$, with $k\geq 2$. On the other hand, to make the scaling dimension and $SU(4)$ Dynkin labels work out right, the only allowed choice is $[p,q,p+2]$, or $k=1$.

[13] But see footnote 1.


i, a
i, a'
j, b
j, b'
+
=
= $f^{apa'}f^{bpb'}B\ (x,y)\ G(x,y)^2$
= $f^{apb}f^{a'pb}\tilde{B}\ (x,y)\ G(x,y)^2$
= $\delta^{ab}\delta^{ij}\ N\ A(x,y)\ G(x,y)$

Figure 2.1: Structures contributing to two-point functions of scalars at order g^2 through four-scalar blocks and the propagator. Thick lines correspond to exchanges of the gauge boson, and of the auxiliary fields F_i and D (in the $\mathcal{N}$=1 formulation; after integrating out F_i and D, the $zz\bar{z}\bar{z}$ vertex). The scalar propagator remains diagonal in both color and flavor indices at order g^2. At order g^2, there are corrections to the scalar propagator coming from a fermion loop (dashed line) and a gauge boson semi-loop (wiggly line). Also, blocks involving four scalars get contributions from a single gauge boson exchange, and from the four-scalar vertex. Gauge fixing and ghost terms in the Lagrangian do not contribute to (2.12) at $\mathcal{O}(g^2)$.

a combinatorial factor.

From the Lagrangian (2.1) we can read off the structures contributing to the four-scalar blocks, and the leading correction to the propagator at order g^2. These are shown in Figure 2.1, where they are categorized according to their gauge group (color) index structure (we will use the same notation as in [3]). The scalar propagator remains diagonal in both color and flavor indices at order g^2. Notice that the corrections proportional to $\tilde{B}$ are antisymmetric in i and j, hence they are absent when the scalars in the four legs have the same flavor.



Figure 2.2: Diagrams contributing to two-point functions of scalars at order g^2.

Thus we will have to compute contributions of six types[14] (see Figure 2.2).

Most Feynman diagrams we come across are easier to evaluate in position space, where they factorize into products of free propagators and the blocks shown in Figure 2.1, and everything except for the combinatorial factors out front is almost trivial. In momentum space, on the other hand, even the simplest $\mathcal{O}(g^2)$ graphs contain divergent subdiagrams.

The functions A and B will be discussed in detail in Section 2.5. Coordinate dependence of $\tilde{B}$ is parametrically determined by conformal invariance, $\tilde{B}(x,0) = \tilde{a}\log(x^2\mu^2) + \tilde{b}$. The coefficients $\tilde{a}$ and $\tilde{b}$ can be found using, for example, differential regularization [53], or a simpler equivalent prescription: replace $1/x^2 \to 1/(x^2+\epsilon^2)$ for propagators inside integrals ($\epsilon \sim \mu^{-1}$ is related to the renormalization scale). With this,

$$\begin{aligned}\tilde{B}(x,0) &= -\tfrac{1}{4}Y^2 \int \frac{(d^4z)\,[4\pi^2 x^2]^2}{[4\pi^2((z-x)^2+\epsilon^2)]^2\,[4\pi^2(z^2+\epsilon^2)]^2} \\ &= -Y^2\frac{1}{32\pi^2}\left[\log(x^2/\epsilon^2) - 1\right] \qquad (2.13)\end{aligned}$$

(for $\mathcal{N}$=4 SUSY, $Y^2 = 2g^2$); the same result is obtained in dimensional regularization.

---

[14] If all scalars were of the same flavor, say 1 (as is the case for $\frac{1}{2}$-BPS operators considered in [3]and [5]), we would only have to consider diagrams of types (a) and (d).

2.4 The simplest cases

We begin by considering scalar composite operators in representations $[p, q, p]$ of the color $SU(4)$, which have $2p + q = 4$ and 5.

The case of $\Delta = 4 + \mathcal{O}(g^2)$ has been studied before. For example, the authors of [7] argued that there are two operators[15] $\mathcal{O}_1^{[2,0,2]}$ and $\mathcal{O}_2^{[2,0,2]}$, which are made of four scalars and annihilated by four supercharges. $\mathcal{O}_1^{[2,0,2]}$ is a descendant of the Konishi scalar $\left(\sum_{I=1}^{6} \operatorname{tr} \phi^I \phi^I\right)$ and therefore is pure (i.e. is an eigenstate of the dilatation operator), since the Konishi scalar is pure. The other operator, $\mathcal{O}_2^{[2,0,2]}$, contains a piece proportional to $\mathcal{O}_1^{[2,0,2]}$, but the rest is a chiral primary. The method in [7] was to analyze four-point correlators of certain $\frac{1}{2}$-BPS operators, and to look at the possible operators in exchange channels. They found that there is a $\frac{1}{4}$-BPS operator exchanged by demonstrating that there is a pole corresponding to an operator of scaling dimension $\Delta = 4$. They determined this operator to be $\mathcal{Y}_{[2,0,2]} = \mathcal{O}_2^{[2,0,2]} - \frac{4}{N}\mathcal{O}_1^{[2,0,2]}$.

Unfortunately, this method does not generalize to chiral primaries with scaling dimension $\Delta \geq 6$, as we shall see in Section 2.6. So instead we explicitly compute two-point functions of scalar composite operators of a given scaling dimension, and find the ones which do not get corrected. This allows us to fix the normalization of $\frac{1}{4}$-BPS operators as well.

2.4.1 Scalar composites with weight [2,0,2]

The simplest operators annihilated by four out of sixteen Poincaré supercharges correspond to the highest weight state of the **84** = [2,0,2] of $SU(4)$. The $SO(6)$ Young tableau for representation is ⊞. An $SO(6)$ irreducible tensor T with this symmetry is made from the corresponding $Gl(6)$ irreducible tensor T^0 by subtracting all possible $SO(6)$ traces:

---

[15] For the notation and definitions, see Section 2.4.1.

$$T_{\begin{array}{|c|c|}\hline a&b\\\hline c&d\\\hline\end{array}} = T^0_{\begin{array}{|c|c|}\hline a&b\\\hline c&d\\\hline\end{array}} - \frac{1}{4}\Big(T^0_{\begin{array}{|c|c|}\hline \bullet&\bullet\\\hline a&b\\\hline\end{array}}\delta_{cd} + T^0_{\begin{array}{|c|c|}\hline \bullet&\bullet\\\hline c&d\\\hline\end{array}}\delta_{ab} + T^0_{\begin{array}{|c|c|}\hline \bullet&\bullet\\\hline a&d\\\hline\end{array}}\delta_{bc} + T^0_{\begin{array}{|c|c|}\hline \bullet&\bullet\\\hline b&c\\\hline\end{array}}\delta_{ad}\Big)$$
$$+\frac{1}{20}\left(\delta_{ab}\delta_{cd} - \delta_{ad}\delta_{bc}\right) T^0_{\begin{array}{|c|c|}\hline \bullet&\bullet\\\hline \bullet&\bullet\\\hline\end{array}} \tag{2.14}$$

where $\begin{array}{|c|c|}\hline \bullet&\bullet\\\hline\end{array} = \sum_{a=1}^{6} \begin{array}{|c|c|}\hline a&a\\\hline\end{array}$. Recall that the $SU(4) \to SU(3) \times U(1)$ projection (2.2) is realized on the elementary fields as $\phi_a = \frac{1}{\sqrt{2}}(z_a + \bar{z}_a)$, $\phi_{a+3} = \frac{1}{i\sqrt{2}}(z_a - \bar{z}_a)$, $a = 1, 2, 3$. Under this "3+1 split," the highest weight state of [2,0,2] becomes

$$T_{\begin{array}{|c|c|}\hline 1&1\\\hline 2&2\\\hline\end{array}} = \frac{1}{4}\Big(T_{\begin{array}{|c|c|}\hline \mathbf{1}&\mathbf{1}\\\hline \mathbf{2}&\mathbf{2}\\\hline\end{array}} + 2T_{\begin{array}{|c|c|}\hline \bar{\mathbf{1}}&\mathbf{1}\\\hline \mathbf{2}&\mathbf{2}\\\hline\end{array}} + 2T_{\begin{array}{|c|c|}\hline \mathbf{1}&\mathbf{1}\\\hline \bar{\mathbf{2}}&\mathbf{2}\\\hline\end{array}} + 2T_{\begin{array}{|c|c|}\hline \bar{\mathbf{1}}&\mathbf{1}\\\hline \bar{\mathbf{2}}&\mathbf{2}\\\hline\end{array}} + T_{\begin{array}{|c|c|}\hline \bar{\mathbf{1}}&\bar{\mathbf{1}}\\\hline \mathbf{2}&\mathbf{2}\\\hline\end{array}}\Big) + \text{c.c.}$$
$$= \frac{1}{4}T^0_{\begin{array}{|c|c|}\hline \mathbf{1}&\mathbf{1}\\\hline \mathbf{2}&\mathbf{2}\\\hline\end{array}} + \text{terms with lower } U(1) \text{ charge} \tag{2.15}$$

where in the left hand side, $1 = \phi^1$; and in the right hand side, $\mathbf{1} = z_1$, $\bar{\mathbf{1}} = \bar{z}_1$, etc. We see that after this projection, we don't have to worry about subtracting the $SO(6)$ traces, if we are only interested in the highest $U(1)$ charge operators. Henceforth, we will consider operators made by applying the Young (anti)symmetrizers corresponding to this tableau, to the string of z_i-s.[16]

We can construct one single trace and one double trace operators with the highest [2,0,2] weight. When projected onto $SU(3) \times U(1)$, they become

$$\mathcal{O}_1^{[2,0,2]} = \text{tr}\, z_1 z_1 z_2 z_2 - \text{tr}\, z_1 z_2 z_1 z_2 = -\tfrac{1}{2}\text{tr}\,[z_1, z_2][z_1, z_2] \tag{2.16}$$
$$\mathcal{O}_2^{[2,0,2]} = 2\left(\text{tr}\, z_1 z_1\ \text{tr}\, z_2 z_2 - \text{tr}\, z_1 z_2\ \text{tr}\, z_1 z_2\right) \tag{2.17}$$

$\mathcal{O}_1^{[2,0,2]}$ is a descendant; by consecutively applying four SUSY transformations,[17] it can be obtained from the Konishi scalar, $\mathcal{K}_0 = \text{tr}\, z_j \bar{z}^j$. More explicitly, acting with the SUSY generator $\bar{Q}_{\bar{\zeta}}$ gives

---

[16] Computing two point functions of operators in representations of $SU(3) \times U(1)$ is as good as computing two point functions of the original $SU(4)$ irreducible operators, see Section 2.1. In the following, we will neglect the $SO(6)$ traces without a comment.

[17] Supersymmetry transformations are listed in Appendix 2.9.1, see equations (2.90-2.96).

$$\delta_{\bar{\zeta}} z_j = 0, \quad \delta_{\bar{\zeta}} \bar{z}^j = \sqrt{2}\bar{\zeta}\bar{\psi}^j, \quad \delta_{\bar{\zeta}} \bar{\psi}^j = i\epsilon^{jkl}[z_k, z_l]\bar{\zeta}, \tag{2.18}$$

and with Q_{ζ_3},

$$\delta_{\zeta_3} z_j = -\sqrt{2}\zeta_3\lambda\delta_{3j}, \quad \delta_{\zeta_3}\lambda = 2i[z_1, z_2]\zeta_3, \tag{2.19}$$

thus

$$\begin{aligned}
(\bar{Q}_{\bar{\zeta}})^2\mathcal{K}_0 &= \bar{Q}_{\bar{\zeta}}\mathrm{tr}\, z_j\sqrt{2}\bar{\zeta}\bar{\psi}^j = 6i\sqrt{2}(\bar{\zeta}\bar{\zeta})\mathrm{tr}\,[z_1, z_2]z_3, \\
(Q_{\zeta_3})^2(\bar{Q}_{\bar{\zeta}})^2\mathcal{K}_0 &= -12i(\bar{\zeta}\bar{\zeta})Q_{\zeta_3}\mathrm{tr}\,[z_1, z_2]\zeta_3\lambda = 24(\bar{\zeta}\bar{\zeta})(\zeta_3\zeta_3)\mathrm{tr}\,[z_1, z_2]^2 \\
&= -48(\bar{\zeta}\bar{\zeta})(\zeta_3\zeta_3)\mathcal{O}_1^{[2,0,2]}
\end{aligned} \tag{2.20}$$

The four generators which annihilate $\mathcal{O}_1^{[2,0,2]}$, are the ones we acted with to obtain it from $\mathcal{K}_0$. However, since $\mathcal{K}_0$ is a non-chiral primary, its descendant $\mathcal{O}_1^{[2,0,2]}$ is not $\frac{1}{4}$-BPS, despite being annihilated by a quarter of SUSY generators.

The free field results for two point functions of $\mathcal{O}_1^{[2,0,2]}$ and $\mathcal{O}_2^{[2,0,2]}$ are

$$\begin{pmatrix} \langle\mathcal{O}_1\bar{\mathcal{O}}_1\rangle & \langle\mathcal{O}_1\bar{\mathcal{O}}_2\rangle \\ \langle\mathcal{O}_2\bar{\mathcal{O}}_1\rangle & \langle\mathcal{O}_2\bar{\mathcal{O}}_2\rangle \end{pmatrix}_{\text{free}} = \frac{3(N^2-1)G^4}{16}\begin{pmatrix} N^2 & 4N \\ 4N & 8(N^2-2) \end{pmatrix} \tag{2.21}$$

while the leading corrections are found to be

$$\begin{pmatrix} \langle\mathcal{O}_1\bar{\mathcal{O}}_1\rangle & \langle\mathcal{O}_1\bar{\mathcal{O}}_2\rangle \\ \langle\mathcal{O}_2\bar{\mathcal{O}}_1\rangle & \langle\mathcal{O}_2\bar{\mathcal{O}}_2\rangle \end{pmatrix}_{g^2} = \frac{9(N^2-1)G^4(\tilde{B}N)}{16}\begin{pmatrix} N^2 & 4N \\ 4N & 16 \end{pmatrix}; \tag{2.22}$$

here $\langle\mathcal{O}_i\bar{\mathcal{O}}_j\rangle \equiv \langle\mathcal{O}_i^{[2,0,2]}(x)\bar{\mathcal{O}}_j^{[2,0,2]}(y)\rangle$, and $G \equiv G(x,y)$, $\tilde{B} \equiv \tilde{B}(x,y)$. Some helpful formulae we used for deriving (2.21-2.22) are collected in Appendix 2.9.2.

By looking at (2.22), we conclude that neither $\mathcal{O}_1^{[2,0,2]}$ nor $\mathcal{O}_2^{[2,0,2]}$ are chiral. However, there is a linear combination of these two operators which has protected two point functions at order g^2. The operator

$$\mathcal{Y}_{[2,0,2]}(x) \equiv \mathcal{O}_2^{[2,0,2]}(x) - \frac{4}{N}\mathcal{O}_1^{[2,0,2]}(x) \tag{2.23}$$

satisfies $\langle \mathcal{Y}\bar{\mathcal{Y}} \rangle = \langle \mathcal{Y}\bar{\mathcal{O}}_1 \rangle = 0$, so $\mathcal{Y}_{[2,0,2]}$ is orthogonal to the descendant of the Konishi operator $\mathcal{O}_1^{[2,0,2]}$, and has protected dimension $\Delta_{\mathcal{Y}} = 4$ at order g^2. Computationally, this cancellation is rather intricate: all $\langle \mathcal{O}_i \bar{\mathcal{O}}_j \rangle$ have very different large N behavior.

We can also calculate the two point function of the Konishi scalar with itself:

$$\langle \mathcal{K}_0(x)\bar{\mathcal{K}}_0(y) \rangle = 3(N^2-1)[G(x,y)]^2 \left\{ 1 + 3\tilde{B}(x,y)N + \mathcal{O}(g^4) \right\} \quad (2.24)$$

so $\langle \mathcal{O}_1^{[2,0,2]}(x)\bar{\mathcal{O}}_1^{[2,0,2]}(y) \rangle = \frac{1}{16}N^2[G(x,y)]^2 \langle \mathcal{K}_0(x)\bar{\mathcal{K}}_0(y) \rangle + \mathcal{O}(g^4)$, which is just the free theory result. In particular, $\mathcal{K}_0$ and its descendant $\mathcal{O}_1^{[2,0,2]}$ have the same normalization and their scaling dimensions differ by 2, as they must: with $\tilde{B}(x,0)$ given by (2.13), the scaling dimension of $\mathcal{O}_1^{[2,0,2]}$ is $\Delta_1 = 4 + \frac{3g^2N}{16\pi^2} + \mathcal{O}(g^4)$, and that of $\mathcal{K}_0$ is $\Delta_{\mathcal{K}} = 2 + \frac{3g^2N}{16\pi^2} + \mathcal{O}(g^4)$, in agreement with [7]. The mixture $\mathcal{O}_2^{[2,0,2]} = \mathcal{Y}_{[2,0,2]} + \frac{4}{N}\mathcal{O}_1^{[2,0,2]}$ has the two-point function with itself which breaks down into two pieces,

$$\begin{aligned} \langle \mathcal{O}_2^{[2,0,2]}(x)\bar{\mathcal{O}}_2^{[2,0,2]}(y) \rangle &= \langle \mathcal{Y}_{[2,0,2]}(x)\bar{\mathcal{Y}}_{[2,0,2]}(y) \rangle + \frac{16}{N^2} \langle \mathcal{O}_1^{[2,0,2]}(x)\bar{\mathcal{O}}_1^{[2,0,2]}(y) \rangle \\ &= \frac{C_{\mathcal{Y}}}{(x-y)^{2\Delta_{\mathcal{Y}}}} + \frac{16}{N^2}\frac{C_1}{(x-y)^{2\Delta_1}} \end{aligned} \quad (2.25)$$

Logarithmic corrections to $\langle \mathcal{O}_2^{[2,0,2]}(x)\bar{\mathcal{O}}_2^{[2,0,2]}(y) \rangle$ are due entirely to the second term $\frac{16}{N^2}\langle \mathcal{O}_1^{[2,0,2]}(x)\bar{\mathcal{O}}_1^{[2,0,2]}(y) \rangle$.

$\mathcal{O}(g^2)$ corrections to all two-point functions just considered are proportional to $\tilde{B}$. In other words, only the contributions due to diagrams of type (c) in Figure 2.2 survive, and all other corrections cancel. As we will show in Section 2.5, this is a general phenomenon. Also, in the large N limit the fraction $\langle \mathcal{O}_1\bar{\mathcal{O}}_2 \rangle / \langle \mathcal{O}_1\bar{\mathcal{O}}_1 \rangle$ is suppressed; it vanishes in the limit $N \to \infty$, g^2N fixed.[18]

One can show that conformal dimension of $\mathcal{Y}_{[2,0,2]}$ is protected perturbatively (to order g^4) and nonperturbatively (for any instanton number) as well, see [7].

---

[18] The large N limit will be analyzed in more detail in Section 2.7.

Non-renormalization of scaling dimension of $\mathcal{Y}$ hints at its BPS property. Following the authors of [7], we suggest that it is indeed a $\frac{1}{4}$-BPS chiral primary operator.

2.4.2 Scalar composites with weight [2,1,2]

The story is similar in the case of the **300** $= [2,1,2]$ of $SU(4)$. The only scalar composite operators corresponding to the highest weight of this representation are (after the projection onto $SU(3) \times U(1)$, as discussed in Sections 2.1 and 2.4.1)

$$\begin{aligned} \mathcal{O}_1^{[2,1,2]} &= \mathrm{tr}\, z_1 z_1 z_1 z_2 z_2 - \mathrm{tr}\, z_1 z_1 z_2 z_1 z_2 = -\tfrac{1}{2}\mathrm{tr}\,[z_1, z_2][z_1^2, z_2] \qquad (2.26) \\ \mathcal{O}_2^{[2,1,2]} &= \mathrm{tr}\, z_1 z_1 z_1 \;\mathrm{tr}\, z_2 z_2 - 2\mathrm{tr}\, z_1 z_2 \;\mathrm{tr}\, z_1 z_2 z_1 + \mathrm{tr}\, z_1 z_2 z_2 \;\mathrm{tr}\, z_1 z_1 \qquad (2.27) \end{aligned}$$

$\mathcal{O}_1^{[2,1,2]}$ is a descendant; $(Q_{\zeta_3})^2(\bar{Q}_{\bar{\zeta}})^2 \mathrm{str}\, z_1 z_i \bar{z}^i \propto \mathcal{O}_1^{[2,1,2]}$ just like in Equation (2.20). Born level and order g^2 two point functions are

$$\begin{pmatrix} \langle \mathcal{O}_1 \bar{\mathcal{O}}_1 \rangle & \langle \mathcal{O}_1 \bar{\mathcal{O}}_2 \rangle \\ \langle \mathcal{O}_2 \bar{\mathcal{O}}_1 \rangle & \langle \mathcal{O}_2 \bar{\mathcal{O}}_2 \rangle \end{pmatrix} = \frac{(N^2-1)(N^2-4)G^5}{16N} \left\{ \begin{pmatrix} N^2 & 6N \\ 6N & 6(N^2-3) \end{pmatrix} + 4\tilde{B}N \begin{pmatrix} N^2 & 6N \\ 6N & 36 \end{pmatrix} + \mathcal{O}(g^4) \right\} \qquad (2.28)$$

where $\langle \mathcal{O}_i \bar{\mathcal{O}}_j \rangle \equiv \langle \mathcal{O}_i^{[2,1,2]}(x) \bar{\mathcal{O}}_j^{[2,1,2]}(y) \rangle$, and $G \equiv G(x,y)$, $\tilde{B} \equiv \tilde{B}(x,y)$ as before. (Note that corrections proportional to A and B cancel again.) There is a linear combination of $\mathcal{O}_1^{[2,1,2]}$ and $\mathcal{O}_2^{[2,1,2]}$

$$\mathcal{Y}_{[2,1,2]}(x) \equiv \mathcal{O}_2^{[2,1,2]}(x) - \frac{6}{N}\mathcal{O}_1^{[2,1,2]}(x) \qquad (2.29)$$

whose two point functions with arbitrary operators do not receive perturbative order g^2 corrections. Again, it seems reasonable to conclude that $\mathcal{Y}_{[2,1,2]}$ is a $\frac{1}{4}$-BPS operator, as it is annihilated by four out of sixteen supercharges, has a protected scaling dimension $\Delta_{\mathcal{Y}} = 5$ (at order g^2), and contains no descendant pieces, $\langle \mathcal{Y}_{[2,1,2]}(x) \bar{\mathcal{O}}_1^{[2,1,2]}(y) \rangle = 0$.

2.5 A Gauge Invariance Argument

In Sections 2.4.1 and 2.4.2, we explicitly calculated the $\mathcal{O}(g^2)$ corrections to two point functions of scalar composite operators. We found that corrections proportional to A and B cancel, i.e. gauge group combinatorics demands that diagrams containing a gauge boson exchange do not arise in the correlator. Here we give a general derivation of this fact, which boils down to gauge invariance of the operators in question, and gauge dependence of A and B.

The two point functions we have been considering are of the form

$$\langle [z^m](x)\ [\bar{z}^m](y) \rangle \tag{2.30}$$

where $[z^m](x)$ is some gauge-invariant homogeneous polynomial (of degree m) in the z_i^a-s. Diagrams involving a gauge boson exchange which contribute to the two-point functions of the form (2.30), are proportional to either $A(x,y)$ or $B(x,y)$, see Figure 2.1. By using nonrenormalization of the two point function $\langle \mathrm{tr}\, z_1 z_2(x)\ \mathrm{tr}\, \bar{z}_1 \bar{z}_2(y) \rangle$, one can immediately see [3] that $B(x,y) = -2A(x,y)$, so

$$\langle [z^m]\,(x)\,[\bar{z}^m]\,(y) \rangle_{(A+B)} \quad = \quad c_g A(x,y) [G(x,y)]^m \tag{2.31}$$

where c_g is some combinatorial coefficient.

Conformal invariance restricts $A(x,y) = a \log x^2\mu^2 + b$. The constants a and b turn out to be gauge dependent. The gauge fixing parameter ξ enters the expression for the scalar propagator as[19]



$$\begin{aligned} &= \xi g^2 \mu^{4-d} \frac{1}{p^4} \int \frac{(d^dk)}{(2\pi)^d} \frac{[(2p+k)\cdot k]^2}{k^4 (p+k)^2} + (\xi\text{-independent}) \\ &= \xi g^2 \tfrac{1}{2} \pi^{2+\epsilon} \frac{1}{(p^2)^{1-\epsilon} \mu^{2\epsilon}} \left[\frac{1}{\epsilon} + \gamma + \mathcal{O}(\epsilon) \right] + (\xi\text{-independent}) \end{aligned} \tag{2.32}$$

---

[19] By changing ξ, we can vary both the pole piece and the $\mathcal{O}(\epsilon^0)$ term (but we can't make them both zero simultaneously; the ξ-independent part is proportional to a different integral). Compare this with [54], where order g^2 corrections to the scalar propagators were found to vanish in super-Feynman gauge of $\mathcal{N}$=1 formulation of the theory.

in momentum space (in dimensional regularization [55]; $\epsilon = \frac{d}{2} - 2$ and γ is Euler's gamma constant), so in position space

$$\begin{aligned} A(x,0) &= \tfrac{1}{2}\pi^2 g^2 \xi \left[\log x^2\mu^2 + \log 4\pi - \gamma\right] + (\xi\text{-independent}) \\ &\equiv a \log x^2\mu^2 + b \end{aligned} \tag{2.33}$$

after factoring out the free propagator $G(x,0)$. Both a and b have pieces linear in the gauge fixing parameter ξ.

Since a correlator of gauge invariant operators must be gauge independent, the combinatorial coefficient multiplying $A(x,y)$ in equation (2.31) must vanish; we necessarily have $c_g = 0$. This is a general phenomenon, illustrated by an explicit calculation of Sections 2.4.1 and 2.4.2: gauge dependent contributions are proportional to $2A + B = 0$.

In the $\mathcal{O}(g^2)$ calculations of correlators of $\frac{1}{2}$-BPS operators [3] and [5], there were no other contributions to two-point functions except for A and B. Thus, gauge invariance together with $\mathcal{N}$=4 SUSY (which is needed to make $2A+B=0$) guarantees that the correlators of [3] and [5] receive no order g^2 corrections.

2.6 Operators of dimension 6 and higher

At this point, we would like to consider operators made of $2p+q \geq 6$ scalar fields. According to the classification of [25], these $[p,q,p]$ operators are the candidates for $\frac{1}{4}$-BPS chiral primaries. However, a new complication arises compared to the cases of $2p+q \leq 5$ studied in Sections 2.4.1 and 2.4.2. Now, there are many ways in which we can make gauge invariant combinations of fields, and hence many scalar composites have to be taken into account. Apart from single and double trace operators we have seen so far, operators made of three or more traces also have to be considered.

This phenomenon has a counterpart in the context of $\frac{1}{2}$-BPS operators, see [5]. The crucial difference is that in our case, none of the scalar composites are pure, and only some special mixtures have a well defined scaling dimension. In general, the "naive" scalar composite operators will have nonvanishing two point functions with each other, whenever this is allowed by group theory. Unlike in the simplest cases of Section 2.4, operators containing commutators are not pure, but contain pieces which are descendants of different operators.

To find pure operators, in this Section we calculate the two point functions of highest weight $[p,q,p]$ scalar composites $\mathcal{O}_i^{[p,q,p]}$, and arrange them as[20]

$$\langle \mathcal{O}_i^{[p,q,p]}(x)\bar{\mathcal{O}}_j^{[p,q,p]}(y)\rangle \equiv [G(x,y)]^{(2p+q)}\left[\mathbf{F}_{ij}+\tilde{B}(x,y)N\,\mathbf{G}_{ij}+\mathcal{O}(g^4)\right]. \tag{2.34}$$

Here, $\mathbf{F}$ the matrix of combinatorial factors at free level; and $\mathbf{G}$, of order g^2 correction combinatorial factors. Note that there can be no corrections proportional to A or B, as was argued in Section 2.5. Both $\mathbf{F}$ and $\mathbf{G}$ are matrices of pure numbers; they are still functions of N, but coordinate and g^2 dependence are all absorbed in $\tilde{B}$ and $[G(x,y)]^{(2p+q)}$. Now the problem becomes one of linear algebra: starting with a basis of $\mathcal{O}_i^{[p,q,p]}$, we want to find their linear combinations $\mathcal{Y}_j^{[p,q,p]}$ that are pure operators. The $\mathcal{Y}_j^{[p,q,p]}$ have a well defined renormalized scaling dimension $\Delta_j = \Delta^0 + \Delta_j^1 + \mathcal{O}(g^4)$; $\Delta^0 = 2p+q$ for all $\mathcal{O}_i^{[p,q,p]}$ and hence for all $\mathcal{Y}_j^{[p,q,p]}$. Such operators can be chosen orthogonal at Born level, and so

$$\langle \mathcal{Y}_i^{[p,q,p]}\bar{\mathcal{Y}}_j^{[p,q,p]}\rangle = \frac{C_i^{[p,q,p];0}\delta_{ij}}{x^{2\Delta^0}}\left[1+\beta_i-\Delta_i^1\log\mu^2x^2+\mathcal{O}(g^4)\right] \tag{2.35}$$

to order g^2. Coefficients $\Delta_j^1 \sim \beta_j \sim g^2$ correspond to corrections of $\mathcal{Y}_j^{[p,q,p]}$'s scaling dimension and its normalization; β_j depends on the renormalization scale μ. To distinguish the pure operators which do receive corrections to their scaling dimension, we will denote them by $\tilde{\mathcal{Y}}$, and reserve the notation $\mathcal{Y}$ for the ones that have $\mathcal{O}(g^2)$ protected two point functions.

---

[20] The operators we are working with are after the projection onto $SU(3)\times U(1)$, as discussed in Sections 2.1 and 2.4.1. The $\mathcal{O}_i^{[p,q,p]}$ are made of only z-s and no $\bar{z}$-s.

This is a standard problem, analogous to finding the normal modes of small oscillations of a mechanical system (see [56], for example). We have to diagonalize a symmetric matrix $\mathbf{G}$ of corrections with respect to the symmetric positive definite matrix $\mathbf{F}$ of free correlators. In other words, we need to find the eigenvalues of matrix $\mathbf{F}^{-1}\mathbf{G}$. If some of them vanish, the corresponding eigenvectors are operators whose two point functions (with themselves as well as with other operators) do not get order g^2 corrections. We conjecture that these are in fact the $\frac{1}{4}$-BPS operators we are after.

We will now illustrate how this method works with the example of the [3,1,3] representation of $SU(4)$. The scalar composites corresponding to the highest weight of the [3,1,3] = **960** of $SU(4)$ are the following five linearly independent[21] operators:

$$\mathcal{O}_0^{[3,1,3]} \equiv \mathrm{tr}\, z_1z_1z_1z_2z_1z_2z_2 - \mathrm{tr}\, z_1z_1z_1z_2z_2z_1z_2 \tag{2.36}$$

$$\begin{aligned}\mathcal{O}_1^{[3,1,3]} \equiv\; & \tfrac{1}{3}\mathrm{tr}\, z_1z_1z_1z_1z_2z_2z_2 - \tfrac{1}{2}\mathrm{tr}\, z_1z_1z_1z_2z_1z_2z_2 - \tfrac{1}{2}\mathrm{tr}\, z_1z_1z_1z_2z_2z_1z_2 \\ & +\tfrac{1}{3}\mathrm{tr}\, z_1z_1z_2z_1z_2z_1z_2 + \tfrac{1}{3}\mathrm{tr}\, z_1z_1z_2z_2z_1z_1z_2\end{aligned} \tag{2.37}$$

$$\begin{aligned}\mathcal{O}_2^{[3,1,3]} \equiv\; & \mathrm{tr}\, z_1z_1z_1z_1\ \mathrm{tr}\, z_2z_2z_2 - 3\,\mathrm{tr}\, z_1z_1z_1z_2\ \mathrm{tr}\, z_1z_2z_2 \\ & + \left(2\,\mathrm{tr}\, z_1z_1z_2z_2 + \mathrm{tr}\, z_1z_2z_1z_2\right)\ \mathrm{tr}\, z_1z_1z_2 - \mathrm{tr}\, z_1z_2z_2z_2\ \mathrm{tr}\, z_1z_1z_1\end{aligned} \tag{2.38}$$

$$\begin{aligned}\mathcal{O}_3^{[3,1,3]} \equiv\; & -\left(\mathrm{tr}\, z_1z_1z_2z_2z_2 - \mathrm{tr}\, z_1z_2z_1z_2z_2\right)\ \mathrm{tr}\, z_1z_1 \\ & + \left(\mathrm{tr}\, z_1z_1z_1z_2z_2 - \mathrm{tr}\, z_1z_1z_2z_1z_2\right)\ \mathrm{tr}\, z_1z_2\end{aligned} \tag{2.39}$$

$$\begin{aligned}\mathcal{O}_4^{[3,1,3]} \equiv\; & \mathrm{tr}\, z_1z_2 \left(2\,\mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_2z_2 - \mathrm{tr}\, z_2z_2\ \mathrm{tr}\, z_1z_1z_1 - 3\mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_2z_2\right) \\ & +\mathrm{tr}\, z_1z_1 \left(\mathrm{tr}\, z_2z_2\ \mathrm{tr}\, z_1z_1z_2 + \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_2z_2z_2\right)\end{aligned} \tag{2.40}$$

The $\mathcal{O}_i^{[3,1,3]}$ are constructed by applying Young symmetrizers to all possible gauge invariant combinations of $(z_1)^4(z_2)^3$. The symmetrizers correspond to the tableau  of $SO(6)$, while gauge invariant combinations amount to grouping the z_i-s into traces. Operators resulting from other partitions and symmetrizations turn out to be linear combinations of the $\mathcal{O}_i^{[3,1,3]}$ above.

---

[21] For $N \leq 4$, the number of independent gauge invariant operators is smaller.

More explicitly, the operators listed in (2.36-2.40) are constructed as

$$7 = 7 : \qquad \mathcal{O}_0^{[3,1,3]}, \mathcal{O}_1^{[3,1,3]} \sim$$



(2.41)

$$7 = 4 + 3 : \qquad \mathcal{O}_2^{[3,1,3]} \sim$$



(2.42)

$$7 = 5 + 2 : \qquad \mathcal{O}_3^{[3,1,3]} \sim$$



(2.43)

$$7 = 3 + 2 + 2 : \qquad \mathcal{O}_4^{[3,1,3]} \sim$$



(2.44)

where each continuous group of boxes stands for a single trace. Although it does not appear in this example, in general it matters not only how we partition the string of letters, but exactly which letters we put in the groups before we apply the symmetrizers. Also, there can be more than one operator corresponding to the same partition, provided there are multiple Young symmetrisers for a given tableau ($\mathcal{O}_0^{[3,1,3]}$ and $\mathcal{O}_1^{[3,1,3]}$ here).

Here we should mentions that operator $\mathcal{O}_0^{[3,1,3]}$ has some special properties. It satisfies $(\mathcal{O}_0^{[3,1,3]})^\dagger = -(\mathcal{O}_0^{[3,1,3]})^*$ while for all other operators $(\mathcal{O}_i^{[3,1,3]})^\dagger = +(\mathcal{O}_i^{[3,1,3]})^*$. As a result, $\mathcal{O}_0^{[3,1,3]}$ has zero correlators with everything else. It is also the only pure operator out of all the $\mathcal{O}_i^{[3,1,3]}$ appearing in (2.36-2.40). It is a descendant, and we will not consider it below.

So we calculate explicitly the $\frac{1}{2} \cdot 4 \cdot (4+1) = 10$ two point functions of [3,1,3] operators, and arrange them as in Equation (2.34). We calculate the matrix $\mathbf{F}$ of free correlator combinatorial factors; and the matrix $\mathbf{G}$, of order g^2 correction combinatorial factors. The matrix $\mathbf{F}^{-1}\mathbf{G}$ is 4×4; it has two zero eigenvalues, while the other two satisfy a quadratic equation. We find

$$\frac{768}{5N^3\left(N^2-1\right)\left(N^2-4\right)}\mathbf{F}=\begin{pmatrix} \frac{N^2+3}{N^2} & \frac{12}{N} & -\frac{12}{N} & \frac{72}{N^2} \\ \frac{12}{N} & \frac{36\left(N^4-8N^2+18\right)}{N^4} & -\frac{108}{N^2} & \frac{72\left(2N^2-9\right)}{N^3} \\ \frac{-12}{N} & -\frac{108}{N^2} & \frac{36\left(N^2+6\right)}{5N^2} & -\frac{72}{N} \\ \frac{72}{N^2} & \frac{72\left(2N^2-9\right)}{N^3} & -\frac{72}{N} & \frac{72\left(N^2-3\right)}{N^2} \end{pmatrix} \tag{2.45}$$

for the matrix of free combinatorial factors, and

$$\frac{128}{25N^3\left(N^2-1\right)\left(N^2-4\right)}\mathbf{G}=\begin{pmatrix} \frac{N^2+7}{N^2} & \frac{12\left(N^2+3\right)}{N^3} & -\frac{72\left(N^2+1\right)}{5N^3} & \frac{96}{N^2} \\ \frac{12\left(N^2+3\right)}{N^3} & \frac{144}{N^2} & -\frac{144}{N^2} & \frac{864}{N^3} \\ -\frac{72\left(N^2+1\right)}{5N^3} & -\frac{144}{N^2} & \frac{144\left(N^2+16\right)}{25N^2} & -\frac{288\left(N^2+6\right)}{5N^3} \\ \frac{96}{N^2} & \frac{864}{N^3} & -\frac{288\left(N^2+6\right)}{5N^3} & \frac{576}{N^2} \end{pmatrix} \tag{2.46}$$

for the matrix of corrections proportional to $\tilde{B}(x,y)N$.

The vectors killed by $\mathbf{F}^{-1}\mathbf{G}$ work out to be

$$\mathcal{Y}_1^{[3,1,3]} = -\frac{12N}{N^2-2}\,\mathcal{O}_1^{[3,1,3]}+\mathcal{O}_2^{[3,1,3]}-\frac{5}{N^2-2}\,\mathcal{O}_3^{[3,1,3]} \tag{2.47}$$

$$\mathcal{Y}_2^{[3,1,3]} = \frac{96}{N^2-4}\mathcal{O}_1^{[3,1,3]}-\frac{4N}{N^2-4}\mathcal{O}_2^{[3,1,3]}+\frac{10N}{N^2-4}\mathcal{O}_3^{[3,1,3]}+\mathcal{O}_4^{[3,1,3]} \tag{2.48}$$

They correspond to zero eigenvalues of $\mathbf{F}^{-1}\mathbf{G}$, and so are the candidates for $\frac{1}{4}$-BPS primaries in the [3,1,3].[22] The remaining eigenvectors of $\mathbf{F}^{-1}\mathbf{G}$

$$\tilde{\mathcal{Y}}_3^{[3,1,3]} = \mathcal{O}_1^{[3,1,3]}-\frac{10}{3\left(N+\sqrt{N^2+160}\right)}\mathcal{O}_3^{[3,1,3]} \tag{2.49}$$

$$\tilde{\mathcal{Y}}_4^{[3,1,3]} = \mathcal{O}_3^{[3,1,3]}-\frac{3\left(N-\sqrt{N^2+160}\right)}{10}\mathcal{O}_1^{[3,1,3]} \tag{2.50}$$

correspond to eigenvalues $27+\frac{3\sqrt{160+N^2}}{N}$ for $\tilde{\mathcal{Y}}_3^{[3,1,3]}$, and $27-\frac{3\sqrt{160+N^2}}{N}$ for $\tilde{\mathcal{Y}}_4^{[3,1,3]}$. Expressions (2.47-2.50) are exact in N, and are not just large N approximations. As expected, the descendants $\tilde{\mathcal{Y}}$ are mixtures of operators involving commutators.

---

[22] We chose $\mathcal{Y}_2^{[3,1,3]}$ to be orthogonal to $\mathcal{Y}_1^{[3,1,3]}$, in the sense that $\langle\mathcal{Y}_2^{[3,1,3]}\bar{\mathcal{Y}}_1^{[3,1,3]}\rangle=0$.

Note that both the g^2 corrections to the scaling dimension of $\tilde{\mathcal{Y}}$ and their expansion coefficients involve radicals (so there is really no way to "guess" the pure primaries such operators came from). Also, radiative corrections to all Δ-s are non-negative, since at free field level the $\mathcal{O}_i^{[3,1,3]}$ are annihilated by a quarter of the supercharges, and hence saturate the BPS bound.

Other representations are similar and we do not give the details here. However, it gets quite cumbersome as the number of $\mathcal{O}$'s increases. The prescription of Sections 2.4.1 and 2.4.2 (find the pure non-BPS primaries, list their descendants, then subtract these pieces from the candidate $\frac{1}{4}$-BPS operator) also becomes difficult to implement in the $\mathcal{N}$=1 component approach. On the other hand, harmonic superspace gives us valuable insigths, and this more efficient approach is used in Chapter 3 to analyse these representations.

2.7 Large N analysis

As we have seen, computations get more and more cumbersome as one tries to find $\frac{1}{4}$-BPS operators for bigger representations of the color group; even the number of operators one has to consider is a nontrivial function of the representation. Symmetry factors multiplying the Feynman graphs show no immediate pattern, and most of the results presented in Sections 2.4.1, 2.4.2, and 2.6 had to be calculated using *Mathematica*.[23]

The next best thing we can do is consider the large N limit. Specifically, we shall concentrate on the leading behavior as $N \rightarrow \infty$, plus the first $1/N$ correction.

---

[23] The calculations took from 0.003 hours for the [2,0,2] representation to 23 hours for [3,1,3], per single $\mathcal{O}(g^2)$ two point function. We used a Sparc 10 with 2048 M memory and 440 MHz speed. Born level calculations were considerably (about 20 times) faster.

2.7.1 Operators $\mathcal{O}_{[p,q,p]}$ and $\mathcal{K}_{[p,q,p]}$

Let us take another look at the results of Section 2.6, where we managed to perform $\mathcal{O}(g^2)$ analysis exactly in N rather than in the large N approximation. In all cases considered so far, there is a special $\frac{1}{4}$-BPS chiral primary $\mathcal{Y}_1^{[2,3,2]}$ which is made of only the double trace and single trace operators. At large N, this operator is a combination of only a particular double trace operator, and the single trace operator, whose contribution is $1/N$ suppressed. The goal of Section 2.7 is to show that this is in fact what happens for general $[p,q,p]$ representations. Here, we begin by defining these operators.

Recall that the $SO(6)$ Young tableau for the $[p,q,p]$ of $SU(4)$ consists of two rows (one of length $p+q$, and the other of length p). Among the possible partitions of the highest weight tableau, there are two special ones

$$\mathcal{O}_{[p,q,p]} \sim \left(\begin{array}{l} \boxed{1}\boxed{\;\cdots\;}\boxed{1}\underbrace{\boxed{1}\boxed{\cdots}\boxed{1}}_{q} \\[4pt] \underbrace{\boxed{2}\boxed{\;\cdots\;}\boxed{2}}_{p} \end{array} \right), \quad \mathcal{K}_{[p,q,p]} \sim \left(\begin{array}{l} \boxed{1}\boxed{\;\cdots\;}\boxed{1}\underbrace{\boxed{1}\boxed{\cdots}\boxed{1}}_{q} \\ \underbrace{\boxed{2}\boxed{\;\cdots\;}\boxed{2}}_{p} \end{array} \right) \qquad (2.51)$$

where each continuous group of boxes stands for a single trace, as before. Explicitly, the corresponding operators are

$$\begin{aligned} \mathcal{O}_{[p,q,p]} &= \sum_{k=0}^{p} \frac{(-1)^k p!}{k!(p-k)!} \operatorname{tr}\left({z_1}^{p+q-k}{z_2}^{k}\right)_s \operatorname{tr}\left({z_1}^{k}{z_2}^{p-k}\right)_s && (2.52) \\ \mathcal{K}_{[p,q,p]} &= \sum_{k=0}^{p} \frac{(-1)^k p!}{k!(p-k)!} \operatorname{tr}\left[\left({z_1}^{p+q-k}{z_2}^{k}\right)_s \left({z_1}^{k}{z_2}^{p-k}\right)_s\right] && (2.53) \end{aligned}$$

(after projecting onto $SU(4) \to SU(3) \times U(1)$ and keeping only the highest $U(1)$-charge pieces, as discussed in Sections 2.1 and 2.4.1). Made of only z_1 and z_2, both types of operators are annihilated by four out of the sixteen Poincaré supersymmetry generators: using the SUSY transformations spelled out in Appendix 2.9.1, we find $\bar{Q}_{\bar{\zeta}} z_j = 0$, $Q_{\zeta_3} z_j = -\sqrt{2}(\lambda\zeta_3)\delta_{3j}$, so

$$\bar{Q}_{\bar{\zeta}}\mathcal{O}_{[p,q,p]} = Q_{\zeta_3}\mathcal{O}_{[p,q,p]} = \bar{Q}_{\bar{\zeta}}\mathcal{K}_{[p,q,p]} = Q_{\zeta_3}\mathcal{K}_{[p,q,p]} = 0. \qquad (2.54)$$

It is clear why $\mathcal{K}_{[p,q,p]}$ is special: it is the only single trace $[p,q,p]$ operator which can be constructed out of these fields. On the other hand, $\mathcal{O}_{[p,q,p]}$ is "the most natural" double trace composite operator in this representation. We also recognize it as the free theory chiral primary from the classification of [25].

As we have seen, neither the single trace $\mathcal{K}_{[p,q,p]}$ nor the double trace $\mathcal{O}_{[p,q,p]}$ are eigenstates of the dilation operator, for general N. Below we calculate correlators $\langle \mathcal{O}\bar{\mathcal{O}}\rangle$, $\langle \mathcal{O}\bar{\mathcal{K}}\rangle$, $\langle \mathcal{K}\bar{\mathcal{O}}\rangle$, and $\langle \mathcal{K}\bar{\mathcal{K}}\rangle$, in the large N limit, and determine the pure operators and their scaling dimension in this approximation.

2.7.2 General correlators $\langle \mathcal{O}_{[p,q,p]}\bar{\mathcal{O}}_{[p,q,p]}\rangle$ to order g^2

Let us first consider the $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle$ correlators. The free contribution is just a power of the free scalar propagator $G(x,y) = [4\pi(x-y)^2]^{-1}$, times a combinatorial factor:

$$\begin{aligned} \langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle|_{\text{free}} &= \sum_{k,l=0}^{p} \frac{(-1)^k p!}{k!(p-k)!}\,\frac{(-1)^l p!}{l!(p-l)!}\,(\mathcal{R}^{p+q,p}_{k,l})|_{\text{free}} \\ &= [G(x,y)]^{(2p+q)} \sum_{k,l=0}^{p} \frac{(-1)^k p!}{k!(p-k)!}\,\frac{(-1)^l p!}{l!(p-l)!}\,\mathcal{F}^{p+q,p}_{k,l} \end{aligned} \tag{2.55}$$

where

$$\begin{aligned} \mathcal{R}^{p+q,p}_{k,l} &= \langle \left[\operatorname{str}(z_1)^{(p+q-k)}(z_2)^k\right](x)\left[\operatorname{str}(z_1)^k(z_2)^{(p-k)}\right](x) \\ &\qquad \left[\operatorname{str}(z_1)^{(p+q-l)}(z_2)^l\right](y)\left[\operatorname{str}(z_1)^l(z_2)^{(p-l)}\right](y)\rangle, \end{aligned} \tag{2.56}$$

$$\begin{aligned} \mathcal{F}^{p+q,p}_{k,l} &= \sum_{\sigma,\rho}\left[\operatorname{str} t^{a_{k+1}}\ldots t^{a_{p+q}}t^{b_1}\ldots t^{b_k}\right]\left[\operatorname{str} t^{a_1}\ldots t^{a_k}t^{b_{k+1}}\ldots t^{b_p}\right] \\ &\qquad \left[\operatorname{str} t^{a_{\sigma(l+1)}}\ldots t^{a_{\sigma(p+q)}}t^{b_{\rho(1)}}\ldots t^{b_{\rho(l)}}\right]\left[\operatorname{str} t^{a_{\sigma(1)}}\ldots t^{a_{\sigma(l)}}t^{b_{\rho(l+1)}}\ldots t^{b_{\rho(p)}}\right] \end{aligned} \tag{2.57}$$

(σ and ρ sample over groups of permutations S_{p+q} and S_p on $p+q$ and p letters, respectively).

Like in the $\frac{1}{2}$-BPS case, the leading contribution[24] to $\mathcal{F}^{p+q,p}_{k,l} \sim (N/2)^{(2p+q)}$ comes from terms in which generators appear in reverse order for z-s and $\bar{z}$-s. To estimate the large N behavior we can use equation (2.118) to "merge traces," $(\mathrm{tr}\, t^{d_1}...t^{d_s}t^c)(\mathrm{tr}\, t^c t^{d_s}...t^{d_1}) \sim \frac{1}{2}\mathrm{tr}\, t^{d_1}...t^{d_s}t^{d_s}...t^{d_1} \sim (N/2)^{s+1}$. In order to find the numerical factor out front (which does not scale with N but depends on p and q), we should determine exactly which terms have this structure.

The generators can appear in opposite order in two pairs of traces in (2.57) under the following circumstances. First, it can happen when $k = l$ and the traces are merged as 1 with 3 and 2 with 4. The factors which arise are: $[1/p!]^2$ from symmetrizations in the 2-nd and 4-th traces; $[1/(p+q)!]^2$ from symmetrizations in the 2-nd and 4-th traces; $p!$ because for any ordering in the 1-st trace there is an identical one in the 3-d trace; $(p+q)!$ for the same reason for the 2-nd and 4-th trace; $k!(p-k)!$ since any permutation of just t^a-s or just t^b-s in the 1-st trace can be "undone" by σ-s and ρ-s in the 3-d trace; and similarly $k!(p+q-k)!$ for the 2-nd and 4-th trace; $p(p+q)$ because of trace cyclicity. There is also an overall factor from the definition (2.52). Second, if $q = 0$, we can merge traces the other way: 1 with 4 and 2 with 3; in this case $k = p - l$ and all other factors are the same. Thus, the leading contributions[25] add up to

$$\begin{aligned}
\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle|_{\text{free}} &\sim \left(\tfrac{1}{2}NG(x,y)\right)^{(2p+q)} \\
&\quad \times \sum_{k=0}^{p}(-)^k \left[\frac{p!}{k!(p-k)!}\right]^2 \frac{k!(p-k)!k!(p+q-k)!}{(p-1)!(p+q-1)!}\left[(-)^k + \delta_{q,0}(-)^{p-k}\right] \\
&= \left(\tfrac{1}{2}NG(x,y)\right)^{(2p+q)}\left[1+\delta_{q,0}(-)^p\right]\frac{p(p+q)(p+q+1)}{(q+1)}
\end{aligned} \tag{2.58}$$

The reproduces the leading order correlators in the low dimensional cases considered in Sections 2.4.1, 2.4.2, and 2.6. Also note that if $q = 0$ and p is odd, both operators $\mathcal{O}_{[p,q,p]}$ and $\mathcal{K}_{[p,q,p]}$ vanish identically, in agreement with (2.58).

---

[24] See Appendix 2.9.2 for useful $SU(N)$ identities.

[25] The error we are committing is of order $\mathcal{O}(N^{-2})$.

Now consider the corrections to this result. Diagrams contributing to two point functions of scalar composite operators at $\mathcal{O}(g^2)$ fall into two categories, see Figure 2.2. On the one hand, there are Feynman graphs involving a gauge boson exchange (these are proportional to A or B). On the other hand, we also have gauge independent ones (proportional to $\tilde{B}$) coming entirely from the $zz\bar{z}\bar{z}$-vertex. These two types of corrections have different combinatorial (index) structure, and we shall handle them separately.

2.7.2.1 Gauge dependent contributions: Combinatorial Argument

In Section 2.5, we argued that two point functions of gauge-invariant operators can not contain pieces proportional to the gauge dependent functions A and B. Here we show this explicitly for operators $\mathcal{O}_{[p,q,p]}$ and $\mathcal{K}_{[p,q,p]}$. This is the only part of Section 2.7 which is exact in N, and is not just a large N approximation.

The simplest order g^2 contribution to $\langle \mathcal{O}_{[p,q,p]} \bar{\mathcal{O}}_{[p,q,p]} \rangle$ comes from corrections to the scalar propagator (diagrams of type (a) and (b) in Figure 2.2). It has the same index structure as the free field result, and so is the same up to a factor $(p+q)NA$ for (a)- and pNA for (b)-type diagrams. These factors simply count the number of z_1-s and z_2-s.

Next consider the other diagrams where the correction comes from blocks with the same flavor in the four legs, ones of type (d) and (e). Each term in the k, l sum in (2.55) receives corrections of the form

$$\begin{aligned}(\tfrac{1}{2})(\tfrac{1}{2})(-1)(2)B \sum_{\sigma,\rho} \sum_{i\neq j=1}^{p+q} \left[\mathrm{tr}\, t^{b_{k+1}} ... t^{b_p} t^{a_1} ... t^{a_k}\right] \left[\mathrm{tr}\, t^{a_{k+1}} ... t^{a_{p+q}} t^{b_1} ... t^{b_k}\right] \\ \left[t^{b_{\rho(l+1)}} ... t^{b_{\rho(p)}} t^{a_{\sigma(1)}} ... \left[t^{a_{\sigma(i)}}, t^c\right] ... \left[t^{a_{\sigma(j)}}, t^c\right] ... t^{a_{\sigma(p+q)}} t^{b_{\rho(1)}} ... t^{b_{\rho(l)}}\right] \qquad (2.59)\end{aligned}$$

where all traces have to be symmetrized, and the second line also contains two traces.[26] The prefactor multiplying the sum comes about in the following manner: a factor of $(\frac{1}{2})$, since the sum is on $i \neq j$ rather than on $i < j$; similarly the other $(\frac{1}{2})$ arises because using $\{\sigma(i), \sigma(j)\}$ and $\{\sigma(j), \sigma(i)\}$ counts the same pair twice; a factor of 2 has to be taken into account as the two cross-symmetric pairs give the same contribution; and finally (-1) is there from two factors of i which are needed to convert f-s to commutators.

The structure in the sum (2.59) consists of four kinds of terms. The two commutators can be both in the third trace, or both in the fourth trace, or one in either trace. When both commutators are in the same trace, we can play the same game as for $\frac{1}{2}$-BPS operators. Fix i and do the sum on j first; this assembles $(...[t^{a_{\sigma(1)}}, t^c]...) + ... + (...[t^{a_{\sigma(l)}}, t^c]...) = (...[t^{a_{\sigma(1)}}...t^{a_{\sigma(l)}}, t^c]...)$ for example. Then, use trace cyclicity in the form $\mathrm{tr}\,[A, B]C = \mathrm{tr}\,A[B, C]$ to move one of the traces over to the other commutator and t^{b_ρ}-s; here we pick up a minus sign which cancels the (-1) in (2.59). As $[[t^a, t^c], t^c] = N t^a$, the first bit is easy — just like in the $\frac{1}{2}$-BPS case, it is proportional to $(+\frac{1}{2}BN)$ times the free-field combinatorial factor; when we do the sum on i we also get a factor of $(p+q)$ here. As $B + 2A = 0$ (by $\mathcal{N}$=4 SUSY), this part cancels the diagrams of type (a). The leftovers, together with the terms with commutators in different traces, add up to

$$\begin{aligned}
&\tfrac{1}{2}B \left[\mathrm{tr}\, t^{b_{k+1}}...t^{b_p} t^{a_1}...t^{a_k}\right] \left[\mathrm{tr}\, t^{a_{k+1}}...t^{a_{p+q}} t^{b_1}...t^{b_k}\right] \sum_{\sigma,\rho} \\
&\quad \left[\mathrm{tr}\, t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}} t^{a_{\sigma(1)}}...t^{a_{\sigma(l)}}\right] \left[\mathrm{tr}\, \left[t^{a_{\sigma(l+1)}}...t^{a_{\sigma(p+q)}}, t^c\right] \left[t^{b_{\rho(1)}}...t^{b_{\rho(l)}}, t^c\right]\right] \\
&+ \left[\mathrm{tr}\, \left[t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}, t^c\right] \left[t^{a_{\sigma(1)}}...t^{a_{\sigma(l)}}, t^c\right]\right] \left[\mathrm{tr}\, t^{a_{\sigma(l+1)}}...t^{a_{\sigma(p+q)}} t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right] \\
&-2 \left[\mathrm{tr}\, t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}} \left[t^{a_{\sigma(1)}}...t^{a_{\sigma(l)}}, t^c\right]\right] \left[\mathrm{tr}\, \left[t^{a_{\sigma(l+1)}}...t^{a_{\sigma(p+q)}}, t^c\right] t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right]
\end{aligned} \tag{2.60}$$

Similarly, there are diagrams of type (e) in Figure 2.2, where all of the flavors are “2” in all four legs. Here, we have a term proportional to the free-field result

---

[26] We will omit the $[G(x, y)]^{(2p+q)}$ factor which is common to all diagrams.

(the only difference being an overall factor of p rather than $p+q$), which cancels contributions from diagrams of type (b). The leftover term is the same as what we have just computed. This removes the factor of $\frac{1}{2}$ from (2.60).

Finally consider the diagrams of type (f). Now we have both flavors "1" and "2" in the four-scalar blocks, while the index structure is the same as that of (d) and (e). The discussion goes through as above, but with a few small modifications. First, the prefactor is just $(-1)B$ as now the indices i and j run over different flavors (so there is no "$i \neq j$ overcounting," no "$\{\sigma(i), \sigma(j)\}$ overcounting," and no factor of 2 from crossing symmetry). Second, we do not pick up a minus sign when transferring commutators under the traces (before, both commutators with t^c were on, say, t^a-s, whereas now one is on t^a and one on t^b). Therefore, the result of adding the diagrams of type (f) is to precisely cancel the whole leftover structure of (twice that given in) equation (2.60).

Thus we have explicitly reproduced the general result of Section 2.5, but with a lot more work: A and B contributions to two-point functions of gauge-invariant scalar composite operators combine as $2A + B$, which vanishes by $\mathcal{N}$=4 SUSY.

2.7.2.2 Contributions proportional to $\tilde{B}$

So far we have established that adding all gauge dependent Feynman graphs, i.e. diagrams of types (a), (b), (d), (e), and (f) shown in Figure 2.2, gives a vanishing $\mathcal{O}(g^2)$ contribution to the two-point functions (2.55), once we impose $\mathcal{N}$=4 SUSY. Just as in the cases of low dimensional operators considered in Sections 2.4 and 2.6, the $\mathcal{O}(g^2)$ corrections to the two-point function of $[p,q,p]$ operators come exclusively from diagrams of type (c), and are proportional to $\tilde{B}$. To find the combinatorial factor multiplying $\tilde{B}$, we need to perform a calculation similar to the one in Section 2.7.2.1.

Here, "contracted with f-s" are z-s with z-s and $\bar{z}$-s with $\bar{z}$-s (unlike in diagrams of types (d), (e) and (f), where it was one z and one $\bar{z}$), and it is more convenient to label the generators slightly differently. For example, the free-field result (2.57) can be rewritten as

$$\begin{aligned}\mathcal{F}^{p+q,p}_{k,l} \;=\; & \sum_{\sigma,\rho}\left[\operatorname{str} t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\right]\left[\operatorname{str} t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\right] \\ & \left[\operatorname{str} t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right]\left[\operatorname{str} t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\right] \qquad (2.61)\end{aligned}$$

For the Born level combinatorial factor it makes no difference, but in calculating the order g^2 diagrams of type (c) we "f-contract" i-th z_1 with $\rho(j)$-th z_2, and $\sigma(i)$-th $\bar{z}_1$ with j-th $\bar{z}_2$. This will exhaust all pairs without overcounting (because i and j are again on different flavors), so the prefactor will be just $(-1)\tilde{B}(x,y)$. Apart from this prefactor (and a factor of $[G(x,y)]^{(2p+q)}$), the (c)-type correction reads (sums on σ and ρ implied)

$$\begin{aligned}&\sum\left[\operatorname{tr} t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_k}\right]\left[\operatorname{tr} t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\right] \\ &\quad\left[\operatorname{tr} t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right]\left[\operatorname{tr} t^{a_1}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\right] \\ +&\sum\left[\operatorname{tr} t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_k}\right]\left[\operatorname{tr} t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\right] \\ &\quad\left[\operatorname{tr} t^{a_{l+1}}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right]\left[\operatorname{tr} t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\right] \\ +&\sum\left[\operatorname{tr} t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\right]\left[\operatorname{tr} t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_p}\right] \\ &\quad\left[\operatorname{tr} t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right]\left[\operatorname{tr} t^{a_1}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\right] \\ +&\sum\left[\operatorname{tr} t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\right]\left[\operatorname{tr} t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_p}\right] \\ &\quad\left[\operatorname{tr} t^{a_{l+1}}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\right]\left[\operatorname{tr} t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\right] \\ &\qquad\qquad (2.62)\end{aligned}$$

with all traces symmetrized again; the sums on $\{i,j\}$ are: in the first line, from $\{1,1\}$ to $\{l,k\}$, in the second line, from $\{l+1,1\}$ to $\{p+q,k\}$, in the third line, from $\{1,k+1\}$ to $\{l,p\}$, and in the last line, from $\{l+1,k+1\}$ to $\{p+q,p\}$.

In the large N limit, such terms can scale as $\frac{1}{2}(N/2)^{(2p+q-1)}$, at best. (Because of the commutators with t^c, we have to merge traces three times: they don't eat each other up in pairs as they did before). After including the factor of $\tilde{B}$, together with (2.58) this means that

$$\begin{aligned}\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle \;&=\; \left(\frac{N}{2}G(x,y)\right)^{(2p+q)} \qquad (2.63)\\ &\times\left([1+\delta_{q,0}(-)^p]\,\frac{p(p+q)(p+q+1)}{(q+1)} + \tilde{B}(x,y)N\times\mathcal{O}(N^{-2})\right)\end{aligned}$$

to order g^2. We might be tempted to stop here. By observing that since to working precision, the two point function of $\mathcal{O}_{[p,q,p]}$ with itself does not get $\mathcal{O}(g^2)$ corrections, we could try to conclude it is chiral, and in particular has a protected $\Delta = 2p+q$. However, as the explicit examples of Sections 2.4 and 2.6 show, $\mathcal{O}_{[p,q,p]}$ may not be a pure operator, in which case it doesn't make sense to talk about its scaling dimension.

2.7.3 General correlators $\langle\mathcal{K}_{[p,q,p]}\bar{\mathcal{O}}_{[p,q,p]}\rangle$

The analysis exactly parallels that of the previous section. Again,

$$\langle \mathcal{K}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle \;=\; \sum_{k,l=0}^{p}\frac{(-1)^k p!}{k!(p-k)!}\,\frac{(-1)^l p!}{l!(p-l)!}\,\mathcal{P}^{p+q,p}_{k,l} \qquad (2.64)$$

with

$$\begin{aligned}\mathcal{P}^{p+q,p}_{k,l} \;&=\; \langle \mathrm{tr}\,\left[\left((z_1)^{(p+q-l)}(z_2)^l\right)_s\left((z_1)^l(z_2)^{(p-l)}\right)_s\right](x)\\ &\left[\mathrm{str}\,(z_1)^{(p+q-k)}(z_2)^k\right](y)\left[\mathrm{str}\,(z_1)^k(z_2)^{(p-k)}\right](y)\rangle \qquad (2.65)\end{aligned}$$

The leading large N contributions to the free correlators now come from terms which contain the combinatorial factor similar to

$$\begin{aligned}&(\mathrm{tr}\,t^{a_1}...t^{a_k}t^{b_{k+1}}...t^{b_p}t^{a_{k+1}}...t^{a_{p+q}}t^{b_1}...t^{b_k})\;(\mathrm{tr}\,t^{b_p}...t^{b_{k+1}}t^{a_k}...t^{a_1})(\mathrm{tr}\,t^{b_k}...t^{b_1}t^{a_{p+q}}...t^{a_{k+1}})\\ &\sim (\tfrac{1}{2})^2N(N/2)^{(2p+q-2)} = \tfrac{1}{2}(N/2)^{(2p+q-1)} \qquad (2.66)\end{aligned}$$

(the two halves and -2 in the exponent are because we have to merge traces twice, and the factor of $N = \operatorname{tr}\mathbf{1}$ is there as usual). To get the other numerical factors we have to carefully analyze which terms in the sums and symmetrizations scale with N this way. So far, we do not need them.

As before, individual terms in the k, l sum get corrections similar to (2.59):

$$\begin{aligned}(\tfrac{1}{2})(\tfrac{1}{2})(-1)(2)B\sum_{\sigma,\rho}\sum_{i\neq j=1}^{p+q} \operatorname{tr}\,\Big[\Big(t^{b_{k+1}}...t^{b_p}t^{a_1}...t^{a_k}\Big)\Big(t^{a_{k+1}}...t^{a_{p+q}}t^{b_1}...t^{b_k}\Big)\Big]& \\ \Big[t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}t^{a_{\sigma(1)}}...\big[t^{a_{\sigma(i)}},t^c\big]...\big[t^{a_{\sigma(j)}},t^c\big]...t^{a_{\sigma(p+q)}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big]&\end{aligned} \tag{2.67}$$

with proper symmetrizations (and omitted factor of $[G(x,y)]^{(2p+q)}$). The only difference is that now there are three traces (rather than four). The discussion of gauge dependent diagrams (all types other than (c), see Figure 2.2) goes through verbatim, since we were only manipulating the second set of traces. As before, when we impose $\mathcal{N}$=4 SUSY they cancel, and order g^2 corrections to the two-point functions (2.64) are due to diagrams of type (c) only.

Diagrams of type (c) are only slightly different from those contributing to the $\langle\mathcal{O}\bar{\mathcal{O}}\rangle$ correlator; they add up to $(-1)\tilde{B}(x,y)[G(x,y)]^{(2p+q)}$ times

$$\begin{aligned}&\sum\operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\Big)\Big]\\
&\qquad\Big[\operatorname{tr}t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big]\Big[\operatorname{tr}t^{a_1}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big]\\
&+\sum\operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\Big)\Big]\\
&\qquad\Big[\operatorname{tr}t^{a_{l+1}}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big]\Big[\operatorname{tr}t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big]\\
&+\sum\operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_p}\Big)\Big]\\
&\qquad\Big[\operatorname{tr}t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big]\Big[\operatorname{tr}t^{a_1}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big]\\
&+\sum\operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_{j-1}}[t^{a_i},t^c]t^{b_{j+1}}...t^{b_p}\Big)\Big]\\
&\qquad\Big[\operatorname{tr}t^{a_{l+1}}...t^{a_{i-1}}[t^{b_j},t^c]t^{a_{i+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big]\Big[\operatorname{tr}t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big]\end{aligned} \tag{2.68}$$

with proper symmetrizations; the sums on $\{i, j\}$ are: in the first line, from $\{1, 1\}$ to $\{l, k\}$, in the second line, from $\{l + 1, 1\}$ to $\{p + q, k\}$, in the third line, from $\{1, k + 1\}$ to $\{l, p\}$, and in the last line, from $\{l + 1, k + 1\}$ to $\{p + q, p\}$.

We can get correlators $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{K}}_{[p,q,p]}(y)\rangle$ by just complex conjugating the sum (2.68) times the same prefactor; both the free propagator $G(x, y)$ and the function $\tilde{B}(x, y)$ are real and depend only on $(x-y)^2$, so exchanging the arguments $x \leftrightarrow y$ and conjugating doesn't change anything.

Large N dependence of (2.68) can be again estimated by merging traces. This time, we have to merge traces only twice (there are three traces total), so it scales a power of N higher than a similar $\langle \mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle|_{g^2}$ correction (where traces had to be merged three times). Hence, $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{K}}_{[p,q,p]}(y)\rangle|_{g^2} \sim (N/2)^{(2p+q)}$.

2.7.4 General correlators $\langle \mathcal{K}_{[p,q,p]}\bar{\mathcal{K}}_{[p,q,p]}\rangle$

The analysis is similar as for $\langle \mathcal{K}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle$. The only surviving contribution to $\langle \mathcal{K}_{[p,q,p]}\bar{\mathcal{K}}_{[p,q,p]}\rangle|_{g^2}$ is due to diagrams of type (c) again, and equals

$$\begin{aligned}
&\sum \operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_{j-1}}\big[t^{a_i}, t^c\big]t^{b_{j+1}}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\Big)\Big] \\
&\qquad \operatorname{tr}\Big[\Big(t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big)\Big(t^{a_1}...t^{a_{i-1}}\big[t^{b_j}, t^c\big]t^{a_{i+1}}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big)\Big] \\
&+\sum \operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_{j-1}}\big[t^{a_i}, t^c\big]t^{b_{j+1}}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_p}\Big)\Big] \\
&\qquad \operatorname{tr}\Big[\Big(t^{a_{l+1}}...t^{a_{i-1}}\big[t^{b_j}, t^c\big]t^{a_{i+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big)\Big(t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big)\Big] \\
&+\sum \operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_{j-1}}\big[t^{a_i}, t^c\big]t^{b_{j+1}}...t^{b_p}\Big)\Big] \\
&\qquad \operatorname{tr}\Big[\Big(t^{a_{l+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big)\Big(t^{a_1}...t^{a_{i-1}}\big[t^{b_j}, t^c\big]t^{a_{i+1}}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big)\Big] \\
&+\sum \operatorname{tr}\Big[\Big(t^{a_{\sigma(k+1)}}...t^{a_{\sigma(p+q)}}t^{b_1}...t^{b_k}\Big)\Big(t^{a_{\sigma(1)}}...t^{a_{\sigma(k)}}t^{b_{k+1}}...t^{b_{j-1}}\big[t^{a_i}, t^c\big]t^{b_{j+1}}...t^{b_p}\Big)\Big] \\
&\qquad \operatorname{tr}\Big[\Big(t^{a_{l+1}}...t^{a_{i-1}}\big[t^{b_j}, t^c\big]t^{a_{i+1}}...t^{a_{p+q}}t^{b_{\rho(1)}}...t^{b_{\rho(l)}}\Big)\Big(t^{a_1}...t^{a_l}t^{b_{\rho(l+1)}}...t^{b_{\rho(p)}}\Big)\Big]
\end{aligned} \tag{2.69}$$

up to a factor of $(-1)\tilde{B}(x,y)[G(x,y)]^{(2p+q)}$, and proper symmetrizations. The sums on $\{i,j\}$ are as before: in the first line, from $\{1,1\}$ to $\{l,k\}$, in the second line, from $\{l+1,1\}$ to $\{p+q,k\}$, in the third line, from $\{1,k+1\}$ to $\{l,p\}$, and in the last line, from $\{l+1,k+1\}$ to $\{p+q,p\}$).

At Born level, $\langle \mathcal{K}_{[p,q,p]}\bar{\mathcal{K}}_{[p,q,p]}\rangle|_{\text{free}} \sim (\frac{1}{2}N)^{(2p+q)}$ at large N: we have to merge traces once, and there are $2p+q$ generators involved.

2.7.5 Quarter BPS operators

Given the leading large N dependence of a $\langle \mathcal{O}\bar{\mathcal{K}}\rangle|_{\text{free}}$ correlator, it's easy to determine the leading large N dependence of the corresponding $\langle \mathcal{K}\bar{\mathcal{K}}\rangle|_{\text{free}}$. Indeed, suppose that a particular term e.g.

$$\begin{aligned}&(\operatorname{tr} t^a...t^a t^b...t^b)(\operatorname{tr} t^a...t^a t^b...t^b)\operatorname{tr}(t^a...t^a t^b...t^b)(t^a...t^a t^b...t^b) \sim \\ &(2/N)(\operatorname{tr} t^a...t^a t^b...t^b)(\operatorname{tr} t^c t^c t^a...t^a t^b...t^b)\operatorname{tr}(t^a...t^a t^b...t^b)(t^a...t^a t^b...t^b) \qquad (2.70)\end{aligned}$$

contributes to $\langle \mathcal{O}\bar{\mathcal{K}}\rangle$. We can insert contents of the first and second traces into the third trace (using $2(\operatorname{tr} At^r)(\operatorname{tr} Bt^r) \sim \operatorname{tr} AB$) to reduce this expression to a single trace. On the other hand, the term in the $\langle \mathcal{K}\bar{\mathcal{K}}\rangle$ with the same order of generators can be written as

$$\begin{aligned}&\operatorname{tr}(t^a...t^a t^b...t^b)(t^a...t^a t^b...t^b)\operatorname{tr}(t^a...t^a t^b...t^b)(t^a...t^a t^b...t^b) \sim \\ &2(\operatorname{tr} t^a...t^a t^b...t^b t^c)(\operatorname{tr} t^c t^a...t^a t^b...t^b)\operatorname{tr}(t^a...t^a t^b...t^b)(t^a...t^a t^b...t^b) \qquad (2.71)\end{aligned}$$

and we can insert the first and second traces into the third trace again in the same locations. This term gives a leading contribution provided all generators collapse after consecutively applying $2t^r t^r \sim N\mathbf{1}$ without having to commute generators past one another. In this case, the term in (2.70) also gives a leading contribution. The only other terms which have the same large N behavior are the ones that differ from it by cyclic permutations within the first and second traces. This gives an overall factor of $p(p+q)$. Comparing (2.70) and (2.71) then shows that to leading order in N, the difference is a factor of $\beta \equiv p(p+q)/N$.

For large N, the analysis of order g^2 corrections is analogous to the case of free field contributions. The structure of terms in (2.68) and (2.69) is similar, and leading contributions come from terms with the same order of generators (modulo cyclic permutations for $\langle \mathcal{O}\bar{\mathcal{K}}\rangle$ corrections); the difference is the multiplicative factor β, the same for all such terms.

Bringing this together with the results of Sections 2.7.2-2.7.4, we can write down the large N leading order two point functions as

$$\begin{pmatrix} \langle \mathcal{K}\bar{\mathcal{K}}\rangle & \langle \mathcal{K}\bar{\mathcal{O}}\rangle \\ \langle \mathcal{O}\bar{\mathcal{K}}\rangle & \langle \mathcal{O}\bar{\mathcal{O}}\rangle \end{pmatrix} = \alpha(GN)^{2p+q}\left\{ \begin{pmatrix} 1 & \beta \\ \beta & * \end{pmatrix} + \tilde{\alpha}(\tilde{B}N)\begin{pmatrix} 1 & \beta \\ \beta & \mathcal{O}(N^{-2}) \end{pmatrix} \right\} \tag{2.72}$$

where α, $*$, and $\tilde{\alpha}$ are some constants of order $\mathcal{O}(N^0)$, and $\beta = p(p+q)/N$. As before, $G \equiv G(x,y)$; $\tilde{B} \equiv \tilde{B}(x,y)$; and $\langle \mathcal{O}\bar{\mathcal{O}}\rangle \equiv \langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle$, etc. Each (order one) coefficient above is valid to $\mathcal{O}(N^{-2})$, and of course (2.72) is perturbative in the coupling constant — we have neglected $\mathcal{O}(g^4)$ terms.

Diagonalizing the matrix of corrections with respect to the matrix of free correlators as in Section 2.6, we find that

$$\tilde{\mathcal{Y}}_{[p,q,p]} = \mathcal{K}_{[p,q,p]} + \mathcal{O}(N^{-2}) \tag{2.73}$$

$$\mathcal{Y}_{[p,q,p]} = \mathcal{O}_{[p,q,p]} - \frac{p(p+q)}{N}\mathcal{K}_{[p,q,p]} + \mathcal{O}(N^{-2}) \tag{2.74}$$

are pure operators, and as such have well defined scaling dimension. At this order, the scaling dimension of $\tilde{\mathcal{Y}}_{[p,q,p]}$ receives an $\mathcal{O}(g^2N)$ correction proportional to the coefficient $\tilde{\alpha}$ in (2.72), while the scaling dimension $\Delta_{\mathcal{Y}} = 2p+q$ of $\mathcal{Y}_{[p,q,p]}$ is protected. Finally, the normalization of $\mathcal{Y}_{[p,q,p]}$ does not get any g^2 corrections, and is given by the Born level expression

$$\begin{aligned}\langle \mathcal{Y}_{[p,q,p]}(x)\bar{\mathcal{Y}}_{[p,q,p]}(y)\rangle \;=\; & [1+\delta_{q,0}(-)^p]\,\frac{p(p+q)(p+q+1)}{(q+1)}\left[\frac{N}{8\pi(x-y)^2}\right]^{2p+q} \\ & \times\Big(1+\,\mathcal{O}(N^{-2};g^4)\Big) \qquad (2.75)\end{aligned}$$

This is found from formulae (2.58), (2.72), and (2.74). The exact expressions of Sections 2.4 and 2.6 agree with (2.74) and (2.75) in the large N limit. We conclude that to working precision, $\mathcal{Y}_{[p,q,p]}$ is a $\frac{1}{4}$-BPS chiral primary operator.

2.8 Conclusion

In this Chapter we studied local, polynomial, gauge invariant scalar composite operators in $[p,q,p]$ representations of $SU(4)$ in $d=4$, $\mathcal{N}=4$ SYM. We found that certain such operators have protected two-point functions at order g^2, with each other and with other operators. We presented ample evidence that these $\mathcal{O}(g^2)$ protected operators are $\frac{1}{4}$-BPS chiral primaries in the fully interacting theory.

These operators are not just the double trace operators from the classification of [25], but mixtures of all gauge invariant local composite operators made of the same scalars: single trace operators, other double trace operators, triple trace operators, etc. As our exact in N, explicit construction of $\frac{1}{4}$-BPS primaries of low scaling dimension ($\Delta = 2p+q < 8$) shows, the N dependence of the coefficients in these linear combinations is quite complicated.

Apart from operators of low dimensions for arbitrary N, we considered the large N behavior of two point functions for all $[p,q,p]$. A leading and subleading analysis reveals that for every $[p,q,p]$, there is a $\frac{1}{4}$-BPS operator which is a certain linear combinations of double and single trace operators. We give closed form expressions for the operators involved in this linear combination and the coefficients with which they enter, all valid to next-to-leading order in N.

2.9 Appendix

2.9.1 $\mathcal{N}$= 4 SUSY in various forms

In the $\mathcal{N}$=1 component notation, the classical Lagrangian (see [57], p. 158) takes the form (in geometric notation, i.e. with $\frac{1}{g^2}$ multiplying the whole action)

$$\begin{aligned}
\mathcal{L} &= \tfrac{1}{g^2}\ \mathrm{tr}\ \Big\{ -\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + \tfrac{i}{2}\bar{\lambda}\gamma^{\mu}D_{\mu}\lambda + \tfrac{1}{2}D^2 \qquad (2.76)\\
&\quad +\tfrac{1}{2}D_{\mu}A_jD^{\mu}A_j + \tfrac{1}{2}D_{\mu}B_jD^{\mu}B_j + \tfrac{i}{2}\bar{\psi}_j\gamma^{\mu}D_{\mu}\psi_j + \tfrac{1}{2}F_jF_j + \tfrac{1}{2}G_jG_j\\
&\quad -i[A_j,B_j]D - i\bar{\psi}_j[\lambda,A_j] - i\bar{\psi}_j\gamma_5[\lambda,B_j]\\
&\quad -\tfrac{i}{2}\epsilon_{jkl}\,(\bar{\psi}_j[\psi_k,A_l] - \bar{\psi}_j\gamma_5[\psi_k,B_l]\\
&\qquad\quad +[A_j,A_k]F_l - [B_j,B_k]F_l + 2[A_j,B_k]G_l)\}
\end{aligned}$$

(with Lorentz signature); there should be no confusion between the auxiliary field D of the vector multiplet and the covariant derivative $D_{\mu} = \partial_{\mu} + iA_{\mu}$.

This Lagrangian can also be rewritten in a manifestly $SU(4)$-invariant form. Combining the three chiral spinors and the gaugino into

$$\lambda_4 = \lambda; \qquad \lambda_j = \psi_j, \quad j = 1,2,3 \qquad (2.77)$$

and making 4×4 antisymmetric matrices of scalars and pseudoscalars by

$$A_{jk} = -\epsilon_{jkl}A_l; \qquad A_{4j} = -A_{j4} = A_j; \qquad (2.78)$$

$$B_{jk} = \ \ \epsilon_{jkl}B_l; \qquad B_{4j} = -B_{j4} = B_j \qquad (2.79)$$

the Lagrangian becomes (sums on indices j,k,l now run from 1 to 4)

$$\begin{aligned}
\mathcal{L} &= \tfrac{1}{g^2}\ \mathrm{tr}\ \Big\{ -\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + \tfrac{i}{2}\bar{\lambda}_j\gamma^{\mu}D_{\mu}\lambda_j + \tfrac{1}{2}D_{\mu}A_{jk}D^{\mu}A_{jk} + \tfrac{1}{2}D_{\mu}B_{jk}D^{\mu}B_{jk}\\
&\quad +\tfrac{i}{2}\bar{\lambda}_j[\lambda_k,A_{jk}] + \tfrac{i}{2}\bar{\lambda}_j\gamma_5[\lambda_k,B_{jk}] + \tfrac{1}{32}[A_{jk},B_{lm}][A_{jk},B_{lm}]\\
&\quad +\tfrac{1}{64}[A_{jk},A_{lm}][A_{jk},A_{lm}] + \tfrac{1}{64}[B_{jk},B_{lm}][B_{jk},B_{lm}]\Big\} \qquad (2.80)
\end{aligned}$$

after integrating out the auxiliary fields D, F_j, and G_j. The A_{jk} and B_{jk} are self-dual and antiself-dual tensors of $O(4)$:

$$A_{jk} = \tfrac{1}{2}\epsilon_{jklm}A_{lm}; \quad B_{jk} = -\tfrac{1}{2}\epsilon_{jklm}B_{lm} \tag{2.81}$$

Alternatively, the fields A_i and B_i form a **6** of the R-symmetry group $SU(4) \sim SO(6)$: we can group them as $\phi^i = A_i$, $\phi^{i+3} = B_i$, $i = 1, 2, 3$.

This form of the Lagrangian is only manifestly $O(4)$ symmetric, however. If we define a complex matrix of scalars

$$M_{jk} \equiv \tfrac{1}{2}\left(A_{lm} + iB_{jk}\right) \tag{2.82}$$

subject to a reality condition

$$\bar{M}^{jk} \equiv (M_{jk})^\dagger = \tfrac{1}{2}\epsilon^{jklm}M_{lm} \tag{2.83}$$

the $\mathcal{N}$=4 Lagrangian

$$\begin{aligned}\mathcal{L} \;=\; & \tfrac{1}{g^2}\,\mathrm{tr}\,\Big\{-\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\lambda_j\sigma^\mu D_\mu\bar{\lambda}^j + \tfrac{1}{2}D_\mu M_{jk}\bar{D}^\mu\bar{M}^{jk} \\ & +i\lambda_j[\lambda_k, \bar{M}^{jk}] + i\bar{\lambda}^j[\bar{\lambda}^k, M_{jk}] + \tfrac{1}{4}[M_{jk}, M_{lm}][\bar{M}^{jk}, \bar{M}^{lm}]\Big\}\end{aligned} \tag{2.84}$$

is then manifestly $SU(4)$ covariant, as are the SUSY transformation laws

$$\delta A_\mu = i\zeta_j\sigma_\mu\bar{\lambda}^j - i\lambda_j\sigma_\mu\bar{\zeta}^j \tag{2.85}$$

$$\delta M_{jk} = \zeta_j\lambda_k - \zeta_k\lambda_j + \epsilon_{jklm}\bar{\zeta}^l\bar{\lambda}^m \tag{2.86}$$

$$\delta\lambda_j = -\tfrac{i}{2}\sigma^{\mu\nu}F_{\mu\nu}\zeta_j + 2i\sigma^\mu D_\mu M_{jk}\bar{\zeta}^k + 2i[M_{jk}, \bar{M}^{kl}]\zeta_l \tag{2.87}$$

(notice that now λ_j and $\bar{\lambda}^j$ are Weyl spinors; there should be no confusion: when spinors are multiplied by 2×2 σ matrices they are Weyl, and when the 4×4 γ matrices are used, they are Dirac).

We can also rewrite this Lagrangian in terms of three unconstrained (unlike the M_{jk}) complex scalar fields

$$z_j = \tfrac{1}{\sqrt{2}}\left(A_j + iB_j\right), \quad \bar{z}^j = \tfrac{1}{\sqrt{2}}\left(A_j - iB_j\right) \tag{2.88}$$

and the original fermions ψ_j and λ:

$$\begin{aligned}\mathcal{L} \;=\; \tfrac{1}{g^2}\,\mathrm{tr}\,\Big\{ & -\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\lambda\sigma^\mu D_\mu\bar{\lambda} + i\psi_j\sigma^\mu D_\mu\bar{\psi}^j + D_\mu z_j \bar{D}^\mu \bar{z}^j \qquad (2.89)\\ & +\tfrac{i}{\sqrt{2}}\lambda[\psi_j,\bar{z}^j] - \tfrac{i}{\sqrt{2}}\psi_j[\lambda,\bar{z}^j] - \tfrac{i}{\sqrt{2}}\epsilon^{jkl}\psi_j[\psi_k,z_l]\\ & +\tfrac{i}{\sqrt{2}}\bar{\lambda}[\bar{\psi}^j,z_j] - \tfrac{i}{\sqrt{2}}\bar{\psi}^j[\bar{\lambda},z_j] - \tfrac{i}{\sqrt{2}}\epsilon_{jkl}\bar{\psi}^j[\bar{\psi}^k,\bar{z}^l]\\ & +[z_j,z_k][\bar{z}^j,\bar{z}^k] - \tfrac{1}{2}[z_j,\bar{z}^j][z_k,\bar{z}^k]\Big\}\end{aligned}$$

and the SUSY transformations now are

$$\begin{aligned}\delta A_\mu &= i\zeta_j\sigma_\mu\bar{\psi}^j - i\psi_j\sigma_\mu\bar{\zeta}^j + i\zeta\sigma_\mu\bar{\lambda} - i\lambda\sigma_\mu\bar{\zeta} && (2.90)\\ \delta z_j &= \sqrt{2}\left(\zeta\psi_j - \zeta_j\lambda - \epsilon_{jkl}\bar{\zeta}^k\bar{\psi}^l\right) && (2.91)\\ \delta\lambda &= -\tfrac{i}{2}\sigma^{\mu\nu}F_{\mu\nu}\zeta + i\sqrt{2}\sigma^\mu D_\mu z_j\bar{\zeta}^j + i\epsilon^{jkl}[z_j,z_k]\zeta_l - i[z_j,\bar{z}^j]\zeta && (2.92)\\ \delta\psi_j &= -\tfrac{i}{2}\sigma^{\mu\nu}F_{\mu\nu}\zeta_j + i\sqrt{2}\epsilon_{jkl}\sigma^\mu\bar{D}_\mu\bar{z}^k\bar{\zeta}^l - i\sqrt{2}\sigma^\mu D_\mu z_j\bar{\zeta}\\ &\qquad + i\left([z_k,\bar{z}^k]\zeta_j - 2[z_j,\bar{z}^k]\zeta_k - \epsilon_{jkl}[\bar{z}^k,\bar{z}^l]\zeta\right) && (2.93)\end{aligned}$$

and their conjugates

$$\begin{aligned}\delta\bar{z}^j &= \sqrt{2}\left(\bar{\zeta}\bar{\psi}^j - \bar{\zeta}^j\bar{\lambda} - \epsilon^{jkl}\zeta_k\psi_l\right) && (2.94)\\ \delta\bar{\lambda} &= +\tfrac{i}{2}\bar{\sigma}^{\mu\nu}F_{\mu\nu}\bar{\zeta} - i\sqrt{2}\bar{\sigma}^\mu\bar{D}_\mu\bar{z}^j\zeta_j - i\epsilon_{jkl}[\bar{z}^j,\bar{z}^k]\bar{\zeta}^l + i[\bar{z}^j,z_j]\bar{\zeta} && (2.95)\\ \delta\bar{\psi}^j &= +\tfrac{i}{2}\bar{\sigma}^{\mu\nu}F_{\mu\nu}\bar{\zeta}^j - i\sqrt{2}\epsilon^{jkl}\bar{\sigma}^\mu D_\mu z_k\zeta_l + i\sqrt{2}\bar{\sigma}^\mu\bar{D}_\mu\bar{z}^j\zeta\\ &\qquad - i\left([\bar{z}^k,z_k]\bar{\zeta}^j - 2[\bar{z}^j,z_k]\bar{\zeta}^k - \epsilon^{jkl}[z_k,z_l]\bar{\zeta}\right) && (2.96)\end{aligned}$$

This way of writing the Lagrangian and SUSY transformations hides the full $SU(4)$ R-symmetry of the theory; now, only the $SU(3)\times U(1)$ subgroup of it is manifest.

2.9.2 Miscellaneous identities for $SU(N)$

We can use the following property of generators of $SU(N)$ (for $N \geq 3$) in the fundamental representation:

$$\{t^a, t^b\} = \frac{1}{N}\delta^{ab} + d^{abc}t^c \tag{2.97}$$

Together with $[t^a, t^b] = if^{abc}t^c$ (valid in any representation), we find

$$t^a t^b = \frac{1}{2N}\delta^{ab}\mathbf{1} + \frac{1}{2}\left(d^{abc} + if^{abc}\right)t^c \tag{2.98}$$

Let

$$g^{a_1 \ldots a_k} \equiv \operatorname{tr} t^{a_1} \ldots t^{a_k}, \quad g^{(0)} = \operatorname{tr} \mathbf{1} = N. \tag{2.99}$$

Then with the standard normalization $\operatorname{tr} t^a t^b = \frac{1}{2}\delta^{ab}$ for $SU(N)$ generators in the fundamental, we can in principle recursively determine the trace of any string of generators in terms of δ^{ab}, d^{abc}, and f^{abc}:

$$g^a = \operatorname{tr} t^a = 0,\ g^{ab} = \tfrac{1}{2}\delta^{ab},\ g^{abc} = \tfrac{1}{4}\left(d^{abc} + if^{abc}\right), \quad \text{and} \tag{2.100}$$

$$g^{a_1 \ldots a_k} = \tfrac{1}{2N}\,\delta^{a_1 a_2} g^{a_3 \ldots a_k} + 2\, g^{a_1 a_2 c} g^{c a_3 \ldots a_k} \tag{2.101}$$

Now we can set up a recursion relation for

$$P_k \equiv g^{a_1 \ldots a_k} g^{a_1 \ldots a_k} \quad \text{and} \quad \tilde{P}_k \equiv g^{a_1 \ldots a_k} g^{a_k \ldots a_1} \tag{2.102}$$

(with sums on repeated $a_1, \ldots, a_k$ implied). Using $t^a t^a = \frac{N^2-1}{2N}\mathbf{1}$, we find

$$P_k = \frac{N^2-1}{4N^2} P_{k-2} + \frac{4}{N^2-1} P_3 P_{k-1} \tag{2.103}$$

and similarly

$$\tilde{P}_k = \frac{N^2-1}{4N^2} \tilde{P}_{k-2} + \frac{4}{N^2-1} \tilde{P}_3 \tilde{P}_{k-1} \tag{2.104}$$

The values of P_2, $\tilde{P}_2$, P_3 and $\tilde{P}_3$ have to be computed explicitly; they are

$$P_2 = \tilde{P}_2 = \frac{N^2-1}{4}, \quad P_3 = -\frac{N^2-1}{4N}, \quad \text{and} \quad \tilde{P}_3 = \frac{(N^2-1)(N^2-2)}{8N} \tag{2.105}$$

For large N, the leading behavior is given by

$$g^{a_1 \ldots a_k} \sim \left(2^{k-3}\right) g^{a_1 a_2 c_3} g^{c_3 a_3 c_4} \ldots g^{c_{k-2} a_{k-2} c_{k-1}} g^{c_{k-1} a_{k-1} a_k} \tag{2.106}$$

and

$$P_{2k+1} \sim -\frac{Nk}{4^k}, \quad P_{2k} \sim \frac{N^2}{4^k}; \quad \tilde{P}_k \sim \frac{N^k}{2^k} \tag{2.107}$$

Dependence on N is[27] very different for P_k and $\tilde{P}_k$; in fact, taking generators in reverse order in the second trace (such as in $\tilde{P}_k$) grows the fastest with k, and taking them in the same order (as in P_k), the slowest.

Here are a few more identities we may have a need for in calculating two-point functions. First, the normalizations of $SU(N)$ generators in an arbitrary representation is defined in terms of a constant $C(r)$ as

$$\mathrm{tr}_r \, T_r^a T_r^b = C(r) \delta^{ab} \tag{2.108}$$

and there is a quadratic Casimir,

$$T_r^c T_r^c = C_2(r) \, \mathbf{1} \tag{2.109}$$

For the adjoint and fundamental representations, $C_2(\text{adj}) = N$, $C_2(\text{fund}) = (\frac{N^2-1}{2N})$, $C(\text{adj}) = N$, $C(\text{fund}) = \frac{1}{2}$. Then, for example,

$$\begin{aligned}
T^a T^a T^b T^b &= \left[C_2(r)\right]^2 \, \mathbf{1} && (2.110)\\
T^a T^b T^b T^a &= \left[C_2(r)\right]^2 \, \mathbf{1} && (2.111)\\
\left[T^a, T^b\right]\left[T^a, T^b\right] &= -N C_2(r) \, \mathbf{1} && (2.112)\\
T^a T^b T^a T^b &= C_2(r) \left(C_2(r) - \tfrac{N}{2}\right) \, \mathbf{1} && (2.113)
\end{aligned}$$

---

[27] Note that the way the recursion formulae (2.102) work out together with the initial values (2.105), leading order large N results are accurate to order $\mathcal{O}(N^{-2})$ and not to $\mathcal{O}(N^{-1})$, as one could have thought naively.

(we have omitted the label "r" on the generators, e.g. $T^a = T^a_r$). Longer expressions are just a little more complicated but not by much:

$$\begin{aligned}
\mathrm{tr}\, T^b \left[T^a, T^b\right] T^c \left[T^a, T^c\right] &= \tfrac{1}{4} N^2 (N^2-1) C(r) && (2.114)\\
\mathrm{tr}\, T^a T^b T^c T^a T^c T^b &= (N^2-1) C(r) (C_2(r) - \tfrac{N}{2})^2 && (2.115)\\
\mathrm{tr}\, T^a T^b T^c T^a T^b T^c &= (N^2-1) C(r) (C_2(r) - \tfrac{N}{2})(C_2(r) - N) && (2.116)
\end{aligned}$$

In particular, the last expression vanishes in the adjoint representation.

Using the fact that $U(N)_C = Gl(N, C)$, any $N \times N$ matrix A can be decomposed into generators (in the fundamental) of $SU(N)$ plus the unit matrix:

$$A = (2\mathrm{tr}\, A t^c)\, t^c + \left(\tfrac{1}{N} \mathrm{tr}\, A\right) \mathbf{1} \qquad (2.117)$$

Then, for example, we can write down the "trace merging formula"

$$2\, (\mathrm{tr}\, A t^c)\, (\mathrm{tr}\, B t^c) = \mathrm{tr}\, AB - \tfrac{1}{N}\, (\mathrm{tr}\, A)\, (\mathrm{tr}\, B) \qquad (2.118)$$

and we can arrive at an even simpler recursion relations for $\tilde{P}_k$:

$$\begin{aligned}
\tilde{P}_{k+1} &= (\mathrm{tr}\, t^{a_1} ... t^{a_k} t^c)\, (\mathrm{tr}\, t^{a_k} ... t^{a_1} t^c)\\
&= \tfrac{1}{2}\, (\mathrm{tr}\, t^{a_1} ... t^{a_k} t^{a_k} ... t^{a_1}) - \tfrac{1}{2N}\, (\mathrm{tr}\, t^{a_1} ... t^{a_k})\, (\mathrm{tr}\, t^{a_k} ... t^{a_1})\\
&= \tfrac{N}{2} (\tfrac{N^2-1}{2N})^k - \tfrac{1}{2N} \tilde{P}_k && (2.119)
\end{aligned}$$

with $\tilde{P}_1 = 0$. (Naturally, this gives the same values for $\tilde{P}_k$ as before.)

Another useful relation satisfied by the generators of $SU(N)$ in the fundamental, is

$$(t^a)_{ij} (t^a)_{kl} = \tfrac{1}{2} \left(\delta_{il} \delta_{jk} - \tfrac{1}{N} \delta_{ij} \delta_{kl}\right). \qquad (2.120)$$

Using this identity, one can easily reproduce the trace merging formula (2.118), as well as the expressions (2.105) for P_2, $\tilde{P}_2$, P_3 and $\tilde{P}_3$.

CHAPTER 3

Systematics of Quarter BPS operators in $\mathcal{N} = 4$ SYM

In this Chapter we use the machinery of (4,1,1) harmonic superspace to describe 1/4 BPS operators. The use of extended supersymmetry dramatically simplifies the counting and construction of scalar composite operators in the $[q, p, q]$ representations of the R-symmetry group $SU(4)$.

The construction of all 1/4 BPS operators in the fully interacting quantum theory is carried out as follows. First, in the classical interacting theory, a basis is produced of all the scalar operators in the representations of the R-symmetry group $SU(4)$ suitable for 1/4 BPS operators with Dynkin labels $[q, p, q]$, $q \geq 1$ and with classical dimension $p + 2q$. In the classical interacting theory there are natural candidates for 1/4 BPS operators as well as other operators which can be identified as descendants. The basis may be regrouped into operators of these two types. Harmonic superspace techniques reduce this construction down to elementary group theory. The (4,1,1) superspace notation also makes distinguishing candidate 1/4 BPS operators from descendant operators simple.

After subtracting off all the descendant pieces we will recover protected operators. These 1/4 BPS primaries are the same protected operators as the ones found in Chapter 2, but the method described here is more elegant and efficient.

3.1 $\mathcal{N} = 4$ SYM in harmonic superspace

The leading component fields of the 1/4 BPS multiplets in $\mathcal{N} = 4$ SCFT are given by scalar fields which transform under the internal symmetry group $SU(4)$ in representations with Dynkin labels of the form $[q, p, q]$. The complete supermultiplets can be very simply described in harmonic superspace, and we briefly recall how this construction works.

For $\mathcal{N}$ extended supersymmetry in four dimensions, $(\mathcal{N}, \mathrm{p}, \mathrm{q})$ harmonic superspace is obtained from ordinary Minkowski superspace M by the adjunction of a compact manifold of the form $K \equiv H \backslash SU(\mathcal{N})$ where $H = S(U(\mathrm{p}) \times U(\mathcal{N} - (\mathrm{p} + \mathrm{q})) \times U(\mathrm{q}))$. This construction allows one to construct p projections of the $\mathcal{N}$ supercovariant derivatives $D_{\alpha i}$ and q projections of their conjugates $\bar{D}^i_{\dot{\alpha}}$ which mutually anticommute. We can therefore define generalized chiral or G-analytic superfields (G for Grassmann) in such superspaces which are annihilated by these derivatives. Now the superfields in harmonic superspace will also depend on the coordinates of K, and as this space is a complex manifold, they can be analytic in the usual sense (H-analytic) in their dependence on these coordinates. As the internal manifold is compact H-analytic fields will have finite harmonic expansions. The Lorentz scalar superfields which are both G-analytic and H-analytic, which we shall refer to as analytic, are the fields we are interested in. They can be shown to carry short irreducible unitary representations of the superconformal group (provided that they transform under irreducible representations of H). For original papers and detailed accounts of harmonic and analytic superspaces, see for example [58, 59, 60, 61].

The $\mathcal{N}$=4 extended formulation is an on-shell formulation, so it is not appropriate for loop computations. This is just as well, since will use it only for classifying operators, which is a question of kinematics, and not dynamics.

3.1.1 (4,1,1) Harmonic Superspace

For the 1/4 BPS operators in $\mathcal{N} = 4$ the most appropriate harmonic superspace has $(\mathcal{N}, \mathrm{p}, \mathrm{q}) = (4, 1, 1)$. Since the analytic fields in this space will be annihilated by one D and one $\bar{D}$ it follows that they will only depend at most on 3/4 of the odd coordinates of M. Instead of working directly on the coset defined by the isotropy group $H = S(U(1) \times U(2) \times U(1))$ we shall follow the standard practice of working on the group $SU(4)$ which amounts to the same thing provided that all the fields have their dependence on H fixed. We denote an element of $SU(4)$ by ${u_I}^i$ and its inverse by ${(u^{-1})_i}^I$. The group H is taken to act on the capital index I which we decompose as $I = (1, r, 4),\ r \in \{2, 3\}$, while $SU(4)$ acts on the small indices $i, j, \ldots$. Using u and its inverse we can convert $SU(4)$ indices into H indices and vice versa. Thus we can define

$$\begin{aligned} D_{\alpha I} &\equiv {u_I}^i D_{\alpha i} = (D_{\alpha 1}, D_{\alpha r}, D_{\alpha 4}) \\ \bar{D}^I_{\dot{\alpha}} &\equiv \bar{D}^i_{\dot{\alpha}} {(u^{-1})_i}^I = (\bar{D}^1_{\dot{\alpha}}, \bar{D}^r_{\dot{\alpha}}, \bar{D}^4_{\dot{\alpha}}) \end{aligned} \tag{3.1}$$

Clearly, we have $\{D_{\alpha 1}, \bar{D}^4_{\dot{\beta}}\} = 0$. To differentiate in the coset space directions we use the right-invariant vector fields on $SU(4)$ which we denote by ${D_I}^J$, and which satisfy ${\bar{D}^I}_J = -{D_J}^I$ and ${D_I}^I = 0$. These derivatives obey the Lie algebra relations of $su(4)$ and act on ${u_K}^k$ by

$${D_I}^J {u_K}^k = {\delta_K}^J {u_I}^k - \frac{1}{4} {\delta_I}^J {u_K}^k \tag{3.2}$$

The basic differential operators on $SU(4)$ can be divided into three sets: the derivatives $({D_1}^1, {D_r}^s, {D_4}^4)$ correspond to the isotropy group, while the derivatives $({D_1}^r, {D_1}^4, {D_r}^4)$ can be thought of as essentially the components of the $\bar{\partial}$ operator on K and the derivatives $({D_r}^1, {D_4}^1, {D_4}^r)$ are the complex conjugates of these. Note that the derivatives $({D_1}^r, {D_1}^4, {D_r}^4)$ commute with $D_{\alpha 1}$ and $\bar{D}^4_{\dot{\alpha}}$. G-analytic fields are annihilated by $D_{\alpha 1}$ and $\bar{D}^4_{\dot{\alpha}}$, H-analytic fields are annihilated by $(D_1{}^r, D_1{}^4, D_r{}^4)$ and analytic fields are annihilated by both of these sets of operators.

The $\mathcal{N} = 4$ Yang-Mills theory is described in Minkowski superspace by a scalar superfield $W_{ij} = -W_{ji}$ which transforms under the six-dimensional representation of $SU(4)$ and also under the adjoint representation of the gauge group which we take to be $SU(N)$. It is real in the sense that $\bar{W}^{ij} = \frac{1}{2}\epsilon^{ijkl}W_{kl}$. This superfield satisfies the constraints

$$\begin{aligned} \nabla_{\alpha i} W_{jk} &= \epsilon_{ijkl}\Lambda_\alpha{}^l \\ \bar{\nabla}^i_{\dot{\alpha}} W_{jk} &= 2\delta^i_{[j}\bar{\Lambda}_{\dot{\alpha}k]} \end{aligned} \tag{3.3}$$

where Λ is a superfield whose leading component is the spinor field of the multiplet.and where $\nabla_{\alpha i}$ is a spinorial derivative which is covariant with respect to the gauge group. Using the superspace Bianchi identities one can show that the only other independent spacetime component of W is the spacetime Yang-Mills field strength and that all of the component fields satisfy their equations of motion.

In $(4, 1, 1)$ superspace we can define the superfield $W_{1r} \equiv u_1{}^i u_r{}^j W_{ij}$. Using the properties outlined above one can easily show that

$$\nabla_{\alpha 1} W_{1r} = \bar{\nabla}^4_{\dot{\alpha}} W_{1r} = 0 \tag{3.4}$$

and that the derivatives $(D_1{}^r, D_1{}^4, D_r{}^4)$ all annihilate W_{1r}, so that W_{1r} is a covariantly analytic field on $(4, 1, 1)$ harmonic superspace. However, if we consider gauge-invariant products of W's, i.e. traces or multi-traces, the resulting objects will be analytic superfields; they will be annihilated by $D_{\alpha 1}$ and $\bar{D}^4_{\dot{\alpha}}$ rather than the gauge-covariant versions. These are the superfields which we shall use to construct the 1/4 BPS states. To make the formulas less cluttered we shall abbreviate[1] W_{1r} to W_{r-1} and define $W^r \equiv \epsilon^{rs}W_s$.

---

[1] So the index r takes the values $r = 2, 3$ for W_{1r}, and $r = 1, 2$ for W_r.

3.1.2 Quarter BPS Operators

These superfields are easy to describe. The superfield corresponding to the representation $[q, p, q]$ contains $p + 2q$ powers of W in the representation p of $SU(2)$ (this is the $SU(2)$ in the isotropy group), i.e. it has p symmetrized $SU(2)$ indices. If $q = 0$ the single trace operators are the chiral primaries which are 1/2 BPS. These operators we will refer to as CPOs and denote by A_p,

$$A_{r_1 \ldots r_p} \equiv \mathrm{tr}\,(W_{(r_1} \ldots W_{r_p)}) \tag{3.5}$$

The lowest CPO is the stress-tensor multiplet $T_{rs} = A_{rs}$. We can obtain further 1/2 BPS operators by taking products of CPOs and symmetrizing on all of the $SU(2)$ indices. The 1/4 BPS operators (for $q > 0$) fall into two classes. There are operators that can be constructed as products of the CPOs with at least one pair of contracted indices, for example

$$T_{rs}T^{rs}, \quad A_{rst}A^{st}, \quad A_{rst}A^{t}{}_{uvw}, \quad \text{and so on.} \tag{3.6}$$

These operators have no commutators in their definition, and so are the candidate 1/4 BPS operators, up to subtleties which we shall come to in due course. Operators in the other class have at least one single-trace factor in which the indices of two or more pairs of W's are contracted, as in

$$\mathrm{tr}\, W^2W^2, \quad A_{rst}\ \mathrm{tr}\, W^2W^2, \quad \mathrm{tr}\, W_rW_sW^2W^2, \quad \text{etc.} \tag{3.7}$$

where

$$W^2 \equiv W_rW^r = \epsilon^{rs}W_rW_s = \tfrac{1}{2}\,\epsilon^{rs}[W_r, W_s] \tag{3.8}$$

These operators are descendants; the superspace Bianchi identities imply that

$$\epsilon^{\alpha\beta}\nabla_{\alpha i}\nabla_{\beta j}\bar{W}^{kl} = 2\delta_j{}^{[k}[W_{im}, \bar{W}^{l]m}] \tag{3.9}$$

From this formula and its conjugate one can see that W^2 can be written as

$$W^2 = (\nabla_1)^2 \bar{W}^{14} = -(\bar{\nabla}^4)^2 W_{14} \tag{3.10}$$

where $(\nabla_1)^2 \equiv \frac{1}{2}\epsilon^{\alpha\beta}\nabla_{\alpha 1}\nabla_{\beta 1}$, $(\bar{\nabla}^4)^2 \equiv -\frac{1}{2}\epsilon^{\dot{\alpha}\dot{\beta}}\bar{\nabla}^4_{\dot{\alpha}}\bar{\nabla}^4_{\dot{\beta}}$, and where $W_{14} = u_1{}^i u_4{}^j W_{ij}$. Given a product of W_r's containing a factor of W^2, therefore, the latter can be written in terms of derivatives as above and the derivatives can be taken to act on the whole expression with W^2 replaced by either W_{14} or its conjugate. This follows by G-analyticity. Indeed, if there are two factors of W^2 in an operator then all four derivatives can be brought outside. This is because

$$\nabla_{\alpha 1} W_{14} = 0 \qquad \text{and} \qquad \bar{\nabla}^4_{\dot{\alpha}} \bar{W}^{14} = 0 \tag{3.11}$$

In fact, the descendant 1/4 BPS operators always have at least two factors of W^2 so that they can be written explicitly as derivatives of long operators by these means. However, we are not quite finished yet because the ancestor operators will not be H-analytic on $(4,1,1)$ harmonic superspace as they stand. This can be remedied by noting that

$$(\nabla_1)^2(\bar{\nabla}^4)^2 \bar{W}^{1r} = [W^2, W^r] \tag{3.12}$$

with the aid of which we can write, for example, $\mathrm{tr}\,(W^2 W^2)$ as

$$\begin{aligned} \mathrm{tr}\,(W^2 W^2) &= -\frac{1}{3}(\nabla_1)^2(\bar{\nabla}^4)^2 \left(\mathrm{tr}\,(W_{14}\bar{W}^{14}) + \mathrm{tr}\,(W_{1r}\bar{W}^{1r})\right) \\ &= -\frac{1}{12}(\nabla_1)^2(\bar{\nabla}^4)^2 \mathrm{tr}\,(W_{ij}\bar{W}^{ij}) \\ &= -\frac{1}{12}(D_1)^2(\bar{D}^4)^2 \mathrm{tr}\,(W_{ij}\bar{W}^{ij}) \end{aligned} \tag{3.13}$$

with $D_{\alpha 1}D_{\beta 1} = \epsilon_{\alpha\beta}(D_1)^2$, and $\bar{D}^4_{\dot{\alpha}}\bar{D}^4_{\dot{\beta}} = -\epsilon_{\dot{\alpha}\dot{\beta}}(\bar{D}^4)^2$. Hence we see that $\mathrm{tr}\,(W^2 W^2)$ is a descendant of the Konishi operator $K \equiv \mathrm{tr}\,(W_{ij}\bar{W}^{ij})$. In general, one can use (3.4), (3.10), (3.11) and (3.12) to find

$$\frac{1}{2}(\nabla_1)^2(\bar{\nabla}^4)^2\, W_{ij} A \bar{W}^{ij} = W_r A [W^2, W^r] + [W^2, W^r] A W_r - 2W^2 A W^2 \tag{3.14}$$

provided A involves only the W_r. And since we are dealing with gauge invariant operators, we can replace the ∇ by D. We note for future use that each of the descendants that we consider below can be written as an ancestor superfield acted on by the differential operator $(D_1)^2(\bar{D}^4)^2$.

On the other hand, the CPOs themselves cannot be obtained by differentiation from other operators and so the candidate 1/4 BPS operators cannot be (entirely) descendants. An operator annihilated by D_1 and $\bar{D}^4$ can be either a $(D_1)^2(\bar{D}^4)^2$ descendant of a long primary; or a $(D_1)^2$ or $(\bar{D}^4)^2$ descendant of a 1/8 BPS primary; or a 1/4 BPS primary. In Chapter 2 we saw that a $[q,p,q]$ scalar composite operator can not be a descendant of a 1/8 BPS primary. Therefore, we argue that after subtracting off all the descendant pieces from candidate BPS operators, we should be left with a 1/4 BPS primary; it simply can not be anything else!

3.1.3 Examples of systematic description

We begin by outlining a few rules that determine which tensor structures are permitted. First we observe that, since contractions are made using the antisymmetric tensor ϵ^{rs} while the tensors $A_{rs\ldots t}$ are symmetric, contractions within the same A give zero,

$$A^r{}_{rs\ldots t} = 0, \tag{3.15}$$

so we can only contract indices in different A's.

Next, consider T^2. Since T_{rs} transforms under the 3-dimensional representation of $SU(2)$ it follows that the product of two Ts will decompose into the five and one-dimensional representations. The former corresponds to symmetrization on all four indices, i.e. $T_{(rs}T_{uv)}$, while for a single contraction we have

$$T_{rt}T_s{}^t = -T_{st}T_r{}^t = \tfrac{1}{2}\epsilon_{rs}T_{uv}T^{uv} \tag{3.16}$$

Similarly, the product of three Ts contains only the seven-and three-dimensional representations, so that, for example

$$T_r{}^sT_s{}^tT_t{}^r = 0 \tag{3.17}$$

One can look at contractions of other A's in a similar fashion. For example, $(A_3)^2$ contains only the seven- and three-dimensional representations of $SU(2)$ corresponding to tensors obtained by symmetrising on six or two indices with zero or two contractions respectively. So

$$A^{rst}A_{rst} = A_{(rs}{}^vA_{tu)v} = 0 \tag{3.18}$$

while

$$A_r{}^{tu}A_{stu} = A_s{}^{tu}A_{rtu} \tag{3.19}$$

or, equivalently,

$$3A_{rst}A^t{}_{uv} \quad = \quad \epsilon_{ru}(A_{stw}A^{tw}{}_v) + \epsilon_{rv}(A_{stw}A^{tw}{}_u) + \epsilon_{sv}(A_{rtw}A^{tw}{}_u) + \epsilon_{su}(A_{rtw}A^{tw}{}_v). \tag{3.20}$$

These equations generalise in a straightforward manner. Whenever we contract an odd number of indices in two As of the same length and symmetrize on the remaining indices, the resulting tensor vanishes.[2] If As of different length are contracted (as in $A_{rs}{}^tT_{tu}$), there is no such restriction.

We shall now discuss some explicit examples. We shall use the convention that uncontracted $SU(2)$ indices are understood to be totally symmetrized.

---

[2] In general there will be more than one nonvanishing structure. For instance, both $A_{(rs}{}^{vw}A_{tu)vw}$ and $A^{rstu}A_{rstu}$ are independent nonvanishing tensors.

The Representation $[1,p,1]$

Such operators have to have $(p+2)$ W_r's and only one contraction. There are no single trace operators in this class because

$$\mathrm{tr}\,(W_{r_1}\dots W_{r_p}W^2)=0 \tag{3.21}$$

Hence these operators can only be constructed by contracting CPOs. They are all protected. This can also be seen from representation theory because there are no long representations which contain these representations [62]. This result also shows that any single-trace factor in an operator must have at least two contractions.

Here we list the lowest dimensional examples of $[1,p,1]$ representations (for $p \leq 5$). For [1,1,1] and [1,2,1] we can not construct any nonvanishing tensors of this form. For [1,3,1], there is one possible operator,

$$\mathcal{O} \;=\; A_{rst}T^{t}{}_{u} \tag{3.22}$$

Similarly, for [1,4,1] the only possible operator is

$$\mathcal{O} \;=\; A_{rstu}T^{u}{}_{v} \tag{3.23}$$

Higher representations offer more choices, and already in the [1,5,1] we find

$$\begin{aligned}\mathcal{O}_1 &= A_{rstuv}T^{v}{}_{w}\\ \mathcal{O}_2 &= A_{rstv}A^{v}{}_{uw}\\ \mathcal{O}_3 &= T_{rs}T_{tv}A^{v}{}_{uw}\end{aligned} \tag{3.24}$$

All of these operators have protected two-point functions, as we will explicitly verify in Section 3.2.

The Representation [2,0,2]

This operator is realized as an $SU(2)$ scalar in $(4,1,1)$ harmonic superspace. There are just two possibilities

$$\begin{aligned} \mathcal{O}_1 &= T_{rs}T^{rs} \\ \mathcal{O}_2 &= \mathrm{tr}\,(W^2W^2) \end{aligned} \tag{3.25}$$

Using the rules outlined in the beginning of Section 3.1.3, we see that $\mathcal{O}_1$ is the only multiple trace operator one can construct with two pairs of contracted indices, and there is also no other choice for the single trace operator but $\mathcal{O}_2$. $\mathcal{O}_1$ is a candidate 1/4 BPS operator while $\mathcal{O}_2$ is a descendant; as seen in (3.13), it is a descendant of the Konishi operator.

The Representation [2,1,2]

This has 5 fields and forms an $SU(2)$ doublet. There are again only two possibilities

$$\begin{aligned} \mathcal{O}_1 &= A_{rst}T^{st} \\ \mathcal{O}_2 &= \mathrm{tr}\,(W_rW^2W^2) \end{aligned} \tag{3.26}$$

This case is completely parallel to the [2,0,2] representation.

The operator $\mathcal{O}_2$ can be written in the form

$$\mathcal{O}_2 = -\frac{1}{16}(D_1)^2(\bar{D}^4)^2\,\mathrm{tr}\,(W_rW_{ij}\bar{W}^{ij}) \tag{3.27}$$

Note that the ancestor here is defined on (4,1,1) harmonic superspace; it is not G-analytic but it is H-analytic. One can easily remove the harmonic variables to obtain the corresponding superfield on ordinary superspace. In this case it is $\mathrm{tr}\,(W_{ij}W_{kl}\bar{W}^{kl})$.

The Representation [2,2,2]

This has 6 fields and transforms as a triplet under $SU(2)$. Multiple trace operators are constructed in the following way. We can partition the set of six fields as 6 = 4 + 2, 6 = 3 + 3, or 6 = 2 + 2 + 2. Two pairs of indices are contracted, and two remaining indices are symmetrized. The possibilities are

$$\begin{aligned}
\mathcal{O}_1 &= A_{rstu}T^{tu} \\
\mathcal{O}_2 &= A_r{}^{tu}A_{stu} \\
\mathcal{O}_3 &= T_{rs}T_{tu}T^{tu} \\
\mathcal{O}_4 &= \mathrm{tr}\,(W_rW_sW^2W^2) \\
\mathcal{O}_5 &= \mathrm{tr}\,(W_rW^2W_sW^2) \\
\mathcal{O}_6 &= \mathrm{tr}\,(W^2W^2)T_{rs}
\end{aligned} \tag{3.28}$$

The first three are candidate 1/4 BPS operators while the last three are descendants. For the partitions 6 = 4 + 2 and 6 = 3 + 3, these are the only choices because contractions within the same A give zero. For the partition 6 = 2 + 2 + 2, equation (3.17) relates any other triple-trace [2,2,2] operator to $\mathcal{O}_3$.

The operator $\mathcal{O}_6$ is a descendant of the product of the Konishi operator and the supercurrent, while $\mathcal{O}_4$ and $\mathcal{O}_5$ are descendants of the operators

$$\begin{aligned}
\mathcal{A}_1 &= \mathrm{tr}\,(W_rW_sW_{ij}\bar{W}^{ij}) \\
\mathcal{A}_2 &= \mathrm{tr}\,(W_rW_{ij}W_s\bar{W}^{ij})
\end{aligned} \tag{3.29}$$

A short calculation yields

$$\begin{aligned}
\mathcal{O}_4 &= \frac{1}{40}(D_1)^2(\bar{D}^4)^2(\mathcal{A}_2 - 3\mathcal{A}_1) \\
\mathcal{O}_5 &= \frac{1}{20}(D_1)^2(\bar{D}^4)^2(\mathcal{A}_1 - 2\mathcal{A}_2)
\end{aligned} \tag{3.30}$$

In terms of ordinary superfields, both $\mathcal{A}_1$ and $\mathcal{A}_2$ are in the $[0,2,0]$ representation of $SU(4)$.

The Representation [2,3,2]

The possibilities are

$$\begin{aligned}
\mathcal{O}_1 &= A_{rstuv}T^{uv} \\
\mathcal{O}_2 &= A_{rs}{}^{uv}A_{tuv} \\
\mathcal{O}_3 &= A_{rst}T_{uv}T^{uv} \\
\mathcal{O}_4 &= A_r{}^{uv}(T^2)_{stuv} \\
\mathcal{O}_5 &= \mathrm{tr}\,(W_rW_sW_tW^2W^2) \\
\mathcal{O}_6 &= \mathrm{tr}\,(W_rW_sW^2W_tW^2) \\
\mathcal{O}_7 &= T_{rs}\mathrm{tr}\,(W_tW^2W^2) \\
\mathcal{O}_8 &= A_{rst}\mathrm{tr}\,(W^2W^2)
\end{aligned} \tag{3.31}$$

The first four are candidate 1/4 BPS operators while the second four are descendants.

The last of these is a descendant of a product of A_3 and the Konishi operator, while the ancestor of $\mathcal{O}_7$ is a product of T and $\mathrm{tr}\,(W_rW_{ij}\bar{W}^{ij})$. For the other two descendants we have

$$\begin{aligned}
\mathcal{O}_5 &= \frac{1}{24}(D_1)^2(\bar{D}^4)^2(\mathcal{A}_2 - 2\mathcal{A}_1) \\
\mathcal{O}_6 &= \frac{1}{24}(D_1)^2(\bar{D}^4)^2(\mathcal{A}_1 - 2\mathcal{A}_2)
\end{aligned} \tag{3.32}$$

where

$$\begin{aligned}
\mathcal{A}_1 &= \mathrm{tr}\,(W_rW_sW_tW_{ij}\bar{W}^{ij}) \\
\mathcal{A}_2 &= \mathrm{tr}\,(W_rW_sW_{ij}W_t\bar{W}^{ij})
\end{aligned} \tag{3.33}$$

The Representation [3,0,3]

There is only one possibility:

$$\mathcal{O} \;=\; \mathrm{tr}\,(W^2W^2W^2) \tag{3.34}$$

This operator is a descendant. There are no candidate 1/4 BPS operators in this case. As we saw in (3.17) and (3.18), the operators $A_{rst}A^{rst}$ and $T_{rs}T_t{}^rT^{st}$ vanish identically. Explicitly, this operator can be written as

$$\mathcal{O} = -\frac{1}{8}(D_1)^2(\bar{D}^4)^2\mathrm{tr}\,(W^2W_{ij}\bar{W}^{ij}) \tag{3.35}$$

The Representation [3,1,3]

This example again has seven fields but the representation of $SU(2)$ is the doublet. The operators are

$$\begin{aligned}
\mathcal{O}_1 &= A_r{}^{stu}A_{stu} \\
\mathcal{O}_2 &= (T^2)_{rstu}A^{stu} \\
\mathcal{O}_3 &= \mathrm{tr}\,(W_rW^2W^2W^2) \\
\mathcal{O}_4 &= \mathrm{tr}\,(W_rW^2W_sW^2W^s - W_rW_sW^2W^sW^2) \\
\mathcal{O}_5 &= T_r{}^s\mathrm{tr}\,(W_sW^2W^2)
\end{aligned} \tag{3.36}$$

so there are 3 descendants in this case. We have symmetrized $\mathcal{O}_4$ so that $\mathcal{O}_4^\dagger = +(\mathcal{O}_4)^*$. This symmetry amounts to charge conjugation on the fields X in the adjoint representation of the gauge group and the **6** of $SU(4)$. Its effect on the $\mathcal{N}=1$ superfield formulation is to map $z_j \to z_j^t$.

The last operator is again a descendant of a product of operators that we have discussed previously. For the other two we have

$$\mathcal{O}_3 \;=\; \frac{1}{30}(D_1)^2(\bar{D}^4)^2(\mathcal{A}_2 - 5\mathcal{A}_1)$$

$$\mathcal{O}_4 = \frac{1}{6}(D_1)^2(\bar{D}^4)^2(\mathcal{A}_1+\mathcal{A}_2) \tag{3.37}$$

where

$$\begin{aligned}\mathcal{A}_1 &= \mathrm{tr}\,(W_r W^2 W_{ij}\bar{W}^{ij}) \\ \mathcal{A}_2 &= \mathrm{tr}\,(W_r W_s W_{ij} W^s \bar{W}^{ij})\end{aligned} \tag{3.38}$$

3.1.4 Multiplicity of quarter BPS operators

The (classical) quarter BPS operators in the $SU(4)$ representation are built from single trace quarter (and half) BPS operators. These have the form

$$\mathrm{tr}\,(W_{(r_1}\dots W_{r_q)}(W^2)^p) \tag{3.39}$$

for operators in the $[p,q,p]$ $SU(4)$ representation, but the order of the $2p+q$ operators inside the trace is arbitrary.

To find the number of different single trace operators in this representation, N_{pq}, consider the reducible operator

$$X_Q := \mathrm{tr}\,(W_{r_1}\dots W_{r_Q}), \tag{3.40}$$

where the $SU(2)$ indices are no longer taken to be symmetrised. This is in a reducible representation of $SU(2)$, and contains all single trace scalar composite operators of dimension Q. So one obtains the number of operators in each representation by expanding this operator as a sum of irreducible representations. For example to find all single trace operators of dimension 4 consider X_4. This has 6 components (given by (1111), (1112), (1122), (1212), (2221), (2222) where $(r_1r_2r_3r_4)$ is short hand for $\mathrm{tr}\,(W_{r_1}\dots W_{r_4})$.) In terms of irreducible $SU(2)$ representations it splits as $6 = 5 + 1$. In terms of $SU(4)$ representations the 5 corresponds to $[0,4,0]$ and the 1 corresponds to $[2,0,2]$, and so we find that there is only one operator in each of these two representations.

More generally, to split X_Q into irreducibles, consider the components of X_Q. Let $c(Q,p)$ denote the number of components of X_Q with p 1's and $Q-p$ 2's, i.e. the number of ways to arrange a total of Q objects with p of one type and $Q-p$ of another type up to circular permutations.

Then X_Q splits into the following irreducible representations:

$$\sum_{p=0}^{\lfloor Q/2 \rfloor} \left(c(Q,p) - c(Q,p-1)\right)\ [p,q,p] \tag{3.41}$$

where $q = Q - 2p$ and where $\lfloor x \rfloor$ denotes the largest integer less than or equal to x. So the number of single trace operators in the $[p,q,p]$ representation is

$$N_{pq} = c(Q,p) - c(Q,p-1). \tag{3.42}$$

In general the formula for $c(Q,p)$ is quite complicated, but in certain cases it simplifies. For example

$$c(Q,0) = 1, \quad c(Q,1) = 1, \quad c(Q,2) = \lfloor Q/2 \rfloor, \tag{3.43}$$

and if Q and p are co-prime then

$$N_{pq} = \frac{1}{Q}\binom{Q}{p}. \tag{3.44}$$

As an example, consider dimension 6 operators: X_6 has 14 components and $c(6,p)$ is given by:

$$c(6,0) = c(6,1) = 1, \quad c(6,2) = 3, \quad c(6,3) = 4. \tag{3.45}$$

Then (3.42) gives

$$N_{06} = 1, \quad N_{14} = 0, \quad N_{22} = 2, \quad N_{30} = 1, \tag{3.46}$$

reproducing the correct numbers of single-trace operators discussed above (in particular there are two operators in the $[2,2,2]$ representation and 1 in the $[3,0,3]$.)

Since multiple trace operators can be obtained by multiplying together single trace operators, to find the number of multi-trace operators in a given representation one just has to consider all possible ways of obtaining the representation in question from tensor products of other representations and use the formula for single-trace operators.

3.1.5 Relationship between $\mathcal{N} = 4$ and $\mathcal{N} = 1$ superfields

The map between quarter BPS operators in the $\mathcal{N} = 1$ formalism and those in $(4, 1, 1)$ analytic superspace is straightforward. In the $\mathcal{N} = 1$ formalism the quarter BPS operators are given by

$$\left[(z^{2c})^p (z_d)^q\right] \tag{3.47}$$

where $[\ldots]$ denote gauge invariant combinations, $(X_a)^p \equiv X_{(a_1} \ldots X_{a_p)}$ and $z^{2c} \equiv z_a z_b \epsilon^{abc}$. Here the $a, b, \ldots = (1, 2, 3)$ are $SU(3)$ indices. These operators have highest weight state given by

$$[(z_1 z_2 - z_2 z_1)^p (z_1)^q] \,. \tag{3.48}$$

In $(4, 1, 1)$ harmonic superspace on the other hand, this object is given by

$$\left[(W^2)^p (W_r)^q\right] . \tag{3.49}$$

If we relabel the $SU(2)$ indices $r, s, \ldots = 1, 2$ then this operator has highest weight state

$$[(W_1 W_2 - W_2 W_1)^p (W_1)^q] \,. \tag{3.50}$$

The correspondence between the $N = 1$ operators and the harmonic superspace operators is now clear, one simply replaces W with z to obtain the highest weight states of each.

3.2 Explicit Computations

In this section we will explicitly calculate two point functions of the above operators. We will work with the lowest components of superfields, the z_i^a and $\bar{z}_i^a$. (Here, $i = 1, ..., 3$, and a labels the adjoint representation of the gauge group $SU(N)$.) We list operators in a given irrep of the R-symmetry group $SU(4)$, and for the descendant operators write out the corresponding Konishi-like long operator they come from. Then we look at Born level and order g^2 contributions to the two point functions of the highest weight state operators.

In each representation we will have the descendant operators L_i and "candidate 1/4-BPS" operators $\mathcal{O}$. We will compute the order g^0 two point functions $\langle OL_i^\dagger \rangle_{\text{Born}}$ and $\langle L_i L_j^\dagger \rangle_{\text{Born}}$. Then we will consider operators

$$\tilde{\mathcal{O}} \equiv \mathcal{O} - \langle OL_i^\dagger \rangle_{\text{Born}} \Big(\langle LL^\dagger \rangle_{\text{Born}}^{-1} \Big)^{ij} L_j \tag{3.51}$$

By construction, they are orthogonal to all the L_i at Born level, $\langle \tilde{\mathcal{O}} L_i^\dagger \rangle_{\text{Born}} = 0$. Then we will show that these operators $\tilde{\mathcal{O}}$ have protected two point functions at order g^2, $\langle \tilde{\mathcal{O}} L_i^\dagger \rangle_{g^2} = 0$ and $\langle \tilde{\mathcal{O}}' \tilde{\mathcal{O}}^\dagger \rangle_{g^2} = 0$ for all such operators $\tilde{\mathcal{O}}, \tilde{\mathcal{O}}'$. The claim is that these operators $\tilde{\mathcal{O}}$ are 1/4-BPS.

The basis of operators we will choose is slightly different from the one used in Chapter 2. The operators introduced in the preceding section are more natural and intuitive.

The representation $[1, p, 1]$

- There is only one operator in the representation $[1, 3, 1]$ whose highest $SU(4)$ weight state is

$$\mathcal{O} \;\; \equiv \;\; \text{tr}\, z_1 z_1 \, \text{tr}\, z_1 z_1 z_2 - \text{tr}\, z_1 z_2 \, \text{tr}\, z_1 z_1 z_1 \tag{3.52}$$

while acting on $\mathcal{O}$ once with an $SU(4)$ ladder operator gives

$$\mathcal{O}' \equiv 2\,\mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_2z_2 - \mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_1z_2 - \mathrm{tr}\, z_2z_2\ \mathrm{tr}\, z_1z_1z_1 \tag{3.53}$$

This operator has the same weight as the [2,1,2] operators (but is of course orthogonal to them). The Born and order g^2 overlaps are

$$\langle \mathcal{O}'\mathcal{O}'^{\dagger}\rangle_{\mathrm{Born}} = \frac{15}{32}N(N^2-1)(N^2-4), \quad \langle \mathcal{O}'\mathcal{O}'^{\dagger}\rangle_{g^2} = 0. \tag{3.54}$$

So indeed it is a 1/4-BPS operator.

- There is only one operator in the representation $[1,4,1]$. The highest $SU(4)$ weight state operator is

$$\mathcal{O} \equiv \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_1z_1z_2 - \mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_1z_1z_1 \tag{3.55}$$

while acting on $\mathcal{O}$ once with an $SU(4)$ ladder operator gives

$$\begin{aligned}\mathcal{O}' \equiv\ & 2\,\mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_1z_2z_2 + \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_2z_1z_2 \\ & -2\,\mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_1z_1z_2 - \mathrm{tr}\, z_2z_2\ \mathrm{tr}\, z_1z_1z_1z_1\end{aligned} \tag{3.56}$$

This operator has the same weight as the [2,2,2] operators (but is of course orthogonal to them). The Born and order g^2 overlaps are

$$\langle \mathcal{O}'\mathcal{O}'^{\dagger}\rangle_{\mathrm{Born}} = \frac{3}{8}(N^2-1)(N^2-4)(N^2-9), \quad \langle \mathcal{O}'\mathcal{O}'^{\dagger}\rangle_{g^2} = 0. \tag{3.57}$$

So indeed it is a 1/4-BPS operator.

- Finally, there are 3 operators in the representation $[1,5,1]$. Their highest $SU(4)$ weight state operators are

$$\mathcal{O}_1 \equiv \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_1z_1z_1z_2 - \mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_1z_1z_1z_1 \tag{3.58}$$

$$\mathcal{O}_2 \equiv \mathrm{tr}\, z_1z_1z_1\ \mathrm{tr}\, z_1z_1z_1z_2 - \mathrm{tr}\, z_1z_1z_2\ \mathrm{tr}\, z_1z_1z_1z_1 \tag{3.59}$$

$$\mathcal{O}_3 \equiv \mathrm{tr}\, z_1z_1\,(\mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_1z_2 - \mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_1z_1) \tag{3.60}$$

while acting on $\mathcal{O}$ once with an $SU(4)$ ladder operator gives

$$\begin{aligned}
\mathcal{O}'_1 &\equiv 2\,\mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_1z_1z_1z_2z_2 + 2\,\mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_1z_1z_2z_1z_2 \\
&\quad -3\,\mathrm{tr}\,z_1z_2\ \mathrm{tr}\,z_1z_1z_1z_1z_2 - \mathrm{tr}\,z_2z_2\ \mathrm{tr}\,z_1z_1z_1z_1z_1 \qquad (3.61) \\
\mathcal{O}'_2 &\equiv 2\,\mathrm{tr}\,z_1z_1z_1\ \mathrm{tr}\,z_1z_1z_2z_2 + \mathrm{tr}\,z_1z_1z_1\ \mathrm{tr}\,z_1z_2z_1z_2 \\
&\quad -\mathrm{tr}\,z_1z_1z_2\ \mathrm{tr}\,z_1z_1z_1z_2 - 2\,\mathrm{tr}\,z_1z_2z_2\ \mathrm{tr}\,z_1z_1z_1z_1 \qquad (3.62) \\
\mathcal{O}'_3 &\equiv 2\,\mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_1z_2z_2 + \mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_1z_2\ \mathrm{tr}\,z_1z_1z_2 \\
&\quad -2\,\mathrm{tr}\,z_1z_2\ \mathrm{tr}\,z_1z_2\ \mathrm{tr}\,z_1z_1z_1 - \mathrm{tr}\,z_2z_2\ \mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_1z_1z_1 \qquad (3.63)
\end{aligned}$$

These operators have the same weight as the [2,3,2] operators (but are of course orthogonal to them). The Born and order g^2 overlaps are

$$\begin{aligned}
\langle \mathcal{O}'_i \mathcal{O}'^{\dagger}_j \rangle_{\mathrm{Born}} &= \tfrac{35(N^2-1)(N^2-4)}{128N} \\
&\quad \times \begin{pmatrix} N^4-10N^2+72 & -11N^2+36 & 6N(N^2-2) \\ & N^4-4N^2+18 & -2N(2N^2+3) \\ & & 2N^2(N^2+5) \end{pmatrix}, \qquad (3.64) \\
\langle \mathcal{O}'_i \mathcal{O}'^{\dagger}_j \rangle_{g^2} &= 0. \qquad (3.65)
\end{aligned}$$

So indeed they all are 1/4-BPS operators.

The Representation [2,0,2]

The operators corresponding to (3.25) are

$$\begin{aligned}
\mathcal{O}_1 &= 2\left(\mathrm{tr}\,z_1z_1\ \mathrm{tr}\,z_2z_2 - \mathrm{tr}\,z_1z_2\ \mathrm{tr}\,z_1z_2\right) \\
\mathcal{O}_2 &= \mathrm{tr}\,z_1z_1z_2z_2 - \mathrm{tr}\,z_1z_2z_1z_2 \qquad (3.66)
\end{aligned}$$

The single trace operator $\mathcal{O}_2$ is a descendant of the Konishi scalar (see equation 2.20),

$$\mathcal{O}_2 \sim (Q^2\bar{Q}^2)\ \mathrm{tr}\,z_j\bar{z}^j. \qquad (3.67)$$

On the other hand, the operator orthogonal to $\mathcal{O}_2$ at Born level

$$\tilde{\mathcal{O}}_1 = \mathcal{O}_1 - \frac{4}{N}\mathcal{O}_2 \tag{3.68}$$

stays orthogonal to $\mathcal{O}_2$, $\langle \tilde{\mathcal{O}}_1(x)\bar{\mathcal{O}}_2(y)\rangle = 0$; and its two-point function is protected at order g^2, $\langle \tilde{\mathcal{O}}_1(x)\bar{\tilde{\mathcal{O}}}_1(y)\rangle = \langle \tilde{\mathcal{O}}_1(x)\bar{\tilde{\mathcal{O}}}_1(y)\rangle_{\text{Born}}$.

The Representation [2,1,2]

In this representation we again have only two operators

$$\begin{aligned}
\mathcal{O}_1 &= \operatorname{tr} z_1z_1z_1 \ \operatorname{tr} z_2z_2 - 2\operatorname{tr} z_1z_2 \ \operatorname{tr} z_1z_2z_1 + \operatorname{tr} z_1z_2z_2 \ \operatorname{tr} z_1z_1 \\
\mathcal{O}_2 &= \operatorname{tr} z_1z_1z_1z_2z_2 - \operatorname{tr} z_1z_1z_2z_1z_2
\end{aligned} \tag{3.69}$$

and the single trace operator is again a descendant,

$$\mathcal{O}_2 \sim (Q^2\bar{Q}^2)\ \operatorname{tr}\left[z_1z_j\bar{z}^j + z_1\bar{z}^jz_j\right] \tag{3.70}$$

The operator orthogonal to $\mathcal{O}_2$ at Born level

$$\tilde{\mathcal{O}}_1 = \mathcal{O}_1 - \frac{6}{N}\mathcal{O}_2 \tag{3.71}$$

satisfies $\langle \tilde{\mathcal{O}}_1(x)\bar{\mathcal{O}}_2(y)\rangle = 0$, $\langle \tilde{\mathcal{O}}_1(x)\bar{\tilde{\mathcal{O}}}_1(y)\rangle = \langle \tilde{\mathcal{O}}_1(x)\bar{\tilde{\mathcal{O}}}_1(y)\rangle_{\text{Born}}$ at order g^2.

The Representation [2,2,2]

Here we have a total of six operators. The lowest components of superfields (3.28) are

$$\begin{aligned}
\mathcal{O}_1 &\equiv 3\operatorname{tr} z_1z_1z_1z_1 \ \operatorname{tr} z_2z_2 - 6\operatorname{tr} z_1z_1z_1z_2 \ \operatorname{tr} z_1z_2 \\
&\qquad\qquad + \left(2\operatorname{tr} z_1z_1z_2z_2 + \operatorname{tr} z_1z_2z_1z_2\right)\ \operatorname{tr} z_1z_1 \\
\mathcal{O}_2 &\equiv \operatorname{tr} z_1z_1z_1 \ \operatorname{tr} z_1z_2z_2 - \operatorname{tr} z_1z_1z_2 \ \operatorname{tr} z_1z_1z_2
\end{aligned}$$

$$
\begin{aligned}
\mathcal{O}_3 &\equiv \operatorname{tr} z_1 z_1 \left(\operatorname{tr} z_1 z_1 \; \operatorname{tr} z_2 z_2 - \operatorname{tr} z_1 z_2 \; \operatorname{tr} z_1 z_2\right) \\
\mathcal{O}_4 &\equiv \operatorname{tr} z_1 z_1 z_1 z_1 z_2 z_2 - 2 \operatorname{tr} z_1 z_1 z_1 z_2 z_1 z_2 + \operatorname{tr} z_1 z_1 z_2 z_1 z_1 z_2 \\
\mathcal{O}_5 &\equiv \operatorname{tr} z_1 z_1 z_1 z_2 z_1 z_2 - \operatorname{tr} z_1 z_1 z_2 z_1 z_1 z_2 \\
\mathcal{O}_6 &\equiv \operatorname{tr} z_1 z_1 \left(\operatorname{tr} z_1 z_1 z_2 z_2 - \operatorname{tr} z_1 z_2 z_1 z_2\right)
\end{aligned}
\tag{3.72}
$$

(and we didn't bother to keep the same normalization factors for all of them — just whatever looks better). The descendants arise from the Konishi-like long primary operators as

$$
\begin{aligned}
\mathcal{O}_4 &\sim (Q^2 \bar{Q}^2) \operatorname{tr} \left[z_1 z_1 z_j \bar{z}^j + z_1 z_1 \bar{z}^j z_j\right] \\
\mathcal{O}_5 &\sim (Q^2 \bar{Q}^2) \operatorname{tr} \left[z_1 z_j z_1 \bar{z}^j\right] \\
\mathcal{O}_6 &\sim (Q^2 \bar{Q}^2) \; [\operatorname{tr} z_1 z_1] \left[\operatorname{tr} z_j \bar{z}^j\right]
\end{aligned}
\tag{3.73}
$$

Note that another operator exists in a long multiplet, whose descendant coincides with $\mathcal{O}_6$,

$$
\mathcal{O}_6 \sim (Q^2 \bar{Q}^2) \; [\operatorname{tr} z_1 z_j] \left[\operatorname{tr} z_1 \bar{z}^j\right]
\tag{3.74}
$$

This may be established by observing that the difference operator,

$$
3 \, [\operatorname{tr} z_1 z_j] \left[\operatorname{tr} z_1 \bar{z}^j\right] - [\operatorname{tr} z_1 z_1] \left[\operatorname{tr} z_j \bar{z}^j\right]
\tag{3.75}
$$

is semi-short[3]. A similar phenomenon occurs for higher representations, and we will not mention it explicitly.

The linear combinations orthogonal to these operators at Born level can be taken as

$$
\begin{aligned}
\tilde{\mathcal{O}}_1 &= \mathcal{O}_1 - \frac{24}{N}\mathcal{O}_4 - \frac{48(2N^2-3)}{N(3N^2-2)}\mathcal{O}_5 + \frac{40}{3N^2-2}\mathcal{O}_6 \\
\tilde{\mathcal{O}}_2 &= \mathcal{O}_2 - \frac{4}{N}\mathcal{O}_4 - \frac{3(7N^2-8)}{N(3N^2-2)}\mathcal{O}_5 + \frac{5}{3N^2-2}\mathcal{O}_6 \\
\tilde{\mathcal{O}}_3 &= \mathcal{O}_3 \qquad\qquad - \frac{20}{3N^2-2}\mathcal{O}_5 - \frac{10}{3N^2-2}\mathcal{O}_6
\end{aligned}
\tag{3.76}
$$

---

[3] This is the non-renormalised 20' operator discussed in [12, 63, 64]

The matrix of two point functions in this basis is

$$\begin{pmatrix} \langle \tilde{\mathcal{O}}_i \tilde{\mathcal{O}}_j^\dagger \rangle_{\text{Born}} & 0 \\ 0 & \langle L_i L_j^\dagger \rangle_{\text{Born}} \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & \langle L_i L_j^\dagger \rangle_{g^2} \end{pmatrix} \tag{3.77}$$

where the (symmetric) blocks are

$$\langle L_i L_j^\dagger \rangle_{\text{Born}} = \tfrac{(N^2-1)}{64} \begin{pmatrix} 7N^4+20N^2+8 & -4N^4-10N^2+4 & 2N(13N^2-2) \\ & 3N^4+2 & -2N(6N^2+1) \\ & & 2N^2(3N^2+13) \end{pmatrix} \tag{3.78}$$

and

$$\langle L_i L_j^\dagger \rangle_{g^2} = \tfrac{3\tilde{B}N(N^2-1)}{32} \begin{pmatrix} 25N^4+148N^2+8 & 15N^4-66N^2+4 & 4N(27N^2+17) \\ & 10N^4+22N^2+2 & -2N(28N^2+13) \\ & & 2N^2(9N^2+89) \end{pmatrix} \tag{3.79}$$

and

$$\begin{aligned}
\langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_1^\dagger \rangle_{\text{Born}} &= \frac{360\ \mathcal{C}_N}{N^2(3N^2-2)} \times (N^6-11N^4+70N^2-48) \\
\langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_2^\dagger \rangle_{\text{Born}} &= \frac{-120\ \mathcal{C}_N}{N^2(3N^2-2)} \times (5N^4-36N^2+24) \\
\langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_3^\dagger \rangle_{\text{Born}} &= \frac{240\ \mathcal{C}_N}{N(3N^2-2)} \times (N^2-2)(2N^2-3) \\
\langle \tilde{\mathcal{O}}_2 \tilde{\mathcal{O}}_2^\dagger \rangle_{\text{Born}} &= \frac{5\ \mathcal{C}_N}{N^2(3N^2-2)} \times (3N^6-41N^4+160N^2-96) \\
\langle \tilde{\mathcal{O}}_2 \tilde{\mathcal{O}}_3^\dagger \rangle_{\text{Born}} &= \frac{-20\ \mathcal{C}_N}{N(3N^2-2)} \times (13N^2-12) \\
\langle \tilde{\mathcal{O}}_3 \tilde{\mathcal{O}}_3^\dagger \rangle_{\text{Born}} &= \frac{60\ \mathcal{C}_N}{(3N^2-2)} \times (N^2+1)(N^2-2).
\end{aligned} \tag{3.80}$$

Here and below we shall use the abbreviation,

$$\mathcal{C}_N \equiv (N^2-1)(N^2-4)/64. \tag{3.81}$$

As seen from (3.77), the operators defined in (3.76) have protected two-point functions $\langle \tilde{\mathcal{O}}_i \tilde{\mathcal{O}}_j^\dagger \rangle$ at order g^2. This shows that we can argue that $\tilde{\mathcal{O}}_1$, $\tilde{\mathcal{O}}_2$, $\tilde{\mathcal{O}}_3$

are 1/4-BPS. Anomalous scaling dimensions of long operators ($L_i = \mathcal{O}_4, \mathcal{O}_5, \mathcal{O}_6$) match those of their Konishi-like primaries computed in [64].

A Better basis for protected [2,2,2] operators

It may seem odd that the operators mixing of the operators in the representations $[2,0,2]$ and $[2,1,2]$ are in terms of coefficients that are merely inverse powers of N, while the mixing coefficients for the operators we identified in the representation $[2,2,2]$ have more complicated denominators. This distinction would also be surprising from the perspective of AdS/CFT, since the more complicated denominators would suggest that an infinite series of corrections in the string coupling $g_s = \lambda/N$ would appear for given 't Hooft coupling λ. As a matter of fact, the mixing coefficients depend upon the bases chosen for both the $\mathcal{O}_1$, $\mathcal{O}_2$ and $\mathcal{O}_3$ operators as well as the pure descendants. In a different basis, the coefficients are all proportional to inverse powers of N. For the representation $[2,2,2]$, these new operators are found easily, and we have

$$\begin{aligned}
\tilde{\mathcal{O}}'_1 &= \tilde{\mathcal{O}}_1 + \frac{4}{N}\tilde{\mathcal{O}}_3 = \mathcal{O}_1 + \frac{4}{N}\mathcal{O}_3 - \frac{24}{N}\mathcal{O}_4 - \frac{32}{N}\mathcal{O}_5 \\
\tilde{\mathcal{O}}'_2 &= \tilde{\mathcal{O}}_2 + \frac{1}{2N}\tilde{\mathcal{O}}_3 = \mathcal{O}_2 - \frac{4}{N}\mathcal{O}_4 - \frac{7}{N}\mathcal{O}_5 \\
\tilde{\mathcal{O}}'_3 &= \tilde{\mathcal{O}}_3 + \frac{2}{3N}\tilde{\mathcal{O}}_1 - \frac{4}{N}\tilde{\mathcal{O}}_2 = \mathcal{O}_3 + \frac{2}{3N}\mathcal{O}_1 - \frac{4}{N}\mathcal{O}_2 - \frac{10}{3N}\mathcal{O}_6 \qquad (3.82)
\end{aligned}$$

In this new basis, the matrix of 2-pt functions now reads

$$\langle \tilde{\mathcal{O}}'_i \tilde{\mathcal{O}}'^{\dagger}_j \rangle_{\mathrm{Born}} = \frac{5\mathcal{C}_N}{N^2} \begin{pmatrix} 24(N^4+3N^2+32) & -16(N^2+9) & 16N(4N^2-1) \\ & (N^2-3)(N^2+9) & -2N(N^2-9) \\ & & 4(N^4+17N^2-24) \end{pmatrix} \qquad (3.83)$$

The Representation [2,3,2]

Here we have 8 operators. The operators corresponding to the basis (3.31) are

$$\begin{aligned}
\mathcal{O}_1 &\equiv 2\,\mathrm{tr}\, z_1z_1z_1z_1z_1 \;\mathrm{tr}\, z_2z_2 - 4\,\mathrm{tr}\, z_1z_1z_1z_1z_2 \;\mathrm{tr}\, z_1z_2 \\
&\qquad\qquad + (\mathrm{tr}\, z_1z_1z_1z_2z_2 + \mathrm{tr}\, z_1z_1z_2z_1z_2)\,\mathrm{tr}\, z_1z_1 \\
\mathcal{O}_2 &\equiv 3\,\mathrm{tr}\, z_1z_1z_1z_1 \;\mathrm{tr}\, z_1z_2z_2 - 6\,\mathrm{tr}\, z_1z_1z_1z_2 \;\mathrm{tr}\, z_1z_1z_2 \\
&\qquad\qquad + (2\,\mathrm{tr}\, z_1z_1z_2z_2 + \mathrm{tr}\, z_1z_2z_1z_2)\,\mathrm{tr}\, z_1z_1z_1 \\
\mathcal{O}_3 &\equiv \mathrm{tr}\, z_1z_1z_1\,(\mathrm{tr}\, z_1z_1 \;\mathrm{tr}\, z_2z_2 - \mathrm{tr}\, z_1z_2 \;\mathrm{tr}\, z_1z_2) \\
\mathcal{O}_4 &\equiv \mathrm{tr}\, z_1z_1z_1 \;\mathrm{tr}\, z_1z_1 \;\mathrm{tr}\, z_2z_2 - 2\,\mathrm{tr}\, z_1z_1z_2 \;\mathrm{tr}\, z_1z_1 \;\mathrm{tr}\, z_1z_2 \\
&\qquad\qquad + \mathrm{tr}\, z_1z_2z_2 \;\mathrm{tr}\, z_1z_1 \;\mathrm{tr}\, z_1z_1 \\
\mathcal{O}_5 &\equiv \mathrm{tr}\, z_1z_1z_1z_1z_1z_2z_2 - 2\,\mathrm{tr}\, z_1z_1z_1z_1z_2z_1z_2 + \mathrm{tr}\, z_1z_1z_1z_2z_1z_1z_2 \\
\mathcal{O}_6 &\equiv \mathrm{tr}\, z_1z_1z_1z_1z_2z_1z_2 - \mathrm{tr}\, z_1z_1z_1z_2z_1z_1z_2 \\
\mathcal{O}_7 &\equiv (\mathrm{tr}\, z_1z_1z_1z_2z_2 - \mathrm{tr}\, z_1z_1z_2z_1z_2)\;\mathrm{tr}\, z_1z_1 \\
\mathcal{O}_8 &\equiv (\mathrm{tr}\, z_1z_1z_2z_2 - \mathrm{tr}\, z_1z_2z_1z_2)\;\mathrm{tr}\, z_1z_1z_1
\end{aligned} \tag{3.84}$$

Out of these, four are descendants,

$$\begin{aligned}
\mathcal{O}_5 &\sim (Q^2\bar{Q}^2)\;\mathrm{tr}\,\left[2z_1z_1z_1z_j\bar{z}^j + 2z_1z_1z_1\bar{z}^jz_j - z_1z_1z_jz_1\bar{z}^j - z_1z_1\bar{z}^jz_1z_j\right] \\
\mathcal{O}_6 &\sim (Q^2\bar{Q}^2)\;\mathrm{tr}\,\left[z_1z_1z_1z_j\bar{z}^j + z_1z_1z_1\bar{z}^jz_j - 2z_1z_1z_jz_1\bar{z}^j - 2z_1z_1\bar{z}^jz_1z_j\right] \\
\mathcal{O}_7 &\sim (Q^2\bar{Q}^2)\left[\mathrm{tr}\, z_1z_1z_j \;\mathrm{tr}\, z_1\bar{z}^j + \mathrm{tr}\, z_1z_1\bar{z}^j \;\mathrm{tr}\, z_1z_j - 12\,\mathrm{tr}\, z_1z_1z_1 \;\mathrm{tr}\, z_j\bar{z}^j\right] \\
\mathcal{O}_8 &\sim (Q^2\bar{Q}^2)\left[\mathrm{tr}\, z_1z_1z_1 \;\mathrm{tr}\, z_j\bar{z}^j\right]
\end{aligned} \tag{3.85}$$

while the combinations orthogonal to them at Born level can be taken as

$$\begin{aligned}
\tilde{\mathcal{O}}_1 &= \mathcal{O}_1 - \frac{20}{N}\mathcal{O}_5 - \frac{30(N^2-2)}{N^3}\mathcal{O}_6 + \frac{15(N^2-2)}{N^4}\mathcal{O}_7 + \frac{10(N^2+2)}{N^4}\mathcal{O}_8 \\
\tilde{\mathcal{O}}_2 &= \mathcal{O}_2 - \frac{30}{N}\mathcal{O}_5 - \frac{30(2N^2-3)}{N^3}\mathcal{O}_6 + \frac{15(2N^2-3)}{N^4}\mathcal{O}_7 + \frac{10(N^2+3)}{N^4}\mathcal{O}_8 \\
\tilde{\mathcal{O}}_3 &= \mathcal{O}_3 \qquad\quad - \frac{12}{N^2}\mathcal{O}_6 - \frac{3(N^2-2)}{N^3}\mathcal{O}_7 - \frac{2(N^2+2)}{N^3}\mathcal{O}_8 \\
\tilde{\mathcal{O}}_4 &= \mathcal{O}_4 \qquad\quad - \frac{18}{N^2}\mathcal{O}_6 - \frac{(7N^2-9)}{N^3}\mathcal{O}_7 - \frac{2(2N^2+9)}{3N^3}\mathcal{O}_8
\end{aligned} \tag{3.86}$$

The matrix of two point functions in this basis is

$$\begin{pmatrix} \langle \tilde{\mathcal{O}}_i \tilde{\mathcal{O}}_j^\dagger \rangle_{\text{Born}} & 0 \\ 0 & \langle L_i L_j^\dagger \rangle_{\text{Born}} \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & \langle L_i L_j^\dagger \rangle_{g^2} \end{pmatrix} \tag{3.87}$$

so indeed at order g^2 the operators defined in (3.86) have protected correlators. This shows that we can argue that $\tilde{\mathcal{O}}_1$, $\tilde{\mathcal{O}}_2$, $\tilde{\mathcal{O}}_3$, $\tilde{\mathcal{O}}_4$ are the 1/4-BPS primaries we are after. The (symmetric) blocks in equation (3.87) are

$$\langle L_i L_j^\dagger \rangle_{\text{Born}} = \tfrac{1}{2} N \mathcal{C}_N \begin{pmatrix} 6N^2+45 & -3N^2-9 & 18N & 36N \\ & 2N^2-3 & -4N & -15N \\ & & 4N^2+24 & 36 \\ & & & 9N^2+54 \end{pmatrix} \tag{3.88}$$

and

$$\begin{aligned} \langle L_i L_j^\dagger \rangle_{g^2} &= 12 \tilde{B} N \mathcal{C}_N \\ & \quad \times \begin{pmatrix} 9N(2N^2+27) & -9N(N^2+8) & 54(N^2+2) & 27(5N^2+6) \\ & N(5N^2+17) & -6(3N^2+2) & -6(10N^2+3) \\ & & 4N(2N^2+23) & 174N \\ & & & 9N(3N^2+35) \end{pmatrix} \end{aligned} \tag{3.89}$$

The Born level overlaps of protected operators are given by ugly and not particularly illuminating expressions. Here we list them for the sake of completeness:

$$\begin{aligned} \langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_1^\dagger \rangle_{\text{Born}} &= 30\ \mathcal{C}_N N^{-5}(N^8 - 10N^6 + 117N^4 - 720N^2 + 420) \\ \langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_2^\dagger \rangle_{\text{Born}} &= -90\ \mathcal{C}_N N^{-5}(4N^6 - 69N^4 + 395N^2 - 210) \\ \langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_3^\dagger \rangle_{\text{Born}} &= 90\ \mathcal{C}_N N^{-4}(N^2-2)(N^4 - 7N^2 + 14) \\ \langle \tilde{\mathcal{O}}_1 \tilde{\mathcal{O}}_4^\dagger \rangle_{\text{Born}} &= 30\ \mathcal{C}_N N^{-4}(N^2-2)(N^2-9)(3N^2-7) \\ \langle \tilde{\mathcal{O}}_2 \tilde{\mathcal{O}}_2^\dagger \rangle_{\text{Born}} &= 45\ \mathcal{C}_N N^{-5}(N^8 - 29N^6 + 328N^4 - 1290N^2 + 630) \\ \langle \tilde{\mathcal{O}}_2 \tilde{\mathcal{O}}_3^\dagger \rangle_{\text{Born}} &= -630\ \mathcal{C}_N N^{-4}(N^2-1)(N^2-6) \end{aligned}$$

$$\begin{aligned}
\langle\tilde{\mathcal{O}}_2\tilde{\mathcal{O}}_4^\dagger\rangle_{\text{Born}} &= 30\,\mathcal{C}_N N^{-4}(N^2-1)(N^2-9)(2N^2-21) \\
\langle\tilde{\mathcal{O}}_3\tilde{\mathcal{O}}_3^\dagger\rangle_{\text{Born}} &= 9\,\mathcal{C}_N N^{-3}(N^6-N^4-16N^2+56) \\
\langle\tilde{\mathcal{O}}_3\tilde{\mathcal{O}}_4^\dagger\rangle_{\text{Born}} &= 6\,\mathcal{C}_N N^{-3}(N^2-9)(N^4+3N^2-14) \\
\langle\tilde{\mathcal{O}}_4\tilde{\mathcal{O}}_4^\dagger\rangle_{\text{Born}} &= 2\,\mathcal{C}_N N^{-3}(N^2-9)(7N^4+16N^2-63) \qquad (3.90)
\end{aligned}$$

and the constant $\mathcal{C}_N = (N^2-1)(N^2-4)/64$ was defined in (3.81).

The Representation [3,1,3]

Here there are 5 operators. The operators corresponding to the basis (3.36) are

$$\begin{aligned}
\mathcal{O}_1 &\equiv \text{tr}\, z_1z_1z_1z_1\ \text{tr}\, z_2z_2z_2 - 3\,\text{tr}\, z_1z_1z_1z_2\ \text{tr}\, z_1z_2z_2 \\
&\qquad + (2\,\text{tr}\, z_1z_1z_2z_2 + \text{tr}\, z_1z_2z_1z_2)\,\text{tr}\, z_1z_1z_2 - \text{tr}\, z_1z_2z_2z_2\ \text{tr}\, z_1z_1z_1 \\
\mathcal{O}_2 &\equiv \text{tr}\, z_1z_2\,(2\,\text{tr}\, z_1z_2\ \text{tr}\, z_1z_2z_2 - \text{tr}\, z_2z_2\ \text{tr}\, z_1z_1z_1 - 3\,\text{tr}\, z_1z_1\ \text{tr}\, z_1z_2z_2) \\
&\qquad + \text{tr}\, z_1z_1\,(\text{tr}\, z_2z_2\ \text{tr}\, z_1z_1z_2 + \text{tr}\, z_1z_1\ \text{tr}\, z_2z_2z_2) \\
\mathcal{O}_3 &\equiv \text{tr}\, z_1z_1z_1z_2z_1z_2z_2 - \text{tr}\, z_1z_1z_1z_2z_2z_1z_2 \\
\mathcal{O}_4 &\equiv 2\,\text{tr}\, z_1z_1z_1z_1z_2z_2z_2 - 3\,\text{tr}\, z_1z_1z_1z_2z_1z_2z_2 - 3\,\text{tr}\, z_1z_1z_1z_2z_2z_1z_2 \\
&\qquad + 2\,\text{tr}\, z_1z_1z_2z_1z_2z_1z_2 + 2\,\text{tr}\, z_1z_1z_2z_2z_1z_1z_2 \\
\mathcal{O}_5 &\equiv -(\text{tr}\, z_1z_1z_2z_2z_2 - \text{tr}\, z_1z_2z_1z_2z_2)\,\text{tr}\, z_1z_1 \\
&\qquad + (\text{tr}\, z_1z_1z_1z_2z_2 - \text{tr}\, z_1z_1z_2z_1z_2)\,\text{tr}\, z_1z_2 \qquad (3.91)
\end{aligned}$$

Out of these three are descendants,

$$\begin{aligned}
\mathcal{O}_3 &\sim (Q^2\bar{Q}^2)\ \text{tr}\,\left[z_1z_1z_2z_j\bar{z}^j + z_1z_1z_2\bar{z}^jz_j - z_1z_1z_j\bar{z}^jz_2 - z_1z_1\bar{z}^jz_jz_2\right] \\
\mathcal{O}_4 &\sim (Q^2\bar{Q}^2)\ \text{tr}\,\left[z_1z_1z_2z_j\bar{z}^j + z_1z_1z_2\bar{z}^jz_j + z_1z_1z_j\bar{z}^jz_2 + z_1z_1\bar{z}^jz_jz_2\right. \\
&\qquad \left. -2z_1z_2z_1z_j\bar{z}^j - 2z_1z_2z_1\bar{z}^jz_j\right] \qquad (3.92) \\
\mathcal{O}_5 &\sim (Q^2\bar{Q}^2)\ \left[2\,\text{tr}\, z_1z_1z_j\ \text{tr}\, z_2\bar{z}^j + 2\,\text{tr}\, z_1z_1\bar{z}^j\ \text{tr}\, z_2z_j - \text{tr}\, z_1z_2\bar{z}^j\ \text{tr}\, z_1z_j\right. \\
&\qquad \left. -\text{tr}\, z_1z_2z_j\ \text{tr}\, z_1\bar{z}^j - \text{tr}\, z_2z_1\bar{z}^j\ \text{tr}\, z_1z_j - \text{tr}\, z_2z_1z_j\ \text{tr}\, z_1\bar{z}^j\right]
\end{aligned}$$

while the combinations orthogonal to them at Born level can be taken as

$$\begin{aligned}\tilde{\mathcal{O}}_1 &= \mathcal{O}_1 - \frac{2N}{N^2-2}\mathcal{O}_4 - \frac{5}{N^2-2}\mathcal{O}_5 \\ \tilde{\mathcal{O}}_2 &= \mathcal{O}_2 + \frac{8}{N^2-2}\mathcal{O}_4 + \frac{10N}{N^2-2}\mathcal{O}_5 \end{aligned} \tag{3.93}$$

(which are the operators found in Chapter 2). The matrix of two point functions in this basis is

$$\begin{pmatrix} \langle \tilde{\mathcal{O}}_i \tilde{\mathcal{O}}_j^\dagger \rangle_{\text{Born}} & 0 \\ 0 & \langle L_i L_j^\dagger \rangle_{\text{Born}} \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & \langle L_i L_j^\dagger \rangle_{g^2} \end{pmatrix} \tag{3.94}$$

so indeed at order g^2 the operators defined in (3.93) have protected correlators. This shows that we can argue that $\tilde{\mathcal{O}}_1$, $\tilde{\mathcal{O}}_2$, are the 1/4-BPS primaries we are after. The (symmetric) blocks in (3.94) are

$$\begin{aligned}\langle L_i L_j^\dagger \rangle_{\text{Born}} &= \mathcal{C}_N \begin{pmatrix} N(N^2-9) & 0 & 0 \\ & 15N(N^2+3) & -30N^2 \\ & & 3N(N^2+6) \end{pmatrix} \\ \langle L_i L_j^\dagger \rangle_{g^2} &= 6\tilde{B}N\mathcal{C}_N \begin{pmatrix} 5N(N^4-9) & 0 & 0 \\ & 75N(N^2+7) & 180(N^2+1) \\ & & 12N(N^2+16) \end{pmatrix} \\ \langle \tilde{\mathcal{O}}_i \tilde{\mathcal{O}}_j^\dagger \rangle_{\text{Born}} &= \frac{15(N^2-9)}{N^2(N^2-2)}\,\mathcal{C}_N \begin{pmatrix} (N^2-1)(N^2-4) & 4N(N^2-1) \\ & 2N^2(N^2-6) \end{pmatrix} \end{aligned} \tag{3.95}$$

As mentioned before, the operator $\mathcal{O}_3$ has zero correlators with everything else. The reason is that $\mathcal{O}_3^\dagger = -(\mathcal{O}_3)^*$, while for all other operators $\mathcal{O}_i^\dagger = +(\mathcal{O}_i)^*$.

Completeness of the Construction

An important point which remains to be addressed is whether the construction of the 1/4 BPS operators given above is exhaustive. The fact that it is follows from $SU(4)$ group theory in the following manner. Given a 1/4 BPS representation of $SU(4)$ of the type $[q,p,q]$, one begins by listing all possible monomial scalar composite operators built out of $(p+q)$ z_1's and q z_2's.[4] These monomials form a basis for the linear space of scalar composite operators of the form $[(z_1)^{(p+q)}\,(z_2)^q]$. They can occur in representations $[0,p+2q,0]$, $[1,p+2q-2,1]$, ... , $[q,p,q]$. Then we have to show that the number of such monomials matches the total number of operators we constructed in these representations.

Let us illustrate how this works with an example. Consider the [2,2,2] representation. The complete set of scalar composite operators we can build out of 4 z_1's and 2 z_2's is

$$\begin{aligned}
6: \quad & \mathrm{tr}\, z_1z_1z_1z_1z_2z_2, \quad \mathrm{tr}\, z_1z_1z_1z_2z_1z_2, \quad \mathrm{tr}\, z_1z_1z_2z_1z_1z_2 \\
4+2: \quad & \mathrm{tr}\, z_1z_1z_1z_1\ \mathrm{tr}\, z_2z_2, \quad \mathrm{tr}\, z_1z_1z_1z_2\ \mathrm{tr}\, z_1z_2, \quad \mathrm{tr}\, z_1z_1z_2z_2\ \mathrm{tr}\, z_1z_1, \\
& \mathrm{tr}\, z_1z_2z_1z_2\ \mathrm{tr}\, z_1z_1 \\
3+3: \quad & \mathrm{tr}\, z_1z_1z_1\ \mathrm{tr}\, z_1z_2z_2, \quad \mathrm{tr}\, z_1z_1z_2\ \mathrm{tr}\, z_1z_1z_2 \\
2+2+2: \quad & \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_2z_2, \quad \mathrm{tr}\, z_1z_1\ \mathrm{tr}\, z_1z_2\ \mathrm{tr}\, z_1z_2
\end{aligned} \tag{3.96}$$

or the total of 11 operators. By taking linear combinations of these, we can construct:

- 4 tensors in the [0,6,0] corresponding to the partitions of 6 in (3.96);
- 1 tensor in the representation [1,4,1], given in (3.56);
- 6 tensors in the representation [2,2,2], listed in (3.72).

Thus there are no other scalar composite operators of the form $[(z_1)^4\,(z_2)^2]$.

In the same fashion, we can go through all other representations we have considered in this Chapter and verify that we didn't leave out any operators.

---

[4] The remaining scalar field z_3 never enters in the highest weight of a 1/4 BPS representation, though it will also be needed when describing 1/8 BPS operators.

CHAPTER 4

Three-Point Functions of Quarter BPS Operators in $\mathcal{N}$=4 SYM

In Chapters 2 and 3, we constructed $\frac{1}{4}$-BPS chiral primaries in the fully interacting $\mathcal{N}$=4 SYM theory. In general, these operators are linear combinations of all local, polynomial, gauge invariant, scalar composite operators in the $[p, q, p]$ representations of the R-symmetry group $SU(4)$. The coefficients with which they all combine into operators with a well defined scaling dimension are quite involved. The $\frac{1}{4}$-BPS chiral primaries, like the $\frac{1}{2}$-BPS operators extensively studied in the literature, have protected two-point functions (at least at order g^2).

Here we investigate the (non-) renormalization properties of three-point correlators involving $\frac{1}{4}$-BPS operators along with $\frac{1}{2}$-BPS operators. Given the elaborate combinatorics of the problem, we concentrate on the following special cases. First, we discuss several group theoretic simplifications of the combinatorial factors multiplying the Feynman graphs that contribute to three-point functions of chiral primaries. Based on $SU(4)$ group theory and conformal invariance only, we argue that certain classes of such correlators are protected at order g^2, for all N. In particular, this allows us to compute $\mathcal{O}(g^2)$ corrections to correlators of the form $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\mathrm{BPS}} \rangle$, where $\mathcal{O}_{\frac{1}{2}}$ are two $\frac{1}{2}$-BPS operators, and $\mathcal{O}_{\mathrm{BPS}}$ is an arbitrary ($\frac{1}{2}$-, $\frac{1}{4}$-, or $\frac{1}{8}$-BPS) chiral primary. Next, we look at the three-point functions $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$ and $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$, also for general N, where $\mathcal{O}_{\frac{1}{4}}$ are the $\Delta \leq 7$ $\frac{1}{4}$-BPS primaries found in Chapters 2 and 3. Then, we carry out a large N analysis of $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$ correlators involving the special $\frac{1}{4}$-BPS operators (mixtures of single and double trace scalar composite operators), for arbitrary Δ. Also in the large N limit, we identify the corresponding objects in the supergravity description, and compute the correlators on the AdS side of the correspondence.

Finally, we make some speculations. Based on the broad range of special cases, we conjecture that two- and three-point functions of $\frac{1}{2}$- and $\frac{1}{4}$-operators receive no quantum corrections, for arbitrary N. Additionally, a set of group theoretic considerations discussed here extends straightforwardly from three-point functions to extremal correlators. Therefore, we suggest that extremal correlators involving $\frac{1}{2}$- and $\frac{1}{4}$-operators are protected as well.

4.1 Contributing diagrams

The two-point functions of the scalar composites discussed in Chapters 2 and 3, have the schematic form

$$\langle \left[{z_1}^{(p+q)} {z_2}^{p} \right] (x) \left[\bar{z}_1^{(p+q)} \bar{z}_2^{p} \right] (y) \rangle \tag{4.1}$$

where [...] are some gauge invariant combinations. The free field part of (4.1) is given by a power of the free correlator $[G(x,y)]^{(2p+q)}$, times a combinatoric factor. From the Lagrangian (2.1) we can read off the structures for the leading correction to the propagator, and the four-scalar blocks, see Figure 2.1.

Three-point functions to be considered in this Chapter are of the form

$$\langle \left[z^{k+l} \right] (x) \left[\bar{z}^{k+m} \right] (y) \left[\bar{z}^{l} z^{m} \right] (w) \rangle \tag{4.2}$$

The free result is just the product of appropriate powers of free correlators $[G(x,y)]^k [G(x,w)]^l [G(y,w)]^m$. The same structures that contribute to the two-


(x)
i, a
j, b
i, a' (y)
j, b'(w)
= f^{apa'} f^{bpb'} C (x;y,w) G(x,y) G(x,w)
(x)
i, a
j, b
i, a' (y)
j, b'(w)
= f^{apa'} f^{bpb'} C' (x;y,w) G(x,y) G(x,w)
(x)
i, a
i, a'
j, b' (y)
j, b (w)
= f^{apa'} f^{bpb'} C" (x;y,w) G(x,y) G(x,w)
(x)
i, a
j, b'
j, b (y)
i, a'(w)
= f^{apb} f^{a'pb'} C̃ (x;y,w) G(x,y) G(x,w)
(x)
i, a
j, b
i, a' (y)
j, b'(w)
= f^{apb} f^{a'pb'} C̃ (x;y,w) G(x,y) G(x,w)

Figure 4.1: Building blocks for g^2 corrections to three-point functions. The three space-time points are x (with two legs attached) and y and w (single leg each).

point functions at order g^2 (see Figure 2.1), also contribute to the three-point functions (4.2). Apart from these, there are new building blocks, shown in Figure 4.1. They have the same index structure, but are now functions of three space-time coordinates rather than two.

Notice that the F-term corrections proportional to $\tilde{B}(x,y)$ in Figure 2.1, and the last graph (proportional to $\tilde{C}(x;y,w)$) in Figure 4.1, are antisymmetric in i and j, hence they are absent when the scalars in the four legs have pairwise the same flavor. For the same reason, these corrections are also absent when the operator at point x is symmetric in all of its flavor indices. In particular, this is the case when the operator at x is $\frac{1}{2}$-BPS.

4.2 Restrictions from $\mathcal{N}$=4 SUSY and gauge invariance

The form of quantum corrections to two and three-point functions is known [3]. Space-time coordinate dependence of the Feynman diagrams contributing to these correlators at order $\mathcal{O}(g^2)$ is constrained, since all exchanged fields are massless. We know the parametric form of the functions $A(x,y)$, $B(x,y)$, $\tilde{B}(x,y)$; and $C(x;y,w)$, $C'(x;y,w)$, $C''(x;y,w)$, and $\tilde{C}(x;y,w)$, without having to perform integrals explicitly. Functions which depend on two space-time points, are of the form $A(x_1,x_2) = a \log x_{12}^2\mu^2 + b$ with $x_{ij} \equiv x_i - x_j$; three-point contributions look like $C(x_1;x_2,x_3) = a' \log x_{12}^2 x_{13}^2 \mu^4 - a'' \log x_{23}^2 + b'$ (making use of the $x_2 \leftrightarrow x_3$ symmetry of these building blocks).

$\mathcal{N}$=4 SUSY tells us more. From non-renormalization of two and three-point functions of operators in the stress tensor multiplet, one can see [3] that $B(x,y) = -2A(x,y)$, and $C'(0;x,y) + \tilde{C}(0;x,y) = -C(0;x,y)$; the authors of [3] chose to combine these and call it just C'.[1] The coefficients a', a'' and b' are determined[2] in terms of a and b:

$$\begin{aligned} A(x,0) \;=\; -\tfrac{1}{2}B(x,0) &= a \log x^2\mu^2 + b \\ -C(0;x,y) &= a \log \frac{x^2y^2\mu^2}{(x-y)^2} + b \end{aligned} \tag{4.3}$$

Therefore, the net contribution to the three-point function (4.2) of the $\mathcal{O}(g^2)$ diagrams involving a gauge boson exchange (diagrams A, B, and C), is

$$\begin{aligned} &\langle [z^k](x_1)[\bar{z}^l](x_2)[z^m\bar{z}^n](x_3)\rangle|_{(A+B+C)} \\ &= a(c_g^{12}\log x_{12}^2\mu^2 + c_g^{13}\log x_{13}^2\mu^2 + c_g^{23}\log x_{23}^2\mu^2) + bc_g^{123} \end{aligned} \tag{4.4}$$

where and c_g^{ij} and c_g^{123} are some combinatorial coefficients.

---

[1] One way to see this is to consider the protected correlators of [0,2,0] scalar composite operators $\langle \mathrm{tr}\, z_1z_2(x)\ \mathrm{tr}\, \bar{z}_1\bar{z}_2(y)\rangle$, and $\langle [z^2](x)[\bar{z}^2](y)[z\bar{z}](w)\rangle$ and $\langle [z\bar{z}](x)[z\bar{z}](y)[z\bar{z}](w)\rangle$.

[2] This follows from $C(x;y,w) + C(y;x,w) + C(w;x,y) + A(x,y) + A(y,w) + A(x,w) = 0$.

Now we use gauge invariance of the theory. On the one hand, we observe that the coefficients a and b are gauge dependent as we saw in Chapter 2, Section 2.5,

$$A(x,0) \;=\; \tfrac{1}{2}\pi^2 g^2 \xi \left[\log x^2\mu^2 + \log 4\pi - \gamma\right] + (\xi\text{-independent}) \qquad (4.5)$$

where ξ is the gauge fixing parameter. On the other hand, a correlator of gauge invariant operators can not depend on ξ. Therefore, the combinatorial coefficients multiplying a and b in equation (4.4) must vanish, $c_g^{ij} = c_g^{123} = 0$. Hence, the D-term diagrams proportional to A, B, and C all cancel; their net contribution to the three-point functions (4.2) is zero.

So just like in the case of two-point functions, we only have to consider the F-term graphs. They are proportional to $\tilde{B}$ and $\tilde{C}$, the only gauge independent diagrams around ($C' = -(C + \tilde{C})$ and $C'' = C - \tilde{C}$ do not have to be treated separately as they are linear combinations of the other ones).

4.3 Position dependence of $\tilde{B}$ and $\tilde{C}$

Having shown that D-term corrections to three-point functions (4.2) are absent, it remains to consider the F-term interactions. In this Section we derive a relation between functions $\tilde{B}$ and $\tilde{C}$, which will play a key role in the analysis of three-point functions of $\frac{1}{4}$-BPS chiral primaries, see Section 4.5.1.1.

Space-time position dependence of $\tilde{B}$ and $\tilde{C}$ (shown Figures 2.1 and 4.1) is parametrically determined to be $\tilde{B}(x,0) = \tilde{a}\log(x^2\mu^2) + \tilde{b}$ and $\tilde{C}(0;x,y) = \tilde{a}'\log(x^2y^2\mu^4) - \tilde{a}''\log((x-y)^2\mu^2) + \tilde{c}$. Furthermore, the leading divergent behavior can be read off from the integrals unambiguously, and so from the limit $\tilde{C}(0;x,y\to x)$ we infer $\tilde{a}' = \frac{1}{2}\tilde{a}$.

To evaluate the remaining coefficients $\tilde{a}$, $\tilde{a}''$, and $\tilde{b}$, replace $1/x^2 \to 1/(x^2+\epsilon^2)$

for scalar propagators inside integrals.[3] With this prescription

$$\begin{aligned}\tilde{B}(x,0) &= -\tfrac{1}{4}Y^2 \int \frac{(d^4z)\,[4\pi^2x^2]^2}{[4\pi^2((z-x)^2+\epsilon^2)]^2\,[4\pi^2(z^2+\epsilon^2)]^2} \\ &= -Y^2\frac{1}{32\pi^2}\left[\log(x^2/\epsilon^2)-1\right] \qquad (4.6)\end{aligned}$$

is the regularized two-point function, while the three-point function becomes

$$\begin{aligned}\tilde{C}(x;y,0) &= -\tfrac{1}{4}Y^2 \int \frac{(d^4z)\,[4\pi^2x^2]\,[4\pi^2(x-y)^2]}{[4\pi^2((z-x)^2+\epsilon^2)]^2\,[4\pi^2((z-y)^2+\epsilon^2)]\,[4\pi^2(z^2+\epsilon^2)]} \\ &= -Y^2\frac{1}{64\pi^2}\left[\log\frac{x^2(x-y)^2}{y^2\epsilon^2}\right] \qquad (4.7)\end{aligned}$$

(The numerators inside the integrals come about because of the powers of free scalar propagator in the definitions of $\tilde{B}$ and $\tilde{C}$, see Figures 2.1 and 4.1.) Hence,

$$\tilde{C}(x;y,0)+\tilde{C}(y;x,0)-\tilde{B}(x,y) = -Y^2\times\frac{1}{32\pi^2} \qquad (4.8)$$

is a nonzero constant (for $\mathcal{N}$=4 SUSY, $Y^2 = 2g^2$). The value of this constant does not depend on the regulator ϵ. Also note that with the "point splitting regularization" one would get the incorrect result of vanishing constant in (4.8).

4.4 Structure of the three-point functions

With the results of Section 4.2 at hand, we can write down the form of a general three-point function of scalar composite operators (4.2) to order g^2:

$$\begin{aligned}\langle\left[z^{k+l}\right](x)\left[\bar{z}^{k+m}\right](y)\left[\bar{z}^l z^m\right](w)\rangle &= G(x,y)^kG(x,w)^lG(w,y)^m \\ &\times\Big(\alpha_{\text{free}}+\tilde{\beta}_{xy}\tilde{B}(x,y)+\tilde{\beta}_{xw}\tilde{B}(x,w)+\tilde{\beta}_{yw}\tilde{B}(y,w) \\ &\quad+\tilde{\gamma}_x\tilde{C}(x;y,w)+\tilde{\gamma}_y\tilde{C}(y;x,w)+\tilde{\gamma}_w\tilde{C}(w;x,y) \\ &\quad+\mathcal{O}(g^4)\Big) \qquad (4.9)\end{aligned}$$

---

[3] This is the fastest way to calculate the integral for $\tilde{C}$, but one can obtain the same results using dimensional regularization.

where α_{free}, $\tilde{\beta}$-s and $\tilde{\gamma}$-s are some combinatorial coefficients. Using the expressions (4.6) and (4.7) from Section 4.3, we can determine the $\mathcal{O}(g^2)$ position dependence of (4.9) completely — if we know these combinatorial coefficients. Together with conformal invariance, and the $SU(4)$ symmetry properties of the operators in (4.9), we can often go a long way to figuring out which of the combinatorial coefficients must vanish, without doing any actual calculations.

4.4.1 Space-time coordinate dependence

Like in the case of two-point functions, conformal invariance restricts position dependence of three-point correlators of pure operators (i.e. ones which have a well defined scaling dimension). Consider three (gauge invariant Lorentz scalar) operators $\mathcal{O}_1$, $\mathcal{O}_2$, and $\mathcal{O}_3$, of dimensions $\Delta_i = k_i + \delta_i$, inserted at corresponding points x_i. Let k_i be integers; and δ_i, the order g^2 corrections to the scaling dimensions (which may or may not be zero). The three-point function $\langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \rangle$ is completely determined up to a multiplicative constant $C_{123} = C^0_{123} + C^1_{123}$ (where again C^0_{123} is the free field result and $C^1_{123} \sim g^2$),

$$\begin{aligned}
\langle \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \mathcal{O}_3(x_3) \rangle &= \frac{C_{123}}{x_{12}^{\Delta_1+\Delta_2-\Delta_3} x_{13}^{\Delta_1+\Delta_3-\Delta_2} x_{23}^{\Delta_2+\Delta_3-\Delta_1}} \\
&= \langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \rangle_{\text{free}} \Big(1 + C^1_{123}/C^0_{123} \\
&\quad -\delta_1 \log \frac{x_{12}^2 x_{13}^2}{x_{23}^2 \epsilon^2} - \delta_2 \log \frac{x_{21}^2 x_{23}^2}{x_{13}^2 \epsilon^2} - \delta_3 \log \frac{x_{31}^2 x_{32}^2}{x_{12}^2 \epsilon^2} \\
&\quad + \mathcal{O}(g^4) \Big)
\end{aligned} \tag{4.10}$$

with $x_{ij} = x_i - x_j$ as usual.

Suppose that all three operators have protected scaling dimensions, $\delta_i = 0$. Then (4.10) reduces to

$$\langle \mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \mathcal{O}_3(x_3) \rangle = \langle \mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3 \rangle_{\text{free}} \Big(1 + C^1_{123}/C^0_{123} \Big) \tag{4.11}$$

and no logs arise. In this case, the combinatorial factors in (4.9) satisfy

$$\tilde{\gamma}_x = -\left(\tilde{\beta}_{xy} + \tilde{\beta}_{xw}\right) \tag{4.12}$$

$$\tilde{\gamma}_y = -\left(\tilde{\beta}_{xy} + \tilde{\beta}_{yw}\right) \tag{4.13}$$

$$\tilde{\gamma}_w = -\left(\tilde{\beta}_{xw} + \tilde{\beta}_{yw}\right) \tag{4.14}$$

and we need to calculate only three coefficients (the $\tilde{\beta}$-s, for example), to find all the $\mathcal{O}(g^2)$ corrections to this correlator. In fact, the only allowed correction is the constant $\tilde{\beta} \equiv \tilde{\beta}_{xy} + \tilde{\beta}_{xw} + \tilde{\beta}_{yw}$ times $(-Y^2/32\pi^2)$. To show that a three-point function of BPS operators is protected, we have to demonstrate that $\tilde{\beta} = 0$.

4.4.2 Group theory simplifications

There are several simplifications which set some of the combinatorial coefficients in (4.9) to zero. These considerations are based on the underlying $SU(4) \sim SO(6)$ symmetry of the theory only, and are applicable for general N. We will leave aside the trivial case when the correlator is forced to vanish by group theory, and assume that the Born level coefficient in (4.9) $\alpha_{\text{free}} \neq 0$.

The simplest BPS operators are $\frac{1}{2}$-BPS chiral primaries, gauge invariant scalar composites in $[0, q, 0]$ representations of $SO(6)$. These are totally symmetric tensors of $SO(6)$, so if for example the operator $\mathcal{O}_x$ is $\frac{1}{2}$-BPS, the coefficients $\tilde{\beta}_{xy} = \tilde{\beta}_{xw} = \tilde{\gamma}_x = 0$ since the diagrams they multiply are antisymmetric in the flavor indices of $\mathcal{O}_x$.[4] Similarly, if both $\mathcal{O}_x$ and $\mathcal{O}_y$ are $\frac{1}{2}$-BPS and $\mathcal{O}_w$ is any BPS operator, we have $\tilde{\beta}_{xy} = \tilde{\beta}_{xw} = \tilde{\gamma}_x = \tilde{\beta}_{yw} = \tilde{\gamma}_y = 0$, and hence $\tilde{\beta} = \tilde{\beta}_{xy} + \tilde{\beta}_{xw} + \tilde{\beta}_{yw} = 0$; there are no $\mathcal{O}(g^2)$ corrections in this case. In particular,

---

[4] If we chose $\mathcal{O}_w$ as such a $\frac{1}{2}$-BPS operator, we would not be able to conclude that $\tilde{\gamma}_w = 0$ just from the symmetries of $\mathcal{O}_w$: the fourth diagram of Figure 4.1, is not antisymmetric in flavor indices at the vertex where the operator is made of both z-s and $\bar{z}$-s. However, using equation (4.14) we find $\tilde{\gamma}_w = \tilde{\beta}_{xw} + \tilde{\beta}_{yw} = 0$ since $\tilde{\beta}_{xw} = \tilde{\beta}_{yw} = 0$ when all three operators have protected scaling dimensions.



Figure 4.2: F-term contributions to $\langle \mathcal{O}_{\rm BPS}(x)\, \mathcal{O}'_{\rm BPS}(y)\, \mathcal{O}_{\frac{1}{2}}(w)\rangle_{g^2}$ in the case when the correlator can be "partitioned into two flavors:" (a) proportional to $\tilde{B}(x,y)$; (b) proportional to $\tilde{C}(x;y,w)$; (c) proportional to $\tilde{C}(y;x,w)$.

this reproduces the result of [3] when all three $\mathcal{O}_{x,y,w}$ are $\frac{1}{2}$-BPS chiral primaries.

The "standard" way to choose $SU(4)$ weights of operators in a three point function is to take the highest weight state λ in its representation; another, the lowest weight state $-\lambda'$ in its; and the third operator to have weight $\lambda' - \lambda$. But in some cases, the combinatorics simplifies if we chose the weights differently, as will be illustrated below.

Suppose that $\mathcal{O}_w$ is $\frac{1}{2}$-BPS, and furthermore we can "partition the correlator into two flavors," i.e. choose the operators such that $\mathcal{O}_w = [\bar{z}_1^m z_2^n]$ while $\mathcal{O}_x = [z_1^k z_2^l]$ and $\mathcal{O}_y = [\bar{z}_1^{(k-m)} \bar{z}_2^{(l-n)}]$. Consider a diagram proportional to $\tilde{C}(x;y,w)$, see Figure 4.2(b). The sum of all such diagrams is symmetric in the color indices a and b at y (since the scalars at y used in this diagram have the same flavors, z_1^a and z_1^b), and symmetric in colors at w (since $\mathcal{O}_w$ is $\frac{1}{2}$-BPS, it is symmetric in the indices b and c). But it must be antisymmetric in the color indices a and c of the z_1^a and z_2^c leaving the interaction vertex (as this diagram comes with a factor of f^{apc}, see Figure 2.1). We conclude that all such diagrams cancel, and so $\tilde{\gamma}_x = 0$. In the same fashion, we conclude that $\tilde{\gamma}_y = 0$ as well, and together with $\tilde{\beta}_{yw} = \tilde{\beta}_{xw} = 0$ (as $\mathcal{O}_w$ is $\frac{1}{2}$-BPS), we find that $\beta = 0$ when $\mathcal{O}_x$ and $\mathcal{O}_y$ are any



Figure 4.3: Order g^2 corrections to correlators of the form (4.15): (a) and (b) includes a gauge boson exchange; (c) and (d) F-terms. Self energy contributions (not shown) also include a gauge boson exchange.

operators with protected scaling dimensions.

There is another type of three-point functions of BPS chiral primaries which receive no $\mathcal{O}(g^2)$ corrections by similar considerations. Consider a correlator (4.9) such that $\mathcal{O}_x$ is made of z_1 and z_2; $\mathcal{O}_y$ made of $\bar{z}_1$ and $\bar{z}_3$; and $\mathcal{O}_w$, made of $\bar{z}_2$ and z_3, i.e. a correlator of the form

$$\langle [z_1^m z_2^n](x)\ [\bar{z}_1^m \bar{z}_3^k](y)\ [z_3^k \bar{z}_2^n](w)\rangle \tag{4.15}$$

This correlator is "partitioned into three disjoint flavors." Order g^2 contributions to this three-point function are shown in Figure 4.3. There are no corrections proportional to any $\tilde{B}$-s since all lines within any rainbow carry the same flavor, so immediately $\tilde{\beta}_{xy} = \tilde{\beta}_{yw} = \tilde{\beta}_{xw} = 0$ and hence there are no $\mathcal{O}(g^2)$ corrections here, as well.

Finally, extremal three-point functions can be analyzed in a simple way. Here, the scaling dimension of one of the operators is equal to the sum of scaling dimensions of the other two.[5] Suppose that $\Delta_x + \Delta_y = \Delta_w$ in (4.9). At Born level, there are no $G(x, w)$ propagators, and so there are no corrections proportional to

---

[5] In general, $(n+1)$-point functions $\langle \mathcal{O}_0(x_0)\mathcal{O}_1(x_1)...\mathcal{O}_n(x_n)\rangle$ are called extremal if one of the scaling dimensions is the sum of all the others, $\Delta_0 = \Delta_1 + ... + \Delta_n$.



Figure 4.4: Order g^2 corrections to extremal correlators: (a) and (b) within a single peddle; (c) and (d) between the two peddles. Self energy contributions (not shown) and diagrams (a) and (c) are gauge dependent, while (b) and (d) diagrams arise from the F-terms.

$\tilde{B}(x, y)$, $\tilde{C}(x; y, w)$, or $\tilde{C}(y; x, w)$, see Figure 4.4. Together with the constraints (4.12-4.14), this determines $\tilde{\beta} = 0$ when the three (Lorentz scalar) operators inserted x, y, and w are arbitrary operators with protected scaling dimensions.

Another remark about extremal correlators is in order. As it is easy to see, one of the above group theory simplifications generalizes straightforwardly to extremal correlators of chiral primaries. Namely, if all operators except for one are $\frac{1}{2}$-BPS (and the remaining one is an arbitrary chiral primary), extremal correlators receive no order g^2 corrections.

4.5 Three-point functions of BPS operators

We are now ready to discuss correlators of three BPS chiral primaries. The simplest correlators $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\frac{1}{2}} \rangle$ (where each $\mathcal{O}_{\frac{1}{2}}$ stands for a $\frac{1}{2}$-BPS operator), were considered by the authors of [3], who found that three-point functions of $\frac{1}{2}$-BPS operators do not get corrected at order g^2, for any N. These are a special case of correlators of the form $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\frac{1}{2}} \mathcal{O}_{\text{BPS}} \rangle$, which we discussed in Section 4.4.2 (here $\mathcal{O}_{\text{BPS}}$ is an arbitrary BPS operator). Such three-point functions receive no

$\mathcal{O}(g^2)$ corrections by group theory reasoning.

4.5.1 Correlators $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$

Not all three-point functions of chiral primaries can be simplified using the results of Section 4.4.2, so occasionally we will have to actually compute some of the combinatorial coefficients. In this Section we will look at correlators of two $\frac{1}{4}$-BPS operators with one $\frac{1}{2}$-BPS operator.

4.5.1.1 $\langle \mathcal{O}_{[p,q,p]}(x) \bar{\mathcal{O}}_{[p,q,p]}(y) (\mathbf{tr}\, \bar{z} t z)(w) \rangle$

The simplest $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ three-point functions are of the form

$$\langle \mathcal{O}(x) \bar{\mathcal{O}}'(y) (\mathrm{tr}\, X^2)(w) \rangle \qquad (4.16)$$

where the $\frac{1}{2}$-BPS primary $\mathrm{tr}\, X^2$ is a scalar composite operator in the [0,2,0] of $SU(4)$. Group theory restricts the quantum numbers of operators which can have nontrivial three point functions. Tensoring $[p,q,p] \otimes [0,2,0]$ using Young diagrams of $SO(6)$ gives



$$\oplus \quad ... \qquad (4.17)$$

where in the first row there are no contractions (i.e. $SO(6)$ traces), only symmetrizations and antisymmetrizations; in the second row, one contraction; and in the third row, two contractions; the "..." stands for tableaux with more than two rows.[6] In terms of Dynkin labels, equation (4.17) reads

$$\begin{aligned}
[p,q,p]\otimes[0,2,0] &= [p,q+2,p]\oplus[p+1,q,p+1]\oplus[p+2,q-2,p+2]\\
&\oplus [p,q,p]\oplus[p+1,q-2,p+1]\oplus[p-1,q+2,p-1]\\
&\oplus [p,q-2,p]\oplus[p-1,q,p-1]\oplus[p-2,q+2,p-2]\\
&\oplus ... \qquad (4.18)
\end{aligned}$$

Now, the "..." stands for representations with $[r,s,r+2k]$ Dynkin labels with $k\neq 0$. Thus the only three-point functions of the form (4.16) which can possibly have a nonzero value are the extremal correlators

$$\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q-2,p]}(y)(\mathrm{tr}\,\bar{z}_1^2)(w)\rangle \qquad (4.19)$$

$$\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p-2,q+2,p-2]}(y)(\mathrm{tr}\,\bar{z}_2^2)(w)\rangle \qquad (4.20)$$

$$\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p-1,q,p-1]}(y)(\mathrm{tr}\,\bar{z}_1\bar{z}_2)(w)\rangle \qquad (4.21)$$

which correspond to those diagrams in (4.17) with zero or maximal number of contractions; and non-extremal correlators

$$\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)(\mathrm{tr}\,\bar{z}tz)(w)\rangle \qquad (4.22)$$

$$\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p-1,q+2,p-1]}(y)(\mathrm{tr}\,\bar{z}_2 z_1)(w)\rangle \qquad (4.23)$$

where t is a diagonal $SU(3)$ generator. All other correlators of the form (4.16) either vanish because the tensor product of irreps $[p,q,p]$ and $[0,2,0]$ does not contain $[r,s,r]$, or are related to the ones in (4.19-4.23).

Extremal three-point functions were discussed in Section 4.4.2, and were found to be protected at order g^2. The only correlators of the form $\langle \mathcal{O}\bar{\mathcal{O}}'\mathrm{tr}\,X^2\rangle$ we

---

[6] Tensoring representations in the manner of equation (4.17) gets messy for larger representations. Another method (of Berenstein-Zelevinsky triangles) is discussed in Appendix 4.9.1.



Figure 4.5: Nonvanishing Born diagrams for non-extremal 3-point functions $\langle \mathcal{O}_{[p,q,p]}\bar{\mathcal{O}}_{[p,q,p]}(\mathrm{tr}\,\bar{z}tz)\rangle$ (a-b); $\langle \mathcal{O}_{[p,q,p]}\bar{\mathcal{O}}_{[p-1,q+2,p-1]}(\mathrm{tr}\,\bar{z}_2 z_1)\rangle$ (c).

need to consider are those given by (4.22) and (4.23). However, the three-point functions of Figure 4.5(c), must in fact vanish: $\mathrm{tr}\,\bar{z}_2 z_1 = \frac{1}{2}\bar{z}_2^a z_1^a$ is diagonal in color indices, and hence the combinatorial factors for the Born graph of $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p-1,q+2,p-1]}(y)(\mathrm{tr}\,\bar{z}_2 z_1)(w)\rangle$ are proportional to the ones for the two-point function $\langle \mathcal{O}_{[p,q,p]}\bar{\mathcal{O}}_{[p-1,q+2,p-1]}\rangle = 0$. The same thing happens at order g^2, etc.[7] So correlators (4.23), although allowed by (4.18), are in fact forbidden by a combination of $SU(N)$ and $SU(4)$ group theory.

Correlators $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)(\mathrm{tr}\,\bar{z}tz)(w)\rangle$ are the only ones that remain to be considered. The contributing Born level diagrams are shown in Figure 4.5(a,b), and the $\mathcal{O}(g^2)$ graphs appear in Figure 4.6 (corrections to the scalar propagator are not shown, but are also present). Repeating the arguments of [3] from the $\frac{1}{2}$-BPS calculations, we see that the combinatorial structure of this three-point function $\langle \mathcal{O}\bar{\mathcal{O}}(\mathrm{tr}\,\bar{z}tz)\rangle$ is the same as that of the two-point function $\langle \mathcal{O}\bar{\mathcal{O}}\rangle$. At Born level, we find that

$$\begin{aligned}&\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)(\mathrm{tr}\,\bar{z}tz)(w)\rangle|_{\mathrm{free}}\\ &\qquad = \tfrac{1}{2}[(p+q)t_{11}+pt_{22}][\frac{G(x,w)G(y,w)}{G(x,y)}]\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle|_{\mathrm{free}} \qquad (4.24)\end{aligned}$$

---

[7] Explicitly, in Section 4.4.2 we saw that the $\mathcal{O}(g^2)$ part of this three-point function vanishes.



Figure 4.6: Order g^2 corrections to $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)(\operatorname{tr}\bar{z}tz)(w)\rangle$ with $t_{22}=0$: (a) within the rainbow ($i,j=1,2$ in the second diagram); (b) from the rainbow to X^2 (there are similar ones with the other leg of X^2 uncorrected).

At order g^2, the contributions proportional to $\tilde{B}$ and $\tilde{C}$ (diagrams (a1) and (b1) in Figure 4.6) have the same index structure, which in turn is identical to that of the two-point functions $\langle \mathcal{O}_{[p,q,p]}(x)\bar{\mathcal{O}}_{[p,q,p]}(y)\rangle$. Because $\operatorname{tr}\bar{z}_1 z_1$ is diagonal in color indices, its only effect on the combinatorics is to distinguish the pair of indices which go to $\mathcal{O}_w$ rather than stretch directly between $\mathcal{O}_x$ and $\mathcal{O}_y$.

There is a curious relation between the functions $\tilde{B}(x,y)$ and $\tilde{C}(x;y,w)$, which can be graphically expressed as[8]



(4.25)

This is a consequence of conformal invariance and nonrenormalization of the scaling dimension of $\operatorname{tr}X^2$. To see how this comes about, let $\mathcal{O}$ be an arbitrary (not necessarily BPS) scalar operators of same scaling dimension Δ; then

$$\langle \mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle = \frac{C}{(x-y)^{2\Delta}} \tag{4.26}$$

$$\langle \mathcal{O}(x)\bar{\mathcal{O}}(y)\operatorname{tr}X^2(w)\rangle = \frac{\tilde{C}}{(x-y)^{2\Delta}}\frac{G(x,w)G(y,w)}{G(x,y)} \tag{4.27}$$

in $d=4$; $\operatorname{tr}X^2 = \operatorname{tr}\bar{z}tz$ as in Equation (4.24). In other words, coordinate dependence (modulo the ratio of free scalar correlators) is the same, and the difference is just a constant factor. Assume now that $\mathcal{O}$ is constructed of only z_i-s (then $\bar{\mathcal{O}}$ is made of only $\bar{z}_j$-s). Then, the only $\mathcal{O}(g^2)$ contributions to the two-point function $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle$ are proportional to $\tilde{B}(x,y)$. Similarly, the $\mathcal{O}(g^2)$ corrections to the correlator $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\,\mathrm{tr}\,X^2(w)\rangle$ are proportional to $\tilde{B}(x,y)$ and to $[\tilde{C}(x;y,w)+\tilde{C}(y;x,w)]$. Index structure of these building blocks is the same (as discussed after equation 4.24), so

---

[8] The fact that $\tilde{C}(x;y,w)+\tilde{C}(y;x,w)-\tilde{B}(x,y)$ is just a constant was established in Section 4.3 by an explicit calculation. The value of this constant was also found there.

$$\begin{aligned}\frac{\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle|_{g^2}}{\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle|_{\text{free}}} &= \zeta\,\tilde{B}(x,y) &(4.28)\\ \frac{\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\,\mathrm{tr}\,X^2(w)\rangle|_{g^2}}{\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\,\mathrm{tr}\,X^2(w)\rangle|_{\text{free}}} &= \tfrac{1}{2}\zeta'\left[\tilde{C}(x;w,y)+\tilde{C}(y;w,x)+\tilde{B}(x,y)\right] &(4.29)\end{aligned}$$

The difference between $\tilde{B}(x,y)$ contributions to $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\,\mathrm{tr}\,X^2(w)\rangle$ and to $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle$ is that the pair of indices which go to w rather than stretch between x and y directly, is distinguished. Hence, $\zeta'=\zeta$ times some combinatorial factor.[9] As was discussed in Section 4.3, $\tilde{B}$ and $\tilde{C}$ have the form $\tilde{B}(x,y)=\tilde{a}\log\frac{(x-y)^2}{\epsilon^2}+\tilde{b}$, $\tilde{C}(x;y,w)=\tilde{a}'\log\frac{(x-y)^2(x-w)^2}{\epsilon^4}-\tilde{a}''\log\frac{(y-w)^2}{\epsilon^2}+\tilde{b}'$. By comparing (4.27) and (4.26), we see that expression (4.29) must have the same coordinate dependence as (4.28). This restricts $\tilde{a}'=\tilde{a}''=\frac{1}{2}\tilde{a}$, which reproduces the "winking cat" identity (4.25).

Finally, we can relate $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\,\mathrm{tr}\,X^2(w)\rangle$ to $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle$ by a Ward identity. As shown in [65], the ratio

$$\frac{\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)T_{\mu\nu}(0)\rangle}{\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle}=\frac{2\Delta}{3\pi^2}\frac{t_{\mu\nu}(\gamma)(x-y)^4}{x^4y^4} \tag{4.30}$$

depends on the scaling dimension Δ of the operator $\mathcal{O}$ (here, $\gamma=\frac{x}{x^2}-\frac{y}{y^2}$ and $t_{\mu\nu}(\gamma)=\frac{\gamma_\mu\gamma_\nu}{\gamma^2}-\frac{1}{4}\eta_{\mu\nu}$). Since the energy momentum tensor $T_{\mu\nu}$ is in the same $\mathcal{N}=4$ multiplet with $\mathrm{tr}\,X^2$, there is also nothing peculiar about the fact that $\tilde{C}_{12}/C_{12}$ can in general receive $\mathcal{O}(g^2)$ correction. This ratio also depends on Δ.

---

[9] In particular, if $\Delta=\Delta_0$ is not corrected, then neither $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\,\mathrm{tr}\,X^2(w)\rangle$ nor $\langle\mathcal{O}(x)\bar{\mathcal{O}}(y)\rangle$ get any one loop corrections.

4.5.1.2 General $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators

Three-point functions of two $\frac{1}{4}$-BPS operator and one $\frac{1}{2}$-BPS operator are similar to the ones described in Section 4.5.1.1. It suffices to consider a single three-point function (such that the Clebsch-Gordon coefficient[10] for these three vectors in the given irreps of $SU(4)$ is nonzero) for each set of three representations. Without loss of generality, we can choose a $[p, q, p]$ scalar composite $\mathcal{O}(x)$ to be made of only z-s; a $[r, s, r]$ scalar composite $\mathcal{O}'(y)$ to be made of only $\bar{z}$-s; and a $[0, k, 0]$ scalar composite $\operatorname{tr} X^{\alpha_1+\alpha_2}$ at w of the form

$$t_{i_1 \dots i_{\alpha_1}; j_1 \dots j_{\alpha_2}} \operatorname{str} z_{i_1} \dots z_{i_{\alpha_1}} \bar{z}_{j_1} \dots \bar{z}_{j_{\alpha_2}} \tag{4.31}$$

where $\alpha_1 + \alpha_2 = k$, and $t_{i_1 \dots i_{\alpha_1}; j_1 \dots j_{\alpha_2}}$ is the appropriate irreducible $SU(3)$ tensor (like in [3]). The correlators we are after are

$$\langle \mathcal{O}(x) \mathcal{O}'(y) (\operatorname{tr} X^{\alpha_1+\alpha_2})(w) \rangle \tag{4.32}$$

Position dependence of (4.32) is

$$[G(x,y)^{(2p+r)+(2r+s)-k} G(x,w)^{k+(2p+r)-(2r+s)} G(w,y)^{k-(2p+r)+(2r+s)}]^{1/2} \tag{4.33}$$

at Born level. The contributing free diagrams are similar to the ones shown in Figure 4.5; and $\mathcal{O}(g^2)$ diagrams, to those of Figure 4.6, but now there can be a different number of lines stretching between x and w and between w and y. Apart from the factor (4.33), the general $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlator (4.32) is given by

$$\alpha_{\text{free}} + \tilde{\beta}_{xy} \tilde{B}(x,y) + \tilde{\gamma}_x \tilde{C}(x; y, w) + \tilde{\gamma}_y \tilde{C}(y; x, w) + \mathcal{O}(g^4). \tag{4.34}$$

According to the discussion of Section 4.4, the remaining combinatorial coefficients vanish, $\beta_{xw} = \beta_{yw} = \gamma_w = 0$. Moreover, $\tilde{\gamma}_x = \tilde{\gamma}_y = -\tilde{\beta}_{xy}$ as follows from equations (4.12-4.14), so (4.34) reads

---

[10] By Wigner-Eckart theorem, for any three representations we only need to calculate one (nonvanishing) correlator of any representatives from these irreps.

$$\alpha_{\text{free}} - \tilde{\beta}_{xy}\left(\tilde{C}(x;y,w) + \tilde{C}(y;x,w) - \tilde{B}(x,y)\right) + \mathcal{O}(g^4) \tag{4.35}$$

Hence, we only need to verify that $\tilde{\beta}_{xy} = 0$.[11]

The simplifications we can use to deduce that $\tilde{\beta}_{xy} = 0$ without doing calculations, are discussed in Section 4.4.2. Extremal three-point functions are always easy to identify, and with the BPS primaries in representations $[p,q,p]$, $[r,s,r]$, and $[0,k,0]$, the restrictions on the scaling dimension translate into

$$2r+s = 2p+q+k, \quad 2p+q = 2r+s+k, \quad \text{or} \quad 2p+q+2r+s = k, \tag{4.36}$$

depending on which scaling dimension is the sum of the other two.

The "three flavor partition" boils down to being able to choose a single flavor (at Born level) for the lines between the two $\frac{1}{4}$-BPS operators, when the third operator is $\frac{1}{2}$-BPS. This is possible whenever

$$2r+s \leq k+q \quad \text{and} \quad 2p+q \leq k+s. \tag{4.37}$$

Alternatively, if we can choose the $\frac{1}{2}$-BPS operator $\mathcal{O}_w$ to be made of only $\bar{z}_1$-s and z_2-s; and the $\frac{1}{4}$-BPS operators as $\mathcal{O}_x$ of z_1-s and z_2-s, $\mathcal{O}_y$ of $\bar{z}_1$-s and $\bar{z}_2$-s, we get the "two flavor partition" of Section 4.4.2. This can happen if

$$k \leq q+s. \tag{4.38}$$

In all three cases (4.36), (4.37), (4.38) there are no $\mathcal{O}(g^2)$ corrections, as established in Section 4.4.2 using only $SU(4)$ group theory and conformal invariance arguments. However, there are allowed three-point functions of the form $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ where we can not choose irrep representatives in such a nice way.

---

[11] The expression multiplying $\tilde{\beta}_{xy}$ in (4.35) is a nonzero, renormalization scale independent constant. In Section 4.3, its value was computed to be $-\left(Y^2/32\pi^2\right)$.

Throughout the rest of this Section, we will concentrate on the $\frac{1}{4}$-BPS operators with $\Delta \leq 7$, constructed in Chapters 2 and 3. In particular, we will consider scalar composite operators in $SU(4)$ representations of the form $[p,q,p]$, with $2p+q \leq 7$. These are [2,0,2], [2,1,2], [2,2,2], [3,1,3], and [2,3,2]. We will take $\frac{1}{2}$-BPS (single trace) operators as whichever ones are allowed by group theory. Of the triple products of the form $[p,q,p] \otimes [r,s,r] \otimes [0,k,0]$ containing the singlet, most satisfy at least one of the simplifying constraints (4.36), (4.37), or (4.38).[12] The exceptions are $[2,0,2] \otimes [2,0,2] \otimes [0,2,0]$ and $[3,1,3] \otimes [3,1,3] \otimes [0,4,0]$.

Correlators $\langle \mathcal{O}_{[p,q,p]}(x) \bar{\mathcal{O}}_{[p,q,p]}(y) \operatorname{tr} X^2(w) \rangle$ were considered in Section 4.5.1.1, so the only three-point function we have to calculate is $\langle \mathcal{O}_{[3,1,3]} \bar{\mathcal{O}}_{[3,1,3]} \operatorname{tr} X^{2+2} \rangle$. Explicitly, we can choose the [3,1,3] scalar composite operators[13] as

$$\mathcal{O}_x = \sum_{j=1}^{4} C_x^j \mathcal{O}_j, \quad \mathcal{O}_y = \sum_{j=1}^{4} C_y^j \bar{\mathcal{O}}_j \quad \text{with} \quad \mathcal{O}_j \sim [z_1^4 z_2^2 z_3], \tag{4.39}$$

$$\mathcal{O}_w \sim [z_2^2 \bar{z}_2^2] - SO(6) \text{ traces.} \tag{4.40}$$

The free combinatorial factor for this three-point function is then

$$\begin{aligned}
\alpha_{\text{free}} \;=\; & \frac{(N^2-1)(N^2-2)}{41472N^2}(189540C_x^2C_y^2 - 4860C_x^2C_y^1N - \\
& 4860C_x^1C_y^2N - 131220C_x^4C_y^2N - 131220C_x^2C_y^4N + \\
& 360C_x^1C_y^1N^2 + 13500C_x^4C_y^1N^2 - 79380C_x^2C_y^2N^2 - \\
& 22680C_x^3C_y^2N^2 - 22680C_x^2C_y^3N^2 + 5184C_x^3C_y^3N^2 + \\
& 13500C_x^1C_y^4N^2 - 30780C_x^4C_y^4N^2 + 2700C_x^2C_y^1N^3 - \\
& 270C_x^3C_y^1N^3 + 2700C_x^1C_y^2N^3 + 43740C_x^4C_y^2N^3 - \\
& 270C_x^1C_y^3N^3 - 9720C_x^4C_y^3N^3 + 43740C_x^2C_y^4N^3 - \\
& 9720C_x^3C_y^4N^3 - 115C_x^1C_y^1N^4 - 2760C_x^4C_y^1N^4 +
\end{aligned}$$

---

[12] We omit the tedious details here. In order to find the allowed triple products, we used the method of BZ triangles, see Appendix 4.9.1. Then we just went through the list and checked if any of the conditions (4.36-4.38) applied.

[13] $\mathcal{O}_{1,\ldots,4}$ were discussed in Chapter 2. They are written out explicitly in Section 2.6.

$$\begin{aligned}
&13500C_x^2C_y^2N^4 + 4410C_x^3C_y^2N^4 + 4410C_x^2C_y^3N^4 - \\
&1332C_x^3C_y^3N^4 - 2760C_x^1C_y^4N^4 + 13680C_x^4C_y^4N^4 - \\
&450C_x^2C_y^1N^5 + 240C_x^3C_y^1N^5 - 450C_x^1C_y^2N^5 - \\
&4500C_x^4C_y^2N^5 + 240C_x^1C_y^3N^5 + 2340C_x^4C_y^3N^5 - \\
&4500C_x^2C_y^4N^5 + 2340C_x^3C_y^4N^5 - 15C_x^1C_y^1N^6 - \\
&990C_x^2C_y^2N^6 - 126C_x^3C_y^3N^6 - 1980C_x^4C_y^4N^6)
\end{aligned} \tag{4.41}$$

so $\langle \mathcal{O}_x \mathcal{O}_y \mathcal{O}_w \rangle \neq 0$ in general (and when $\mathcal{O}_x$ and $\mathcal{O}_y$ are $\frac{1}{4}$-BPS, in particular). We have also explicitly calculated[14] the $\mathcal{O}(g^2)$ combinatorial factor in (4.35):

$$\begin{aligned}
\tilde{\beta}_{xy} \;=\; & \frac{(N^2-1)(N^2-4)}{13824}(-10800C_x^2C_y^1 + 4320C_x^3C_y^1 - \\
&10800C_x^1C_y^2 - 259200C_x^4C_y^2 + 4320C_x^1C_y^3 + \\
&103680C_x^4C_y^3 - 259200C_x^2C_y^4 + 103680C_x^3C_y^4 - \\
&2025C_x^1C_y^1N - 27000C_x^4C_y^1N - 32400C_x^2C_y^2N + \\
&38880C_x^3C_y^2N + 38880C_x^2C_y^3N - 25920C_x^3C_y^3N - \\
&27000C_x^1C_y^4N - 129600C_x^4C_y^4N - 600C_x^2C_y^1N^2 + \\
&2940C_x^3C_y^1N^2 - 600C_x^1C_y^2N^2 + 50400C_x^4C_y^2N^2 + \\
&2940C_x^1C_y^3N^2 - 7200C_x^4C_y^3N^2 + 50400C_x^2C_y^4N^2 - \\
&7200C_x^3C_y^4N^2 + 175C_x^1C_y^1N^3 + 5400C_x^4C_y^1N^3 + \\
&7200C_x^2C_y^2N^3 - 7920C_x^3C_y^2N^3 - 7920C_x^2C_y^3N^3 + \\
&3888C_x^3C_y^3N^3 + 5400C_x^1C_y^4N^3 + 28800C_x^4C_y^4N^3 + \\
&600C_x^2C_y^1N^4 - 780C_x^3C_y^1N^4 + 600C_x^1C_y^2N^4 - \\
&780C_x^1C_y^3N^4 - 2880C_x^4C_y^3N^4 - 2880C_x^3C_y^4N^4 + \\
&50C_x^1C_y^1N^5 + 288C_x^3C_y^3N^5)
\end{aligned} \tag{4.42}$$

---

[14] This calculation was done using *Mathematica* and took about 200 hours. The choice of flavors (4.39-4.40) was optimal from the computational efficiency point of view.

If we choose the coefficients $(C_x^1, C_x^2, C_x^3, C_x^4)$ and $(C_y^1, C_y^2, C_y^3, C_y^4)$ independently from the set $\{(-\frac{12N}{N^2-2}, 1, -\frac{5}{N^2-2}, 0), (\frac{96}{N^2-4}, -\frac{4N}{N^2-4}, \frac{10N}{N^2-4}, 1)\}$, we recover $\tilde{\beta}_{xy} = 0$. This corresponds to taking $\mathcal{O}_x$ and $\mathcal{O}_y$ as the $\frac{1}{4}$-BPS chiral primaries found in Chapters 2 and 3, so there are no $\mathcal{O}(g^2)$ corrections in this case either.

4.5.2 Three-point functions of $\frac{1}{4}$-BPS operators

When all three operators are $\frac{1}{4}$-BPS, the arguments get more tedious. We will chose $2l + k \leq 2p + q \leq 2r + s$. The simplifications discussed in Section 4.4.2 applicable to correlators $\langle \mathcal{O}_{[p,q,p]} \mathcal{O}_{[r,s,r]} \mathcal{O}_{[l,k,l]} \rangle$ are: the extremality condition

$$2r + s = 2p + q + 2l + k; \tag{4.43}$$

and the "partition into three disjoint flavors" condition,[15]

$$\begin{cases} 2r + s & \leq & 2l + k + q \\ 2r + s & \leq & 2p + q + k \\ 2p + q & \leq & 2l + k + s \end{cases} \tag{4.44}$$

(all three inequalities have to be satisfied simultaneously). For example, (4.44) are true when all the $\frac{1}{4}$-BPS operators are in the $84 = [2, 0, 2]$ of $SU(4)$; take[16]

$$\mathcal{Y}(x) = \left\{ (\mathrm{tr}\, z_1^2)(\mathrm{tr}\, z_2^2) - (\mathrm{tr}\, z_1 z_2)(\mathrm{tr}\, z_1 z_2) \right\} + \frac{1}{N} \left\{ \mathrm{tr}\, [z_1, z_2]^2 \right\}, \tag{4.45}$$

$$\mathcal{Y}(y) = \left\{ (\mathrm{tr}\, \bar{z}_1^2)(\mathrm{tr}\, \bar{z}_3^2) - (\mathrm{tr}\, \bar{z}_1 \bar{z}_3)(\mathrm{tr}\, \bar{z}_1 \bar{z}_3) \right\} + \frac{1}{N} \left\{ \mathrm{tr}\, [\bar{z}_1, \bar{z}_3]^2 \right\}, \tag{4.46}$$

$$\mathcal{Y}(w) = \left\{ (\mathrm{tr}\, z_3^2)(\mathrm{tr}\, \bar{z}_2^2) - (\mathrm{tr}\, z_3 \bar{z}_2)(\mathrm{tr}\, z_3 \bar{z}_2) \right\} + \frac{1}{N} \left\{ \mathrm{tr}\, [z_3, \bar{z}_2]^2 \right\}. \tag{4.47}$$

The Born amplitude does not vanish[17] for $N > 2$, so we can't blame the lack of corrections on group theory,

$$\langle \mathcal{Y}(x)\, \mathcal{Y}(y)\, \mathcal{Y}(w) \rangle_{\text{free}} \;\propto\; (N^2 - 1)(N^2 - 4)(2N^2 - 15) \tag{4.48}$$

---

[15] Which just says that the number of scalars exchanged between each pair of $\mathcal{O}$-s is no larger than the length of the first column in the corresponding Young tableaux. There are three more inequalities, but they are satisfied trivially since we took $2l + k \leq 2p + q \leq 2r + s$.

[16] As shown in Appendix 4.9.2, such operators are in fact in the 84 of $SU(4)$.

[17] For $N \leq 2$ there are no $\frac{1}{4}$-BPS operators in the 84 of $SU(4)$.

and since $\langle \mathcal{Y}(x)\ \mathcal{Y}(y)\ \mathcal{Y}(w)\rangle$ is of the form (4.15), it receives $\mathcal{O}(g^2)$ corrections.

Of the allowed $\langle \mathcal{O}_{\frac{1}{4}}(x)\mathcal{O}_{\frac{1}{4}}(y)\mathcal{O}_{\frac{1}{4}}(w)\rangle$ three-point functions where each $\mathcal{O}_{\frac{1}{4}}$ is a scalar composite in a $[p,q,p]$ of $SU(4)$ with $2p+q\leq 7$, ten more satisfy (4.43) or (4.44).[18] For the remaining five correlators

$$\begin{aligned}
&\langle \mathcal{Y}_{[2,0,2]}(x)\mathcal{Y}_{[2,0,2]}(y)\mathcal{Y}_{[2,2,2]}(w)\rangle \\
&\langle \mathcal{Y}_{[2,0,2]}(x)\mathcal{Y}_{[2,1,2]}(y)\mathcal{Y}_{[2,3,2]}(w)\rangle \\
&\langle \mathcal{Y}_{[2,0,2]}(x)\mathcal{Y}_{[2,1,2]}(y)\mathcal{Y}_{[3,1,3]}(w)\rangle \\
&\langle \mathcal{Y}_{[2,0,2]}(x)\mathcal{Y}_{[3,1,3]}(y)\mathcal{Y}_{[2,3,2]}(w)\rangle \\
&\langle \mathcal{Y}_{[2,0,2]}(x)\mathcal{Y}_{[3,1,3]}(y)\mathcal{Y}_{[3,1,3]}(w)\rangle
\end{aligned} \tag{4.49}$$

we have to verify that there are no contributions proportional to any of the functions $\tilde{B}(x,y)$, $\tilde{B}(x,w)$, or $\tilde{B}(y,w)$. In fact, with $2l+k\leq 2p+q\leq 2r+s$,

$$2l+k\leq 2r+s+q \quad \text{and} \quad 2p+q\leq 2r+s+k \tag{4.50}$$

are automatically satisfied, so we can always choose the operators as[19]

$$\mathcal{Y}_{[l,k,l]}(x) \quad \sim \quad [\bar{z}_1^a\bar{z}_2^b z_3^e] \tag{4.51}$$

$$\mathcal{Y}_{[p,q,p]}(y) \quad \sim \quad [\bar{z}_1^c\bar{z}_2^d\bar{z}_3^e] \tag{4.52}$$

$$\mathcal{Y}_{[r,s,r]}(w) \quad \sim \quad [z_1^{r+s}z_2^r] \tag{4.53}$$

where $e\equiv\frac{1}{2}[(2l+k)+(2p+q)-(2r+s)]\leq l+k, p+q$; and integers a, b, c, d partition $r+s=a+c$, $r=b+d$. Then $\tilde{\beta}_{xy}=0$ since the operators exchanged between $\mathcal{Y}_x$ and $\mathcal{Y}_y$ all have the same flavor, and we only need to calculate $\tilde{\beta}_{xw}$ and $\tilde{\beta}_{yw}$. Details of these calculations are given in Appendix 4.9.3, and here we just quote the result: as in the cases considered so far, $\tilde{\beta}_{xy}=\tilde{\beta}_{xw}=\tilde{\beta}_{yw}=0$, and none of the three-point functions (4.49) receive any $\mathcal{O}(g^2)$ corrections.

---

[18] We used the method of BZ triangles (see Appendix 4.9.1) to find the allowed triple products.

[19] This choice of flavors was motivated by computational efficiency; our *Mathematica* calculations took more computer time with other flavor breakdowns.

4.6 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators in the large N limit

Like the two-point functions studied in Chapters 2 and 3, $\langle \mathcal{O}_x \mathcal{O}_y \mathcal{O}_w \rangle$ calculations get progressively more cumbersome as the representations of the $\mathcal{O}$-s become larger. In this Section we will calculate correlators of two $\frac{1}{4}$-BPS operators with one $\frac{1}{2}$-BPS operator, in the large N limit. The situation when all three operators are $\frac{1}{4}$-BPS is even less tractable, and we avoid it here.

4.6.1 Large N operators

We will use the $\frac{1}{4}$-BPS operators found in Chapters 2 and 3. Schematically, the special double and single trace operators can be written as

$$\mathcal{O}_{[p,q,p]} \sim$$



,

$$\mathcal{K}_{[p,q,p]} \sim$$



(4.54)

(each continuous group of boxes stands for an $SU(N)$ trace); explicit formulae for highest $SU(4)$ weight operators of this form are given in Chapter 2, Section 2.7.1. In the large N limit the linear combinations

$$\tilde{\mathcal{Y}}_{[p,q,p]} = \mathcal{K}_{[p,q,p]} + \mathcal{O}(N^{-2}) \tag{4.55}$$

$$\mathcal{Y}_{[p,q,p]} = \mathcal{O}_{[p,q,p]} - \frac{p(p+q)}{N} \mathcal{K}_{[p,q,p]} + \mathcal{O}(N^{-2}) \tag{4.56}$$

are eigenstates of the dilatations operator. $\mathcal{Y}_{[p,q,p]}$ have protected normalization and scaling dimension ($\Delta_{\mathcal{Y}} = 2p + q$) at order g^2, and were argued to be $\frac{1}{4}$-BPS.

We did not specify the $SU(4)$ weights of operators $\mathcal{O}_{[p,q,p]}$ and $\mathcal{K}_{[p,q,p]}$ in (4.54). The choice of weights will depend on the representations in the triple product $[p,q,p] \otimes [r,s,r] \otimes [0,k,0]$ in the following way. Assume $p \leq r$; then it is convenient to choose[20]

$$\begin{aligned} \mathcal{Y}_{[p,q,p]}(x) &\sim [\bar{z}_1^l z_2^n z_3^p] && (4.57) \\ \mathcal{Y}_{[r,s,r]}(y) &\sim [\bar{z}_1^m \bar{z}_2^n \bar{z}_3^p] && (4.58) \\ \mathcal{O}_{[0,k,0]}(w) &\sim [z_1^k] && (4.59) \end{aligned}$$

with $l \equiv \frac{1}{2}[(2p+q)+k-(2r+s)]$, $m \equiv k-l$ and $n \equiv p+q-l$. We will also assume that none of the simplifications (4.36-4.38) apply, since those cases were already discussed in Section 4.5.1.2.

4.6.2 $\langle \mathcal{KKO}_{\frac{1}{2}} \rangle_{\text{free}}$, $\langle \mathcal{KOO}_{\frac{1}{2}} \rangle_{\text{free}}$, $\langle \mathcal{OKO}_{\frac{1}{2}} \rangle_{\text{free}}$, and $\langle \mathcal{OOO}_{\frac{1}{2}} \rangle_{\text{free}}$

We can estimate the leading large N behavior of the combinatorial factors α_{free} and $\tilde{\beta}_{xy}$ using the "trace merging formula"

$$2\,(\operatorname{tr} A t^c)\,(\operatorname{tr} t^c B) = \operatorname{tr} AB - \tfrac{1}{N}\,(\operatorname{tr} A)\,(\operatorname{tr} B) \qquad (4.60)$$

where A and B are arbitrary $N \times N$ matrices and t^c are $SU(N)$ generators in the fundamental (sums on repeated indices are implied). With (4.60) and the expression for the quadratic Casimir[21]

$$T_r^c T_r^c = C_2(r)\,\mathbf{1} \qquad (4.61)$$

of $SU(N)$, we find for example that for $k \geq 2$,

$$(\operatorname{tr} t^{a_1} t^{a_2} ... t^{a_k})(\operatorname{tr} t^{a_k} ... t^{a_2} t^{a_1}) \;=\; \left(\frac{N}{2}\right)^k \left[1 + \mathcal{O}(1/N^2)\right] \qquad (4.62)$$

To have this large N behavior, generators in the two traces should appear in opposite order. When the generators are taken in any other order (except cyclic permutations inside the traces), such products are suppressed at least by $1/N^2$.

---

[20] With this choice of flavors, the manipulations of the following Sections are a simple generalization of the arguments of Section 2.7 of Chapter 2.

[21] For the adjoint and fundamental representations, $C_2(\text{adj}) = N$, $C_2(\text{fund}) = (\frac{N^2-1}{2N})$, $C(\text{adj}) = N$, $C(\text{fund}) = \frac{1}{2}$. See Section 2.9.2 for more details.

Calculations proceed along the same lines as in Chapter 2. We begin by considering correlators of the form $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle$ with the two $\mathcal{O}$-s in the same representation $[p,q,p]$. The leading contribution to α_{free} comes from terms like

$$(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_n})(\operatorname{tr} t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_l}t^{b_n}...t^{b_1})(\operatorname{tr} t^{c_p}...t^{c_1})\ (\operatorname{tr} t^{a_l}...t^{a_1}t^{d_l}...t^{d_1})$$
$$\sim \left(\frac{1}{2}\right)^2 N \left(\frac{N}{2}\right)^{2l-2+n} \times \left(\frac{N}{2}\right)^p = \left(\frac{1}{2}\right)\left(\frac{N}{2}\right)^{2l+n+p-1} \tag{4.63}$$

The factor of $(\frac{1}{2})^2$ comes about because we merge traces twice; the exponent $2l+n-2$ counts how many generators collapse using $t^c t^c \sim \frac{1}{2}N\mathbf{1}$; the extra factor of N is due to $\operatorname{tr}\mathbf{1} = N$; and finally $(N/2)^p$ is from contracting the traces of equal length containing the t^{c_i}-s. All remaining calculations of this and next Sections are analogous, and we won't spell things out as much.

If the representations of $[p,q,p]$ and $[r,s,r]$ are different, a similar situation occurs when for example $p = r+s$, i.e. $\mathcal{O}_{[p,q,p]}$ and $\mathcal{O}_{[r,s,r]}$ contain traces of equal length. Then we merge traces twice, and one set of traces collapses completely as in (4.62). Otherwise, we have to merge traces three times, so the leading contributions to $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{\text{free}}$ are of the form

$$(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_n})(\operatorname{tr} t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_m}t^{b_n}...)(\operatorname{tr}...t^{b_1}t^{c_p}...t^{c_1})\ (\operatorname{tr} t^{a_l}...t^{a_1}t^{d_m}...t^{d_1})$$
$$\begin{aligned} &\sim \left(\frac{1}{2}\right)\left(\frac{N}{2}\right)^{p+l+m+n-1} && \text{if } p=r \text{ or } p+q=r+s \\ & && \text{or } p=r+s \text{ or } r=p+q \\ &\sim \left(\frac{1}{2}\right)^3\left(\frac{N}{2}\right)^{p+l+m+n-3} && \text{otherwise} \end{aligned} \tag{4.64}$$

For the other three types of correlators, no pair of traces ever collapses completely, so the answers are more uniform. We find that the large N behavior of $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{\text{free}}$ is defined by terms like

$$(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_n}t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_m}t^{b_n}...t^{b_1}t^{c_p}...t^{c_1})\ (\operatorname{tr} t^{a_l}...t^{a_1}t^{d_m}...t^{d_1})$$
$$\sim \left(\frac{1}{2}\right)^2\left(\frac{N}{2}\right)^{p+l+m+n-2} N = \left(\frac{1}{2}\right)\left(\frac{N}{2}\right)^{p+l+m+n-1} \tag{4.65}$$

as we merge traces twice. Similarly, $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{\text{free}}$ scales as the terms

$$\begin{aligned}
&(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_n})(\operatorname{tr} t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_m}t^{b_n}...t^{b_1}t^{c_p}...t^{c_1})\ (\operatorname{tr} t^{a_l}...t^{a_1}t^{d_m}...t^{d_1}) \\
&\qquad \sim \left(\frac{1}{2}\right)^2 \left(\frac{N}{2}\right)^{p+l+m+n-2} \qquad (4.66)
\end{aligned}$$

since traces have to be merged three times now. The three-point functions $\langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{\text{free}}$ also have the leading large N dependence (4.66).

4.6.3 $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$, $\langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$, $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$, and $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$

Here there are no special cases to consider. We have to merge traces twice for $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$, three times for $\langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$ or $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$, and four times for $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$. The leading behavior of the $\tilde{\beta}_{xy}$ combinatorial coefficient for the three-point functions $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$ is the same as for terms of the form

$$\begin{aligned}
&(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_{n-1}}[t^{c_1},t^s]t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_m}t^{c_p}...t^{c_2}[t^{b_n},t^s]t^{b_n}...t^{b_1})\ (\operatorname{tr} t^{a_l}...t^{a_1}t^{d_m}...t^{d_1}) \\
&\qquad \sim \left(\frac{1}{2}\right)^2 \left(\frac{N}{2}\right)^{l+m-2} (\operatorname{tr} t^{b_1}...t^{b_{n-1}}[t^{c_1},t^s]t^{c_1}...t^{c_p}t^{c_p}...t^{c_2}[t^{b_n},t^s]t^{b_n}...t^{b_1}) \\
&\qquad \sim \left(\frac{1}{2}\right)^2 \left(\frac{N}{2}\right)^{p+l+m+n-4} (\operatorname{tr}[t^{c_1},t^s]t^{c_1}[t^{b_n},t^s]t^{b_n}) \\
&\qquad \sim \left(\frac{1}{2}\right) \left(\frac{N}{2}\right)^{p+l+m+n} \qquad (4.67)
\end{aligned}$$

which give the leading large N contributions to it. In the same fashion, the most significant terms in the correlators $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$ are

$$\begin{aligned}
&(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_{n-1}}[t^{c_1},t^s])(\operatorname{tr} t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_m}t^{c_p}...t^{c_2}[t^{b_n},t^s]t^{b_n}...t^{b_1})\ (\operatorname{tr} t^{a_l}...t^{a_1}t^{d_m}...t^{d_1}) \\
&\qquad \sim \left(\frac{1}{2}\right)^2 \left(\frac{N}{2}\right)^{p+l+m+n-1} \sim \langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2} \qquad (4.68)
\end{aligned}$$

while $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}}\rangle_{g^2}$ gets its leading N behavior from terms like

$$\begin{aligned}
&(\operatorname{tr} t^{a_1}...t^{a_l}t^{b_1}...t^{b_{n-1}}[t^{c_1},t^s])(\operatorname{tr} t^{c_1}...t^{c_p})\ (\operatorname{tr} t^{d_1}...t^{d_m}t^{c_p}...)(\operatorname{tr} ...t^{c_2}[t^{b_n},t^s]t^{b_n}...t^{b_1}) \\
&\quad \times(\operatorname{tr} t^{a_l}...t^{a_1}t^{d_m}...t^{d_1}) \\
&\qquad \sim \left(\frac{1}{2}\right)^3 \left(\frac{N}{2}\right)^{p+l+m+n-2} \qquad (4.69)
\end{aligned}$$

4.6.4 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators are protected

With just a little more work, we can find the ratios of the $\mathcal{O}(g^2)$ corrections to the three-point functions $\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, $\langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, and $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$. The argument proceeds along the same lines as in Chapter 2. Given a term with generators in a particular order, contributing to $\langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$, such as the one shown in (4.67), we know that a term with the same order of generators also gives a leading contribution to $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$ as in (4.68). However, cyclic permutations within the two traces (of length p and $p+q$) of $\mathcal{O}$, also contribute to $\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2}$ in the same amount as the term (4.68). Therefore,

$$\langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2} / \langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2} = \frac{p(p+q)}{N} + \mathcal{O}(N^{-3}) \equiv \beta \tag{4.70}$$

In the same fashion

$$\begin{aligned} \langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2} / \langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2} &= \frac{r(r+s)}{N} + \mathcal{O}(N^{-3}) \equiv \beta', && (4.71) \\ \langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2} / \langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{g^2} &= \frac{p(p+q)}{N} + \mathcal{O}(N^{-3}). && (4.72) \end{aligned}$$

Next consider the Born level correlators of Section 4.6.2. When a pair of traces collapses completely (see equations 4.64-4.66), we get

$$\langle \mathcal{Y}_{[p,q,p]} \mathcal{Y}_{[r,s,r]} \mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}} \sim \langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}} \sim N^{p+l+m+n-1} \tag{4.73}$$

Otherwise, the contributions add up to

$$\langle \mathcal{O}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}} - \beta \langle \mathcal{K}\mathcal{O}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}} - \beta' \langle \mathcal{O}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}} + \beta\beta' \langle \mathcal{K}\mathcal{K}\mathcal{O}_{\frac{1}{2}} \rangle_{\text{free}} \sim N^{p+l+m+n-3} \tag{4.74}$$

The terms in (4.74) are all of the same order and do not cancel. The factors of β and β' discussed above are still present, but there are other complications. First, the string of t^c-s can be inserted anywhere in the third trace in (4.64), and cyclic permutations of the t^b-s in the same trace give terms of the same order in N.

This results in an extra factor of $(r-p)^2$. Second, different terms in the sum over antisymmetrizations (as in equation (2.52), for example) contribute differently. The combinatorics is more involved, and we do not discuss this case in detail.

Bringing everything together, we see that the order g^2 corrections to the three-point function of the BPS operators in the large N limit add up to

$$\begin{aligned}\langle \mathcal{Y}_{[p,q,p]} \mathcal{Y}_{[r,s,r]} \mathcal{O}_{\frac{1}{2}} \rangle_{g^2} &\propto \left(\frac{1}{2}\right)\left(\frac{N}{2}\right)^{p+l+m+n-1} \tilde{B}(x,y) N \\ &\qquad \times \begin{pmatrix} -\beta \\ 1 \end{pmatrix}^t \begin{pmatrix} 1+\mathcal{O}(N^{-2}) & \beta' \\ \beta & \beta\beta' \end{pmatrix} \begin{pmatrix} -\beta' \\ 1 \end{pmatrix} \\ &= \left(\frac{1}{2}\right)\left(\frac{N}{2}\right)^{p+l+m+n-1} \tilde{B}(x,y) N \times \mathcal{O}(N^{-4}) \qquad (4.75)\end{aligned}$$

A comparison of (4.75) with (4.73) or (4.74) shows that order g^2 corrections to three-point functions of one $\frac{1}{2}$-BPS operator with two $\frac{1}{4}$-BPS operators vanish in the large N limit, within working precision.

4.7 Supergravity considerations

In the AdS/CFT correspondence, there is a duality mapping single trace $\frac{1}{2}$-BPS primary operators $\operatorname{tr} X^k$ of the SYM theory onto canonical supergravity fields, [1]. Given a set of such $\frac{1}{2}$-BPS primary operators, one can compute their two- and three-point functions in SYM. Two-point functions define the normalization of operators, and three-point functions probe the interactions between them. Independently, both the normalization of the SUGRA fields as well as their couplings, can be read off from the supergravity action (or supergravity equations of motion), [2]. So as a check of the AdS/CFT correspondence, one can compare the unambiguously defined three- and higher n-point functions of normalized $\frac{1}{2}$-BPS operators in SYM, with the correlators of the corresponding elementary excitations in supergravity, [2, 12, 17, 50].

We would like to proceed, in the same spirit, with the $\frac{1}{4}$-BPS chiral primaries of the $\mathcal{N}$=4 Super Yang Mills. We argued that these two- and three-point functions are independent of the SYM coupling constant (at least to order g^2), so it is reasonable to expect these correlators to agree with their dual AdS description. However, multiple trace operators do not correspond to any of the fields appearing in the supergravity action, so the discussion will be different than in the case of the previously studied $\frac{1}{2}$-BPS primary operators $\mathrm{tr}\, X^k$.

4.7.1 OPE definition of $\frac{1}{4}$-BPS chiral primaries

One of the ways to see $\frac{1}{4}$-BPS chiral primaries is to consider higher n-point correlators of $\frac{1}{2}$-BPS operators. For example, four-point functions of [0,2,0] operators reveal a pole corresponding to the exchange of a [2,0,2] operator with a protected dimension $\Delta = 4$, [7]. In general, the $\frac{1}{4}$-BPS primaries $\mathcal{Y}_{[p,q,p]}$ show up in

$$\langle \mathrm{tr}\, X^{(p+q)}(x) \mathrm{tr}\, X^p(x+\epsilon)\ \mathrm{tr}\, X^{(p+q)}(y)\ \mathrm{tr}\, X^p(w) \rangle \tag{4.76}$$

in the limit $\epsilon \to 0$, as the $[p,q,p]$ operators with the threshold value of scaling dimension $\Delta = 2p+q = \dim[\mathrm{tr}\, X^{(p+q)}] + \dim[\mathrm{tr}\, X^p]$. In other words, $\frac{1}{4}$-BPS chiral primaries can be defined by the OPE-s of $\frac{1}{2}$-BPS operators as

$$\mathcal{P}^{\Delta=2p+q}_{[p,q,p]} \left[\lim_{\epsilon \to 0} \mathrm{tr}\, X^{(p+q)}(x)\ \mathrm{tr}\, X^p(x+\epsilon) \right] = \sum_i c_i \mathcal{Y}_i^{[p,q,p]}(x) \tag{4.77}$$

Here, $\mathcal{P}^{\Delta}_{[p,q,p]}$ projects onto the $[p,q,p]$ representation of $SU(4)$, and eliminates operators with scaling dimension other than Δ (e.g. the non-chiral descendants with the same $SU(4)$ quantum numbers). Singular terms normally subtracted from an OPE such as (4.77), are automatically removed by applying $\mathcal{P}^{\Delta=2p+q}_{[p,q,p]}$.

On the other hand, one can see by calculating three-point correlators that all $\frac{1}{4}$-BPS primary operators $\mathcal{Y}_i^{[p,q,p]}$ are present in the OPE (4.77). It appears that for general N, there is no canonical definition of the special $\mathcal{Y}_{[p,q,p]}$ that is a linear combination of the single and double-trace scalar composite $[p, q, p]$ operators only. However, this $\mathcal{Y}_{[p,q,p]}$ dominates in the $N \to \infty$ limit. For large N, all other terms in the right hand side of (4.77) are suppressed by at least a factor of $1/N$, and the predominantly double-trace $\frac{1}{4}$-BPS chiral primary operator

$$\mathcal{Y}_{[p,q,p]} = \left(\begin{array}{l} \boxed{}\boxed{\;\cdots\;}\boxed{}\boxed{}\boxed{\cdots}\boxed{} \\ \underbrace{\boxed{}\boxed{\;\cdots\;}\boxed{}}_{p} \qquad q \end{array} \right) + \mathcal{O}(1/N) \tag{4.78}$$

is uniquely defined by the OPE of $\frac{1}{2}$-BPS primaries.

When translated into the SUGRA language, the definition (4.77) implies that $\frac{1}{4}$-BPS primary operators of SYM should be thought of as threshold bound states of elementary SUGRA excitations. A threshold bound state is a state whose mass is precisely equal to the sum of the masses of all its constituents, and thus occurs at the lower end of the spectrum. Any bound state of BPS states which is itself BPS is automatically a threshold bound state. A familiar example is provided by an assembly of like sign charged Prasad-Sommerfield magnetic monopoles, whose classical static solution forms a threshold bound state of monopole constituents.

4.7.2 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$ and $\langle \mathcal{O}_{\frac{1}{2}} \mathcal{O}'_{\frac{1}{2}} \mathcal{O}_{\frac{1}{4}} \rangle$ correlators

We are now going to illustrate the consistency of this dictionary. Specifically, we will look at two- and three-point functions involving $\frac{1}{2}$- and $\frac{1}{4}$-BPS operators in the large N limit, in $\mathcal{N}$=4 SYM. Then we will compare these correlators with their dual supergravity description.

The normalization of $\frac{1}{2}$- and $\frac{1}{4}$-BPS operators comes from their two-point functions, whose leading large N behavior was found in Chapter 2 (also see [2]):

$$\langle \operatorname{tr} X^q(x) \; \operatorname{tr} X^q(y) \rangle \sim \frac{N^q}{(x-y)^{2q}} \tag{4.79}$$

$$\langle \mathcal{Y}_{[p,q,p]}(x) \; \bar{\mathcal{Y}}^{[p,q,p]}(y) \rangle \sim \frac{N^{(2p+q)}}{(x-y)^{2(2p+q)}} \tag{4.80}$$

times some N-independent factors which we omit.

The simplest three point functions involving $\frac{1}{2}$- and $\frac{1}{4}$-BPS operators are of the form $\langle \mathcal{O}_{\frac{1}{2}}(x)\mathcal{O}'_{\frac{1}{2}}(y)\mathcal{O}_{\frac{1}{4}}(w)\rangle$. If the $SU(N)$ traces collapse completely (in which case $\langle \mathcal{O}_{\frac{1}{2}}\mathcal{O}'_{\frac{1}{2}}\mathcal{O}_{\frac{1}{4}}\rangle$ are extremal), the normalized three point-functions are then

$$\frac{1}{\sqrt{N^{(2p+q)+(p+q)+p}}}\langle \mathrm{tr}\, X^{(p+q)}(x)\ \mathrm{tr}\, X^{p}(y)\ \mathcal{Y}_{[p,q,p]}(w)\rangle \sim 1 \tag{4.81}$$

from a field theory calculation; the space-time coordinate dependence is fixed by conformal invariance, so we will not exhibit it anymore. If the traces do not collapse completely, the correlator is suppressed by $1/N^2$ (see the discussion around equations (4.63-4.64) of Section 4.6), and

$$\frac{1}{\sqrt{N^{(2p+q)+(k+l)+k}}}\langle \mathrm{tr}\, X^{(k+l)}(x)\ \mathrm{tr}\, X^{k}(y)\ \mathcal{Y}_{[p,q,p]}(w)\rangle \sim \frac{1}{N^2} \tag{4.82}$$

whenever $k \neq p$ of $l \neq q$. All this matches nicely with the corresponding supergravity diagrams:



(4.83)

Leading AdS diagrams for (a) equation (4.80); (b) equation (4.81); (c) equation (4.82). Each cubic bulk interaction vertex goes like $1/N$.

We denoted $\frac{1}{2}$-BPS primaries by “$\bullet$”; and the predominantly double trace $\frac{1}{4}$-BPS primaries which arise from bringing two $\frac{1}{2}$-BPS operators together by “$\overset{\frown}{\bullet\bullet}$”.

There are also AdS diagrams with quartic interactions in the bulk, which have the same large N dependence as (4.83c); we will not show these.

4.7.3 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}'_{\frac{1}{4}} \mathcal{O}_{\frac{1}{2}} \rangle$ correlators

Other three point functions involving $\frac{1}{4}$-BPS as well as $\frac{1}{2}$-BPS operators can be analyzed similarly. Whenever traces of the SYM operators do not collapse completely, the supergravity counterparts of such correlators have extra bulk interaction vertices. The leading dependence of such correlators is then suppressed by the corresponding power of $1/N$. For example, correlators of the form $\langle \mathcal{O}_{\frac{1}{4}}(x) \mathcal{O}'_{\frac{1}{4}}(y) \mathcal{O}_{\frac{1}{2}}(w) \rangle$, discussed in Section 4.6, behave like

$$\frac{1}{\sqrt{N^{(2p+q)+(2r+s)+k}}} \langle \mathcal{Y}_{[p,q,p]} \mathcal{Y}_{[r,s,r]} \mathrm{tr}\, X^k \rangle \sim \begin{cases} 1/N & \text{(a) if one pair of traces} \\ & \text{collapses completely} \\ 1/N^3 & \text{(b) otherwise} \end{cases} \tag{4.84}$$

From the AdS point of view, this difference is explained by the following diagrams



(4.85)

AdS description of equation (4.84).

4.7.4 $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}'_{\frac{1}{4}} \mathcal{O}''_{\frac{1}{4}} \rangle$ correlators

Similar arguments show that when all operators are $\frac{1}{4}$-BPS, the normalized three-point functions are

$$\frac{1}{\sqrt{N^{(2p+q)+(2r+s)+(2l+k)}}} \langle \mathcal{Y}_{[p,q,p]} \mathcal{Y}_{[r,s,r]} \mathcal{Y}_{[l,k,l]} \rangle \sim \begin{cases} 1 & \text{(a) if all traces} \\ & \text{collapse pairwise} \\ 1/N^2 & \text{(b) if only one pair} \\ & \text{of traces collapses} \\ 1/N^4 & \text{(c) otherwise} \end{cases} \tag{4.86}$$

and the corresponding AdS diagrams



$$(4.87)$$

AdS description of equation (4.86).

show the correct leading large N behavior.

4.7.5 Detailed agreement between SYM and AdS

Unlike the $\frac{1}{2}$-BPS calculations (e.g. [2]), this study does not provide a new independent check or application of the AdS/CFT correspondence. On the one hand, the definition of the predominantly double-trace $\frac{1}{4}$-BPS operators in the SYM theory (in the large N limit) is based on the OPE of $\frac{1}{2}$-BPS primaries. On the other hand, AdS correlators of the duals of the $\frac{1}{4}$-BPS operators (bound states of elementary SUGRA excitations) are defined by the corresponding correlators of primary supergravity fields. Therefore, SYM correlators involving $\frac{1}{4}$-BPS operators agree by construction with their SUGRA counterparts.

This is especially clear in the cases show in diagrams (4.83a,b), (4.85a), and (4.87a). To leading order in N, these two- and three-point functions of $\frac{1}{4}$-BPS scalar composite operators are expressed in terms of the previously studied two- and three-point functions of $\frac{1}{2}$-BPS chiral primary operators.

4.8 Conjectures

Let us summarize what has been done so far. First, $\frac{1}{4}$-BPS primary operators were identified in Chapters 2 and 3: for general representations $[p, q, p]$ in the large N limit; and for general N in the case when $2p+q \leq 7$. Second, three-point functions involving $\frac{1}{2}$-BPS operators as well as several infinite families of $\frac{1}{4}$-BPS operators were considered in this Chapter, also for arbitrary N. It was found that there are no $\mathcal{O}(g^2)$ corrections to such correlators. Next, all three-point functions involving the $\frac{1}{4}$-BPS primaries with $2p+q \leq 7$ were computed for general N, and were shown to be protected at order g^2. In the large N approximation, three point functions involving two $\frac{1}{4}$-BPS primaries and one $\frac{1}{2}$-BPS primary were shown to receive no $\mathcal{O}(g^2)$ corrections, for general representations of the operators involved. Finally, we presented AdS considerations which reproduced many features of the CFT two- and three-point functions in the large N limit.

Collecting the non-renormalization effects established above generates strong evidence for a number of natural conjectures, which we now state:

(1) We conjecture that on the CFT side, for every $[p,q,p]$ representation of $SU(4)$ and arbitrary N, there are $\frac{1}{4}$-BPS chiral primaries. Within each $[p,q,p]$, one of these operators is a linear combination of double and single trace scalar composites only; the other $\frac{1}{4}$-BPS chiral primaries in $[p,q,p]$ also involve operators with higher numbers of traces.

(2) We speculate that two–point functions of $\frac{1}{4}$-BPS operators, as well as three-point functions involving $\frac{1}{2}$-BPS and $\frac{1}{4}$-BPS operators, do not depend on the coupling g^2 of $\mathcal{N}$=4 SYM. This non-renormalization also persist for all N, and is not just a large N approximation.

(3) One of the group theory arguments of Section 4.4.2, and the analysis of Section 4.6, generalize straightforwardly to extremal correlators, i.e. $(n+1)$-point functions of the form $\langle \mathcal{O}_0(x_0)\mathcal{O}_1(x_1)...\mathcal{O}_n(x_n)\rangle$ with $\Delta_0 = \Delta_1 + ... + \Delta_n$. So do the AdS considerations of Section 4.7. Therefore, we conjecture that arbitrary extremal correlators of $\frac{1}{2}$- and $\frac{1}{4}$-BPS chiral primaries are also protected.

4.9 Appendix

4.9.1 $[p,q,p]\otimes[r,s,r]\otimes[0,k,0]$ and BZ triangles

Tensoring irreducible representations using Young tableaux can get quite tedious. Berenstein-Zelevinsky (BZ) triangles [66] provide a powerful way to calculate the multiplicity of the scalar representation[22] in $\lambda\otimes\mu\otimes\nu$. We will discuss the construction for $SU(3)$ and $SU(4)$, the generalization to higher $SU(N)$ (but not to other Lie algebras, which is not currently known) being straightforward.

For $SU(3)$, the triangles are constructed according to the following rules:

$$\begin{array}{ccccccc} & & & m_{13} & & & \\ & & n_{12} & & l_{23} & & \\ & m_{23} & & & & m_{12} & \\ n_{13} & & l_{12} & & n_{23} & & l_{13} \end{array} \tag{4.88}$$

where the nine non-negative integers l_{ij}, m_{ij}, n_{ij} are related to the Dynkin labels (λ_1,λ_2), (μ_1,μ_2), (ν_1,ν_2) of the highest weights of the three representations by

$$\begin{array}{lll} m_{13}+n_{12}=\lambda_1 & n_{13}+l_{12}=\mu_1 & l_{13}+m_{12}=\nu_1 \\ m_{23}+n_{13}=\lambda_2 & n_{23}+l_{13}=\mu_2 & l_{23}+m_{13}=\nu_2 \end{array} \tag{4.89}$$

They must further satisfy the so-called hexagon conditions

$$\begin{aligned} n_{12}+m_{23}&=m_{12}+n_{23} \\ l_{12}+m_{23}&=m_{12}+l_{23} \\ l_{12}+n_{23}&=n_{12}+l_{23} \end{aligned} \tag{4.90}$$

This means that the length of opposite sides in the hexagon formed by n_{12}, l_{23}, m_{12}, n_{23}, l_{12}, and m_{23} in (4.88) are equal, the length of a segment being the sum of its two vertices.

---

[22] It is conventional to choose ν^* instead of ν for the third weight.

The number of such triangles gives the multiplicity $\mathcal{N}_{\lambda\mu\nu}$; if it is not possible to construct such a triangle, ν^* does not occur in the tensor product $\lambda \otimes \mu$.

The integers in the BZ triangles have the following origin. Each pair of indices ij, $i < j$, on the labels of the triangle is related to a positive root of $SU(3)$. For $SU(N)$, positive roots can be written as $\epsilon_i - \epsilon_j$, $1 \leq i \leq j \leq N$, in terms of orthonormal vectors ϵ_i in $\mathbf{R}^N$.

The triangle encodes three sums of positive roots:

$$\begin{aligned} \mu + \nu - \lambda^* &= \textstyle\sum_{i<j} l_{ij}(\epsilon_i - \epsilon_j) \\ \nu + \lambda - \mu^* &= \textstyle\sum_{i<j} m_{ij}(\epsilon_i - \epsilon_j) \\ \lambda + \mu - \nu^* &= \textstyle\sum_{i<j} n_{ij}(\epsilon_i - \epsilon_j) \end{aligned} \tag{4.91}$$

The hexagon relations (4.90) can be seen as consistency conditions for these three expansions.

For $SU(4)$, the BZ triangles are defined in a similar way, in terms of

$$\begin{array}{ccccccccccc} & & & & & m_{14} & & & & & \\ & & & & n_{12} & & l_{34} & & & & \\ & & & m_{24} & & & & m_{13} & & & \\ & & n_{13} & & l_{23} & & n_{23} & & l_{24} & & \\ & m_{34} & & & & m_{23} & & & & m_{12} & \\ n_{14} & & l_{12} & & n_{24} & & l_{13} & & n_{34} & & l_{14} \end{array} \tag{4.92}$$

eighteen non-negative integers, related to the Dynkin labels by

$$\begin{array}{lll} m_{14} + n_{12} = \lambda_1 & n_{14} + l_{12} = \mu_1 & l_{14} + m_{12} = \nu_1 \\ m_{24} + n_{13} = \lambda_2 & n_{24} + l_{13} = \mu_2 & l_{24} + m_{13} = \nu_2 \\ m_{34} + n_{14} = \lambda_3 & n_{34} + l_{14} = \mu_3 & l_{34} + m_{14} = \nu_3 \end{array} \tag{4.93}$$

Furthermore, an $SU(4)$ BZ triangle has three hexagons[23]

$$\begin{array}{ccc} n_{12}+m_{24}=m_{13}+n_{23} & n_{13}+l_{23}=l_{12}+n_{24} & l_{24}+n_{23}=l_{13}+n_{34} \\ n_{12}+l_{34}=l_{23}+n_{23} & n_{13}+m_{34}=n_{24}+m_{23} & n_{23}+m_{23}=m_{12}+n_{34} \\ m_{24}+l_{23}=l_{34}+m_{13} & m_{34}+l_{12}=l_{23}+m_{23} & l_{13}+m_{23}=l_{24}+m_{12} \end{array} \tag{4.94}$$

As an application, consider $\nu=[0,k,0]\subset[p,q,p]\otimes[r,s,r]=\lambda\otimes\mu$ of $SU(4)$; here all representations are self-conjugate. The restrictions on the l_{ij}, m_{ij}, n_{ij} (these integers must all be non-negative) imply that the entries of the BZ triangle are actually

$$\begin{aligned} m_{14}=l_{14}=m_{12}=l_{34} &= 0, \\ n_{12} &= p, \\ n_{23} &= n_{14}, \\ n_{34} &= r, \\ l_{23}=m_{34} &= p-n_{14}, \\ l_{12}=m_{23} &= r-n_{14}, \\ l_{13} &= \tfrac{1}{2}(s+k-(2p+q)+2n_{14}), \\ m_{24} &= \tfrac{1}{2}(q+k-(2r+s)+2n_{14}), \\ n_{13} &= \tfrac{1}{2}(q-k+(2r+s)-2n_{14}), \\ n_{24} &= \tfrac{1}{2}(s-k+(2p+q)-2n_{14}), \\ m_{13} &= \tfrac{1}{2}((2p+q)+k-(2r+s)), \\ l_{24} &= \tfrac{1}{2}((2r+s)+k-(2p+q)). \end{aligned} \tag{4.95}$$

All entries thus depend on a single parameter n_{14} which is subject to restrictions $0\le n_{14}\le p,r,\frac{1}{2}(p+r-k)$; plus we get further constraints $k\ge|(2p+q)-(2r+s)|$,

---

[23] The $SU(N)$ generalization is straightforward; the BZ triangles are built out of three corner vertices and $\frac{1}{2}(N-1)(N-2)$ hexagons.

$p+q \geq r$, and $r+s \geq p$, etc.

Now, recall that $SO(6) \sim SU(4)$, and all our operators are in fact made of the scalars ϕ^I $(I = 1, ..., 6)$ which are in the fundamental $\mathbf{6}$ of $SO(6)$. In terms of the Young diagrams for $SO(6)$, the representations involved are partitioned as

$$\begin{array}{|c|c|c|c|}\hline n_{14} & m_{34} & m_{24} & n_{13} \\ \hline n_{14} & m_{34} & \multicolumn{2}{c}{q} \\ \cline{1-2} \multicolumn{2}{c}{p} \end{array} \qquad \begin{array}{|c|c|c|c|}\hline n_{14} & m_{23} & l_{13} & n_{24} \\ \hline n_{14} & m_{23} & \multicolumn{2}{c}{s} \\ \cline{1-2} \multicolumn{2}{c}{r} \end{array}$$

$$\begin{array}{|c|c|c|c|}\hline m_{34} & m_{24} & m_{23} & l_{13} \\ \hline \multicolumn{4}{c}{k} \end{array} \tag{4.96}$$

An especially convenient decomposition is when the $[0, k, 0]$ state is made up of say only 1-s and $\bar{2}$-s. In which case, by symmetry in the vertices, there will be no contributions proportional to $\tilde{C}$ provided only two flavors are involved in the diagram. Unfortunately, this can be achieved only when $s + q \geq k$.

Alternatively, we can take a $[p, q, p]$ state made up of $n_{14} + m_{34} + m_{24}$ 1-s and $n_{14} + m_{34} + n_{13}$ 2-s; $[r, s, r]$ state made up of $n_{14} + m_{23} + l_{13}$ $\bar{1}$-s and $n_{14} + m_{23} + n_{24}$ $\bar{2}$-s; $[0, k, 0]$ state made up of $m_{34} + m_{24}$ $\bar{1}$-s and $m_{23} + l_{13}$ 1-s, minus contractions. Unlike the previous decomposition, this one works for any $[0, k, 0] \otimes [p, q, p] \otimes [r, s, r]$ containing the singlet.

4.9.2 Partitioning a tableau into 2 flavors

It is often convenient to choose the operators to have only two distinct flavors. Here we shall see that it can always be done.

Consider an operator in the $[p, q, p]$ of $Gl(6)$ made of n_1 1-s and n_2 2-s, to be concrete. We have the following constraints: $p \leq n_2, n_1 \leq p + q$. This state can be assigned an $SU(4)$ weight $w = n_1(0, 1, 0) + n_2(1, -1, 1) = (n_2, n_1 - n_2, n_2)$.

Next, we project this onto the $SU(3) \times U(1)$; for example, we can choose $1 \to 1 + \bar{1}$, $2 \to 2 + \bar{2}$. Then w contains terms with b 1-s, $n_1 - b$ $\bar{1}$-s, c 2-s, $n_2 - c$

$\bar{2}$-s; these have weights $w'_{b,c} = (n_2 - n_1 + 2(b-c), 2c - n_2)^{2(b+c)-(n_1+n_2)}$.

To make an irrep of $Gl(6)$ into one of $SO(6)$, we must subtract traces. Since traces have weight zero, contributions with n contractions instead of 1 and m instead of 2, are equivalent to $n'_1 = n_1 - 2n$, $n'_2 = n_2 - 2m$. They are projected onto $w'^{n,m}_{b,c} = (n_2 - n_1 + 2(b+n) - 2(c+m), 2(c+m) - n_2)^{2(b+n+c+m)-(n_1+n_2)}$. We see that for fixed n_1 and n_2, $w'^{n,m}_{b,c} = w'^{\tilde{n},\tilde{m}}_{\tilde{b},\tilde{c}}$ iff $b + n = \tilde{b} + \tilde{n}, c + m = \tilde{c} + \tilde{m}$.

We are interested in having $b = n_1$, $c = n_2$, for example; then $w'_{n_1,n_2} = (n_1 - n_2, n_2)^{n_1+n_2}$. In order for $w'^{n,m}_{b,c}$ to have the same weight we must have $n_1 - n \geq b + n = n_1$, $n_2 - m \geq c + m = n_2$, or $m = n = 0$; likewise, traces also do not contribute to the projection onto $b = n_1$, $c = 0$.

This means that $[p, q, p]$ states of $SO(6) \sim SU(4)$ which consist of n_1 of any ϕ_a and n_2 of any other ϕ_b minus various contractions, project onto states in irreducible representations of $SU(3) \times U(1)$, ones containing n_1 z_a-s and n_2 z_b-s or n_1 z_a-s and n_2 $\bar{z}_b$-s, etc. *without having to subtract any traces*.

4.9.3 Details of $\langle \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \mathcal{O}_{\frac{1}{4}} \rangle$ calculations

In Section 4.5.2 we needed to explicitly calculate several three-point functions $\langle \mathcal{O}_{[l,k,l]}(x) \mathcal{O}_{[p,q,p]}(y) \mathcal{O}_{[r,s,r]}(w) \rangle$ of $\frac{1}{4}$-BPS operators. The flavor breakdown is discussed in Section 4.5.2, see equations (4.51-4.53). For the five cases of (4.49), we discuss the combinatorial coefficients multiplying the Born diagram, as well as the ones in front of $\tilde{B}(x, w)$ and $\tilde{B}(w, y)$.

When operators $\mathcal{O}_w$ are properly symmetrized, we can mark the z_1-s exchanged between $\mathcal{O}_w$ and $\mathcal{O}_x$ separately from the z_1-s exchanged between $\mathcal{O}_w$ and $\mathcal{O}_y$. As far as the combinatorial factors α_{free}, $\tilde{\beta}_{xw}$ and $\tilde{\beta}_{yw}$ are concerned, the difference is just a multiplicative factor. This would be equivalent to calculating the three-point functions with

$$\begin{aligned}
\mathcal{O}_{[l,k,l]}(x) &\sim [z_1^a z_2^b z_3^e] && (4.97)\\
\mathcal{O}_{[p,q,p]}(y) &\sim [\bar{z}_1^c \bar{z}_2^d \bar{z}_3^e] && (4.98)\\
\mathcal{O}_{[r,s,r]}(w) &\sim [\bar{z}_1^a z_1^c \bar{z}_2^b z_2^d] && (4.99)
\end{aligned}$$

rather than (4.51-4.53) instead,[24] with the same $e \equiv \frac{1}{2}[(2l+k)+(2p+q)-(2r+s)] \leq l+k, p+q$, and integers a, b, c, d partitioning $r+s=a+c$, $r=b+d$. This simplifies the calculations dramatically.

For operators in the [2,0,2] or [2,1,2] representations, we chose the $\frac{1}{4}$-BPS operator from the beginning. In the other cases, several $\frac{1}{4}$-BPS chiral primaries exist in each representation, so instead we choose the operators as $\mathcal{O}_w = \sum_j C_w^j \mathcal{O}_j$ for example (see Sections 2.4, 2.6 and 3.2 for the definitions). With $\mathcal{O}_x$, $\mathcal{O}_y$, $\mathcal{O}_w$ as in (4.97-4.99), we list the representations and choices of flavors below:

$$\begin{aligned}
&[2,0,2]\otimes[2,0,2]\otimes[2,2,2]: && \langle [z_1 z_2^2 z_3]_x [\bar{z}_1 \bar{z}_2^2 \bar{z}_3]_y [z_1 z_2^2 \bar{z}_1 \bar{z}_2^2]_w \rangle && (4.100)\\
&[2,0,2]\otimes[2,1,2]\otimes[2,3,2]: && \langle [z_1 z_2^2 z_3]_x [\bar{z}_1 \bar{z}_2^3 \bar{z}_3]_y [z_1 z_2^3 \bar{z}_1 \bar{z}_2^2]_w \rangle && (4.101)\\
&[2,0,2]\otimes[2,1,2]\otimes[3,1,3]: && \langle [z_1 z_2^2 z_3]_x [\bar{z}_1^2 \bar{z}_2^2 \bar{z}_3]_y [z_1^2 z_2^2 \bar{z}_1 \bar{z}_2^2]_w \rangle && (4.102)\\
&[2,0,2]\otimes[3,1,3]\otimes[2,3,2]: && \langle [z_1^2 z_3^2]_x [\bar{z}_1^2 \bar{z}_2^3 \bar{z}_3^2]_y [z_1^2 z_2^3 \bar{z}_1^2]_w \rangle && (4.103)\\
&[2,0,2]\otimes[3,1,3]\otimes[3,1,3]: && \langle [z_1^2 z_3^2]_x [\bar{z}_1^2 \bar{z}_2^3 \bar{z}_3^2]_y [z_1^2 z_2^3 \bar{z}_1^2]_w \rangle && (4.104)
\end{aligned}$$

The coefficients α_{free} and $\tilde{\beta}_{yw}$ take a long time to compute, and are not particularly illuminating. For example, in the case of $[2,0,2]\otimes[3,1,3]\otimes[3,1,3]$ we find (after 2430 hrs of a *Mathematica* computation on a Pentium-III with 1.4MHz speed)

$$\alpha_{\text{free}} = \frac{(N^2-1)(N^2-4)}{1036800N^2}(2332800C_w^2C_y^2 - 2332800C_y^2C_w^4N - \qquad (4.105)$$

---

[24] As written in (4.99), $\mathcal{O}_w$ is not even a $[r,s,r]$ operator; we need to subtract $SO(6)$ traces. But when calculating whether the three coefficients α_{free}, $\tilde{\beta}_{xw}$ and $\tilde{\beta}_{yw}$ are zero or not, the answers are the same as if we had done it properly.

$$
\begin{aligned}
& 2332800C_w^2C_y^4N + 10800C_w^1C_y^1N^2 + 324000C_w^2C_y^2N^2 - \\
& 492480C_y^2C_w^3N^2 - 492480C_w^2C_y^3N^2 + 114048C_w^3C_y^3N^2 + \\
& 259200C_y^1C_w^4N^2 + 259200C_w^1C_y^4N^2 - 777600C_w^4C_y^4N^2 - \\
& 44100C_y^1C_w^2N^3 - 44100C_w^1C_y^2N^3 - 8280C_y^1C_w^3N^3 - \\
& 8280C_w^1C_y^3N^3 + 950400C_y^2C_w^4N^3 - 250560C_y^3C_w^4N^3 + \\
& 950400C_w^2C_y^4N^3 - 250560C_w^3C_y^4N^3 - 5875C_w^1C_y^1N^4 - \\
& 133200C_w^2C_y^2N^4 + 90720C_y^2C_w^3N^4 + 90720C_w^2C_y^3N^4 - \\
& 22752C_w^3C_y^3N^4 - 52800C_y^1C_w^4N^4 - 52800C_w^1C_y^4N^4 + \\
& 396000C_w^4C_y^4N^4 + 900C_y^1C_w^2N^5 + 900C_w^1C_y^2N^5 + \\
& 4920C_y^1C_w^3N^5 + 4920C_w^1C_y^3N^5 - 100800C_y^2C_w^4N^5 + \\
& 51840C_y^3C_w^4N^5 - 100800C_w^2C_y^4N^5 + 51840C_w^3C_y^4N^5 + \\
& 75C_w^1C_y^1N^6 + 3600C_w^2C_y^2N^6 - 2880C_w^3C_y^3N^6 - \\
& 50400C_w^4C_y^4N^6)
\end{aligned}
$$

$$
\begin{aligned}
\tilde{\beta}_{yw} \;=\; & \frac{(N^2-1)(N^2-4)}{518400}(172800C_y^1C_w^2 + 172800C_w^1C_y^2 - \\
& 69120C_y^1C_w^3 - 69120C_w^1C_y^3 + 4147200C_y^2C_w^4 - \\
& 1658880C_y^3C_w^4 + 4147200C_w^2C_y^4 - 1658880C_w^3C_y^4 + \\
& 33450C_w^1C_y^1N - 837000C_w^2C_y^2N - 79920C_y^2C_w^3N - \\
& 79920C_w^2C_y^3N + 197856C_w^3C_y^3N + 457200C_y^1C_w^4N + \\
& 457200C_w^1C_y^4N + 2678400C_w^4C_y^4N - 107700C_y^1C_w^2N^2 - \\
& 107700C_w^1C_y^2N^2 - 2640C_y^1C_w^3N^2 - 2640C_w^1C_y^3N^2 - \\
& 910800C_y^2C_w^4N^2 + 96480C_y^3C_w^4N^2 - 910800C_w^2C_y^4N^2 + \\
& 96480C_w^3C_y^4N^2 - 13175C_w^1C_y^1N^3 + 18000C_w^2C_y^2N^3 + \\
& 83880C_y^2C_w^3N^3 + 83880C_w^2C_y^3N^3 - 43200C_w^3C_y^3N^3 - \\
& 100800C_y^1C_w^4N^3 - 100800C_w^1C_y^4N^3 - 597600C_w^4C_y^4N^3 +
\end{aligned}
$$

$$1500C_y^1C_w^2N^4 + 1500C_w^1C_y^2N^4 + 9480C_y^1C_w^3N^4 +$$
$$9480C_w^1C_y^3N^4 + 59760C_y^3C_w^4N^4 + 59760C_w^3C_y^4N^4 +$$
$$125C_w^1C_y^1N^5 - 5976C_w^3C_y^3N^5)$$

In all cases (4.100-4.104), Born level correlators are nonzero for general N, and so are the order g^2 contributions for a random set of coefficients C^j. But when we set the C^j to their proper values (to make $\mathcal{O}_y$ and $\mathcal{O}_w$ $\frac{1}{4}$-BPS), we recover correlators which are nonvanishing ($\alpha_{\text{free}} \neq 0$) and protected at order g^2 ($\tilde{\beta}_{xw} = \tilde{\beta}_{yw} = 0$).

CHAPTER 5

Strings in the near plane wave background and AdS/CFT

So far, we have been discussing the properties of supergravity modes, and the corresponding protected SYM operators. But we can do better than that. As it turns out, the GS superstring can be quantized exactly in the plane wave background [31, 32], which can be viewed as the Penrose limit of the $AdS_5 \times S^5$ geometry [30, 33]. The limit involves scaling both the AdS_5 radius $R \to \infty$ and the R-charge $J \sim R^2$. One considers states with finite plane wave light cone energy and momentum. It was proposed by Berenstein, Maldacena and Nastase (BMN) [30] that such string states correspond to single trace operators in the gauge theory with certain phases inserted. Remarkably, the parameter controlling perturbative expansion of scaling dimensions of such operators is $\lambda' = gN/J^2$, which can be made small to allow reliable gauge theory computations. BMN were able to resum the diagrams weighted by powers of λ' and show precise agreement between the scaling dimensions of SYM operators and the light cone energies of corresponding string states. This was further confirmed in [34, 35, 36]. The following development included studying string interactions both in the plane wave string theory and in the gauge theory [36, 37, 38, 39, 40, 41, 42, 43, 44, 45].

The plane wave limit is an improvement over being able to handle only supergravity states and protected operators. But we would still like to get closer to the full AdS string theory. One way to gain insight is to do systematic perturbation theory around the plane wave limit, taking $1/R^2$ as a small parameter. This approach was tested in [47] on the $AdS_3 \times S^3$ background with NS-NS flux. String theory in this background is described by an exactly solvable $SL(2) \times SU(2)$ WZNW model. It was shown [47] that one can recover the exact string spectrum at small coupling g to the next to leading order in $1/R^2$ expansion.

In this Chapter we determine the leading order finite radius corrections to the string spectrum in $AdS_5 \times S^5$. Specifically, we carry out systematic perturbation theory around the plane wave limit, taking $1/R^2$ as a small parameter. On the Yang Mills side, the corresponding calculation involves refining the definition of BMN operators and computing their scaling dimensions. We work at small string coupling g, which corresponds to computing only planar diagrams in the gauge theory. Furthermore, we consider only the leading non-trivial term in the λ' expansion. The calculation of scaling dimensions in SYM then reduces to computing the matrix of two-point functions and its subsequent diagonalization. We identify the gauge theory operator which corresponds to the light cone worldsheet Hamiltonian, and show that its matrix elements relevant for diagonalization agree with the string theory results. Hence we conclude that to the accuracy we are working at, the scaling dimensions of gauge theory operators agree with the spectrum of string states in $AdS_5 \times S^5$.[1]

The Chapter is organized in the following way. In section 5.1 we describe how to quantize the string in the background which includes the $\mathcal{O}(1/R^2)$ corrections to the plane wave metric, and show how to compute the leading corrections to the spectrum of bosonic plane wave states. In section 5.2 we explain how the definition of BMN operators should be extended to include finite J effects.

---

[1] Related issues were also addressed in [68, 69, 70].

There we also establish agreement between string and SYM results for a subset of matrix elements of the light cone Hamiltonian. In section 5.3 we discuss our results and mention possible future developments. In appendix 5.4.1 we present an alternative technique, based on the formalism of [67], for computing $1/R^2$ corrections in string theory. The results for physical quantities are the same as in section 5.1. Appendix 5.4.2 contains the tools we use in the SYM calculations. In appendix 5.4.3 we generalize the results of section 5.2.

5.1 Corrections to the plane wave string spectrum

In this section, we do perturbation theory on the worldsheet following the method described in [47]. We start by outlining the procedure used in [31, 32, 30] for deriving the leading order spectrum in the Penrose limit of $AdS_5 \times S^5$. The $AdS_5 \times S^5$ metric is

$$ds^2 = R^2 \left[-dt^2 \cosh^2\!\rho + d\rho^2 + \sinh^2\!\rho \, d\Omega_3^2 + d\psi^2 \cos^2\!\theta + d\theta^2 + \sin^2\!\theta \, d\Omega_3'^2 \right]. \quad (5.1)$$

The Penrose limit of this geometry is obtained by zooming in on the neighborhood of a lightlike geodesic circling the equator of S^5. This is done by changing variables as

$$X^+ = \frac{1}{2}(t+\psi), \quad X^- = \frac{1}{2}(t-\psi)R^2, \quad \rho = \frac{r}{R}, \quad \theta = \frac{y}{R}, \qquad (5.2)$$

and taking R to be large, while keeping $|X^\pm|, r, y$ finite. At leading order in $1/R^2$, the $AdS_5 \times S^5$ metric (5.1) reads

$$ds_0^2 = -4dX^- dX^+ - (r^2 + y^2) dX^+ dX^+ + dr_i \, dr_i + dy_i \, dy_i. \qquad (5.3)$$

Coordinates y_i and r_i parameterize two copies of R^4, but the $SO(8)$ symmetry of the metric (5.3) is broken down to $SO(4) \times SO(4)$ by the RR flux

$$F_{+1234} = F_{+5678} = \text{const.} \qquad (5.4)$$

We would like to quantize type IIB superstring in the background (5.3), (5.4). As was shown in [31, 32], the way to do this is to look at the sigma-model part of the GS action, and use κ-symmetry in light-cone gauge to determine the rest of the worldsheet action. Bosons and fermions decouple for the plane wave background (5.3) in light-cone gauge [31, 32]. We will only be interested in the bosonic part of the full superstring action. The light cone gauge is specified by

$$\begin{aligned} X^{+} &= \tau, && (5.5)\\ \partial_\sigma \gamma_{\sigma\sigma} &= 0, \\ \det\gamma_{\alpha\beta} &= -1, \end{aligned}$$

for bosonic fields; the worldsheet coordinates are $\tau \in (-\infty, \infty)$, $\sigma \in [0, l]$. The worldsheet metric can be written as [71]

$$\gamma^{\alpha\beta} = \begin{pmatrix} -\gamma_{\sigma\sigma}(\tau) & \gamma_{\sigma\tau}(\tau,\sigma) \\ \gamma_{\sigma\tau}(\tau,\sigma) & \gamma_{\sigma\sigma}^{-1}(\tau)(1-\gamma_{\sigma\tau}^{2}(\tau,\sigma)) \end{pmatrix}. \qquad (5.6)$$

In this section we consider only the y part of the theory. The r part can be included by noticing that (5.3) and (5.4) are invariant under $y \leftrightarrow r$ while in the $\mathcal{O}(1/R^2)$ correction to the plane wave metric y and r terms come with opposite signs (see below). This means that to restore the r terms in the final result one needs to copy the y part, substitute $y \to r$ and flip the sign in front of the $\mathcal{O}(1/R^2)$ terms. This is confirmed in appendix 5.4.1, where explicit calculations are performed.

In the light cone gauge (5.5) the bosonic part of the Lagrangian is

$$\begin{aligned} L_0 &= -\frac{1}{4\pi}\int_0^l \Big\{\gamma_{\sigma\sigma}\Big[4\dot{X}^{-}+\sum_i (y_i y_i - \dot{y}_i\dot{y}_i)\Big] \\ &\qquad -2\gamma_{\sigma\tau}\Big[2(X^{-})' - \sum_i \dot{y}_i y_i'\Big] + \gamma_{\sigma\sigma}^{-1}(1-\gamma_{\sigma\tau}^{2})\sum_i y_i' y_i'\Big\}, \qquad (5.7) \end{aligned}$$

where we used the leading order spacetime metric (5.3). The equation of motion for the worldsheet metric (Virasoro constraints) are

$$(X^-)' = \frac{1}{2}\sum_i \dot{y}_i y_i', \qquad \dot{X}^- = \frac{1}{4}\sum_i \left[\dot{y}^i \dot{y}^i + y_i' y_i' - y_i y_i\right]. \tag{5.8}$$

One can use the equation of motion for X^- and the leftover gauge freedom $\sigma \to \sigma + f(\tau)$ to set $\gamma_{\sigma\tau} = 0$ in (5.7) [71]. The equation of motion for the zero mode of X^- implies that $\gamma_{\sigma\sigma}$ is related to the conserved light cone momentum $\mathcal{P}_- = -i\frac{\partial}{\partial X^-}$. Choosing the gauge

$$l = 2\pi\eta, \quad \text{where } \eta \equiv -\tfrac{1}{2}\mathcal{P}_-, \tag{5.9}$$

sets $\gamma_{\sigma\sigma} = 1$ at the leading order in $1/R^2$. The plane wave Hamiltonian that follows from (5.7) can therefore be written as

$$H_0 = \frac{1}{4\pi}\int_0^l d\sigma \sum_i \left[(2\pi)^2 P_y^i P_y^i + y_i y_i + y_i' y_i'\right]. \tag{5.10}$$

where $P_y^i = \dot{y}_i/2\pi$. The worldsheet theory of a light cone string is massive in the plane wave background. The fields can be expressed in terms of eigenmodes

$$y_i = \frac{i}{\sqrt{2}}\sum_n \frac{1}{\sqrt{w_n}}\left[y_n^i - y_n^{i\,\dagger}\right], \tag{5.11}$$

where the τ, σ-dependent oscillators $y^i, y_n^{i\,\dagger}$ are defined as

$$y_n^i = \alpha_n^i e^{\frac{-iw_n\tau - in\sigma}{\eta}}, \qquad y_n^{i\,\dagger} = \alpha_n^{i\,\dagger} e^{\frac{iw_n\tau + in\sigma}{\eta}}, \tag{5.12}$$

and the frequencies are given by

$$w_n = \sqrt{\eta^2 + n^2}. \tag{5.13}$$

Substituting the field expansions into (5.10) diagonalizes the plane wave Hamiltonian

$$H_0 = \frac{1}{\eta}\sum_{i,n} w_n N_n^i, \tag{5.14}$$

where $N_n^i = y_n^{i\,\dagger} y_n^i$. The normal ordering constant cancels between bosons and fermions by virtue of spacetime supersymmetry, so we do not include it in (5.14). The leading terms in the expansion of H_0 in powers of $1/\eta^2$ are

$$H_0 = \sum_{i,n} N_n^i + \frac{1}{2\eta^2} \sum_{i,n} n^2 N_n^i + \mathcal{O}\left(\frac{1}{\eta^4}\right). \tag{5.15}$$

In addition, we have the level matching condition

$$\sum_{i,n} n N_n^i = 0. \tag{5.16}$$

To compute $\mathcal{O}(1/R^2)$ corrections to the string spectrum in the plane wave background, one would add the $\mathcal{O}(1/R^2)$ correction ds_1^2 to the leading metric ds_0^2, write down the bosonic part of the light cone Lagrangian, and then use κ-symmetry to write the full GS action. Subsequently the system can be quantized perturbatively in $1/R^2$. Expanding (5.1) to next to leading order in $1/R^2$ we have

$$ds_1^2 = \frac{1}{R^2}\left[-2dX^- dX^+ (r^2 - y^2) - \frac{1}{3}(r^4 - y^4) dX^+ dX^+ + \frac{1}{3}(r^4 d\Omega_3{}^2 - y^4 d\Omega_3'{}^2)\right]. \tag{5.17}$$

The bosonic part of the $\mathcal{O}(1/R^2)$ Lagrangian is therefore quartic in the fields. The leading form of the κ-symmetry then implies that the fermionic part of the $\mathcal{O}(1/R^2)$ GS action is at most bi-quadratic in bosons and fermions. We are considering corrections to the spectrum of bosonic states, so the fermionic part of the action can only contribute diagonal matrix elements of the type

$$\frac{1}{R^2} \sum_{i,n} f(w_n) N_n^i, \tag{5.18}$$

where $f(w_n)$ is some function. Fixing the exact form of $f(w_n)$ in (5.18) requires dealing with the $\mathcal{O}(1/R^2)$ fermionic part of the superstring action. This we have not bothered to do. We also drop all terms that are due to the normal ordering of bosonic operators in all subsequent calculations.

Using the identities $dy_i dy_i = dy^2 + y^2 d\Omega^2$ and $y dy = y_i dy_i$ we can write $y^4 d\Omega_3'^2 = y_i y_i dy_j dy_j - y_i y_j dy_i dy_j$ and deduce the correction to the leading order Lagrangian (5.7)

$$L_1 = \frac{1}{4\pi R^2}\int_0^l d\sigma \left[\frac{1}{3}\sum_i y_i^4 - \frac{1}{3}\sum_{i\neq j}\Big[y_i^2(\dot{y}_j^2 - y_j^2 - (y_j')^2) + y_i y_j (y_i' y_j' - \dot{y}_i \dot{y}_j)\Big] + \frac{1}{2} y^2 \dot{X}^-\right]. \tag{5.19}$$

Terms proportional to $\gamma_{\sigma\tau}$ are higher order in $1/R^2$ and do not contribute to (5.19). As explained in [47], for the purpose of computing the leading corrections to the spectrum, the correction to the Hamiltonian equals minus the correction to the Lagrangian.[2] The correction to the plane wave Hamiltonian can therefore be written as

$$\begin{aligned} H_1 \;=\; & \frac{1}{4\pi R^2}\int_0^l d\sigma \Bigg[-\frac{1}{3}\sum_i y_i^4 + \frac{1}{3}\sum_{i\neq j}\Big[y_i^2[(2\pi P_y^j)^2 - y_j^2 - (y_j')^2] \\ & + y_i y_j (y_i' y_j' - (2\pi)^2 P_y^i P_y^j)\Big] - \frac{1}{2}\sum_{i,j} y_i^2[(2\pi P_y^j)^2 + (y_j')^2 - y_j^2]\Bigg], \end{aligned} \tag{5.20}$$

where in rewriting the last term we used the Virasoro constraint [the second equation in (5.8)].

Next we expand (5.20) in modes (5.11). We are interested in first order corrections to the energies, so we only need to compute matrix elements of H_1 between degenerate states. Plane wave string states are

$$y_{n_1}^{i_1\,\dagger} \ldots y_{n_k}^{i_k\,\dagger} \ldots |\eta\rangle. \tag{5.21}$$

They are degenerate only if the two sets of worldsheet momenta $(n_1, \ldots n_k, \ldots)$ and $(n_1', \ldots n_k', \ldots)$ are permutations of one another. Thus the only relevant terms in H_1 are of the form $y_k y_k^\dagger y_l y_l^\dagger$. Diagonal contributions come from $y_k^i y_k^{i\,\dagger} y_l^j y_l^{j\,\dagger}$; they add up to

---

[2] One can convince oneself that this is the case by perturbing the Lagrangian, calculating the canonically conjugate momenta, and keeping only terms up to $\mathcal{O}(1/R^2)$ in the Hamiltonian. In [47] the zero mode of X^- was treated separately, but one can show that this is not necessary.

$$H_1^D = \frac{1}{2\eta R^2}\left(\frac{1}{2}\sum_{i;n}\frac{n^2(N_n^i)^2}{w_n^2} - \sum_{i,j;m,n}\frac{n^2 N_n^i N_m^j}{w_m w_n}\right). \tag{5.22}$$

The relevant off-diagonal terms are of the form ${y_m^i}^\dagger y_n^i {y_n^j}^\dagger y_m^j$, $i \neq j$, $m \neq n$; and ${y_m^i}^\dagger {y_n^i}^\dagger y_m^j y_n^j$, $i \neq j$. These add up to

$$H_1^{OD} = \frac{1}{2\eta R^2}\sum_{i\neq j;\, m\neq n}\frac{nm}{w_n w_m}\left({y_m^i}^\dagger {y_n^i}^\dagger y_m^j y_n^j - {y_m^i}^\dagger y_n^i {y_n^j}^\dagger y_m^j\right) + \frac{1}{4\eta R^2}\sum_{i\neq j;n}\frac{n^2}{w_n^2}{y_n^i}^\dagger {y_n^i}^\dagger y_n^j y_n^j. \tag{5.23}$$

Expanding (5.22) and (5.23) in powers of $1/\eta$ we obtain

$$H_1^D = \frac{1}{2\eta^3 R^2}\left(\frac{1}{2}\sum_{i;n} n^2 (N_n^i)^2 - \sum_{i,j;m,n} n^2 N_n^i N_m^j\right) + \mathcal{O}\left(\frac{1}{\eta^5 R^2}\right) \tag{5.24}$$

and

$$\begin{aligned} H_1^{OD} &= \frac{1}{2\eta^3 R^2}\sum_{i\neq j;\, m\neq n} nm\left({y_m^i}^\dagger {y_n^i}^\dagger y_m^j y_n^j - {y_m^i}^\dagger y_n^i {y_n^j}^\dagger y_m^j\right) \\ &\quad + \frac{1}{4\eta^3 R^2}\sum_{i\neq j;n} n^2 {y_n^i}^\dagger {y_n^i}^\dagger y_n^j y_n^j + \mathcal{O}\left(\frac{1}{\eta^5 R^2}\right), \end{aligned} \tag{5.25}$$

respectively. The leading $1/\eta$ term in H_1 is a sum of these two expressions.

An alternative derivation is given in appendix 5.4.1, where more details are provided.

5.2 Anomalous dimensions and AdS/CFT

We now turn to the boundary $\mathcal{N} = 4$ Super Yang-Mills theory. Our starting point will be the BMN operators [30] which correspond to plane wave states in the Penrose limit. One can still regard plane wave states as belonging to the Hilbert space of the full $AdS_5 \times S^5$ theory, even though they are no longer eigenstates of the full Hamiltonian. As explained in the previous section, departing from the Penrose limit corresponds to turning on perturbative corrections to the plane wave Hamiltonian. Eigenstates of the full Hamiltonian can be found using ordinary quantum-mechanical perturbation theory.

SYM operators which correspond to string eigenstates must have definite conformal dimensions. Such operators may be obtained from a complete set of operators by diagonalizing the matrix of their two-point functions. This procedure is analogous to the diagonalization of the string theory Hamiltonian. We find that the spectra computed on both sides of the correspondence match, and the operator defined by the matrix of two-point functions is the SYM counterpart of the string Hamiltonian.

This section is organized as follows. In section 5.2.1 we define operators that correspond to plane wave states away from the strict Penrose limit. In section 5.2.2 we show how the matrix of two-point functions is related to the string Hamiltonian. In section 5.2.3 we match the matrix elements of the light cone Hamiltonian between the string and the gauge theory. We analyze a simple case where all of the excited modes have distinct $SO(4)$ indices and none of them is excited more than once. The most general case is treated in appendix 5.4.3. Feynman rules are discussed in appendix 5.4.2.

5.2.1 Operators

The important assumption that we start with is that suitably refined BMN operators continue to correspond to plane wave states, regarded as states in the Hilbert space of $AdS_5 \times S^5$, even away from the plane wave limit. To define the right operators we will follow closely the logic of BMN. We start with the operator which corresponds to the light cone vacuum

$$\frac{1}{\sqrt{\Omega}}\mathrm{tr}\,[z^J] \quad \leftrightarrow \quad |\eta\rangle, \tag{5.26}$$

where $z = \frac{1}{\sqrt{2}}(\phi^5 + i\phi^6)$ and Ω is a normalization constant (more about this below). For the ground state (5.26) there is a relation $J = R^2\eta$, but this gets modified by $\mathcal{O}(1/R^2)$ terms for excited states.

SYM operators which correspond to states with excited zero modes can be generated by acting on the light cone ground state (5.26) with generators of the global symmetry group. The generators that we will be interested in are rotations in ij plane, denoted by T_{ij} and their combinations $T_{iz} = \frac{1}{\sqrt{2}}(T_{i5} + iT_{i6})$ and $T_{i\bar{z}} = \frac{1}{\sqrt{2}}(T_{i5} - iT_{i6})$. They act on the fields as

$$\begin{aligned} [T_{iz}, z] &= 0, \qquad [T_{iz}, \bar{z}] = \phi^i, \qquad [T_{iz}, \phi^j] = -z\,\delta_i^j, \\ [T_{i\bar{z}}, z] &= \phi^i, \qquad [T_{i\bar{z}}, \bar{z}] = 0, \qquad [T_{i\bar{z}}, \phi^j] = -\bar{z}\,\delta_i^j. \end{aligned} \tag{5.27}$$

On the worldsheet we have a correspondence

$$T_{iz} \leftrightarrow y_0^i, \quad T_{i\bar{z}} \leftrightarrow y_0^{i\,\dagger}. \tag{5.28}$$

Consider as an example the operator corresponding to the state $y_0^{i\,\dagger} y_0^{j\,\dagger}|\eta\rangle$, $i \neq j$. It is obtained by computing successive commutators of $T_{i\bar{z}}$ and $T_{j\bar{z}}$ with (5.26). Either of these generators can turn any z in the string of z-s into ϕ^i or ϕ^j respectively. The result is therefore the sum of $\mathrm{tr}\,[z, \ldots \phi^i\, z\, \ldots \phi^j\, z \ldots]$ over all possible positions of inserted ϕ's:

$$\frac{1}{\sqrt{\Omega}} \left[\sum_{a=0}^{J} \sum_{b=a}^{J} \mathrm{tr}\,[z^a\, \phi^i\, z^{b-a}\, \phi^j\, z^{J-b}] + (i \leftrightarrow j) \right] \quad \leftrightarrow \quad y_0^{i\,\dagger} y_0^{j\,\dagger}|\eta\rangle. \tag{5.29}$$

This formula has an obvious generalization for higher number of ϕ insertions, as long as no label appears more than once. If some of the ϕ's indices do coincide, $T_{i\bar{z}}$ can act on the same field. In this case, z is first turned into ϕ^i, and then into $-\bar{z}$. For example, in the case of two ϕ insertions we have

$$\frac{1}{\sqrt{\Omega}} \left(2 \sum_{a=0}^{J} \sum_{b=a}^{J} \mathrm{tr}\,[z^a\, \phi^i\, z^{b-a}\, \phi^i\, z^{J-b}] - \sum_{a=0}^{J+1} \mathrm{tr}\,[z^a\, \bar{z}\, z^{J+1-a}] \right) \quad \leftrightarrow \quad y_0^{i\,\dagger} y_0^{i\,\dagger}|\eta\rangle. \tag{5.30}$$

To construct an operator with three ϕ's with the same index inserted, one should act by $T_{i\bar{z}}$ on both terms in (5.30) to produce

$$\frac{1}{\sqrt{\Omega}}\Big(\sum \mathrm{tr}\,[z\ldots\phi^i\ldots\phi^i\ldots\phi^i\ldots] - 3\sum \mathrm{tr}\,[z\ldots\phi^i\ldots\bar{z}\ldots]\Big) \quad \leftrightarrow \quad y_0^{i\,\dagger}y_0^{i\,\dagger}y_0^{i\,\dagger}|\eta\rangle, \tag{5.31}$$

where dots stand for a bunch of z's and the sum is over all possible positions of the insertions. The second sum in (5.31) has $J+1$ times fewer terms than the first sum, and is subleading when it comes to computing two-point functions. Throughout this discussion we are interested in the subleading corrections in $1/J \sim 1/\eta R^2$, and therefore we should keep this term. If we act with $T_{i\bar{z}}$ one more time, a term $3\sum \mathrm{tr}\,[z\ldots\bar{z}\ldots\bar{z}\ldots]$ appears when $T_{i\bar{z}}$ hits the ϕ^i in the second sum in (5.31). This piece is $\mathcal{O}(1/J^2)$ compared to the leading term, so we can drop it.

In general, when an arbitrary number of zero modes excited, the corresponding SYM operator is

$$\mathcal{O} \;=\; \tilde{\mathcal{O}} - \mathcal{O}_*, \tag{5.32}$$

$$\tilde{\mathcal{O}} \;=\; \frac{1}{\sqrt{\Omega}}\sum \mathrm{tr}\,[z\ldots\phi^{i_1}\ldots\phi^{i_k}\ldots], \tag{5.33}$$

$$\mathcal{O}_* \;=\; \frac{1}{\sqrt{\Omega}}\sum_{(p,q):\,i_p=i_q} \mathrm{tr}\,[z\ldots\phi^{i_1}\ldots\phi^{i_k}\ldots\check{\phi}^{i_p}\ldots\check{\phi}^{i_q}\ldots\bar{z}\ldots], \tag{5.34}$$

where $\check{\phi}^{i_p}$ stands for ϕ^{i_p} being omitted from the string of operators and the sum in $\mathcal{O}_*$ runs over all possible pairs of (ϕ^{i_p},ϕ^{i_q}) with the same indices. When writing (5.32), we omitted terms which appear when $T_{i\bar{z}}$ hits the same field more than twice, as such are $\mathcal{O}(1/J^2)$. When all ϕ's inserted have different flavors, the operator $\mathcal{O}_*$ vanishes and we have $\mathcal{O} = \tilde{\mathcal{O}}$.

Next we turn to the construction of operators which correspond to general string states (5.21). Such operators must satisfy a few necessary requirements. First, if only the zero modes are excited, they must reduce to the BPS operators described. Second, they must vanish unless the level matching condition

$$\sum_{i,n} n N^i_n = 0, \tag{5.35}$$

is satisfied. Finally, our operators must reduce to the BMN operators as $J \to \infty$.

Suppose there is a total of N oscillators excited,

$$\mathsf{N} = \sum_i \mathsf{N}_i, \qquad \mathsf{N}_i = \sum_n N^i_n. \tag{5.36}$$

Due to the cyclicity of the trace,

$$\frac{1}{\sqrt{\Omega}} \mathrm{tr}\,[z \ldots \phi^{i_1} \ldots \phi^{i_k} \ldots] \tag{5.37}$$

is equivalent to $(J + \mathsf{N})$ other terms in $\tilde{\mathcal{O}}$ which are related to it by cyclic permutations. According to [30], at the leading order in the $1/J$ expansion, oscillators ${y^i_n}^\dagger$ correspond to insertions of ϕ^{i_k} with the phase $\exp(\frac{2\pi i n_k a_k}{J})$, where a_k counts the number of z's to the left of this ϕ^{i_k}. One has to be be more careful when $1/J$ effects are taken into account. In order for an operator to vanish when the level matching condition is not satisfied, each sum over cyclically related terms in (5.37) must vanish separately. This happens when the phases assigned to the ϕ^{i_k} insertions are

$$q^{a_k}_{n_k} = \exp\left(\frac{2\pi i n_k a_k}{J + \mathsf{N}}\right). \tag{5.38}$$

Here a_k counts *a*ll operators appearing to the left of the ϕ^{i_k} insertion, and not just the z-s. Similar arguments can be made to fix the form of $\mathcal{O}_*$. Again, each ϕ^{i_k} insertion comes with a phase given by (5.38). In order to satisfy the level matching condition we should also assign a phase $q^{a_{\bar{z}}}_{n_k+n_l}$ to $\bar{z}$.

To summarize, we have a correspondence which relates SYM operators and plane wave string states away from the strict Penrose limit

$$\mathcal{O} = (\tilde{\mathcal{O}} - O_*) \quad \leftrightarrow \quad {y^{i_1}_{n_1}}^\dagger \ldots {y^{i_k}_{n_k}}^\dagger \ldots |\eta\rangle, \tag{5.39}$$

where

$$\tilde{\mathcal{O}} \;=\; \frac{1}{\sqrt{\Omega}} \sum \left(\prod_k q_{n_k}^{a_k} \right) \mathrm{tr}\,[z \ldots \phi^{i_1} \ldots \phi^{i_k} \ldots], \tag{5.40}$$

$$\mathcal{O}_* \;=\; \frac{1}{\sqrt{\Omega}} \sum_{(n_p, n_q):\, i_p = i_q} \left(\prod_{k \neq p,q} q_{n_k}^{a_k} \right) q_{n_p+n_q}^{a_{\bar{z}}} \,\mathrm{tr}\,[z \ldots \phi^{i_1} \ldots \phi^{i_k} \ldots \check{\phi}^{i_k} \ldots \check{\phi}^{i_l} \ldots \bar{z} \ldots], \tag{5.41}$$

and the phases $q_{n_k}^{a_k}$ are given by (5.38).

The normalization constant Ω will be chosen so that the leading term in $1/J$ expansion of the $\mathcal{O}(g^0)$ two-point function is normalized to one. This leading term is given by the interaction-free diagrams



, (5.42)

where the subscript n_k in $\phi_{n_k}^{i_k}$ means that the corresponding insertion of ϕ^{i_k} in the string of operators comes with the phase $q_{n_k}^{a_k}$. Expression (5.42) contains only contractions of the same $\phi_{n_k}^{i_k}$. Interaction-free diagrams with contractions of $\phi_{n_k}^{i_k}$ and $\phi_{n_l}^{i_l}$ with $n_k \neq n_l$ are also allowed, as long as $i_k = i_l$. Such diagrams however are subleading in $1/J$.

From (5.42) we infer that

$$\Omega = c N^{J+\mathsf{N}} (J + \mathsf{N}) \tilde{\Omega}, \tag{5.43}$$

where c is an irrelevant numerical prefactor; $N^{J+\mathsf{N}}$ arises from the number of color loops in (5.42); and $(J+\mathsf{N})$ takes care of the fact that performing a cyclic permutation in one of the operators entering the two-point function gives an equivalent diagram. When no oscillators are excited more than once, there is no further choice of contractions and $\tilde{\Omega}$ is equal to the number of ways N ϕ's can be distributed among J z's, $\tilde{\Omega} = \prod_{n=1}^{\mathsf{N}} (J+n)$. When there are multiple excitations of the same mode, there can be $N^i_n!$ inequivalent permutations of the ϕ^i_n in either one of the operators. This gives rise to $N^i_n!$ copies of the diagram (5.42). We conclude that in general,

$$\tilde{\Omega} = \prod_k N^{i_k}_{n_k}! \prod_{n=1}^{\mathsf{N}} (J+n). \tag{5.44}$$

5.2.2 Two-point functions and the light cone Hamiltonian

The light cone energy of a string state and its momentum are related to the anomalous dimension Δ and R-charge J of the corresponding operator as follows

$$\begin{aligned} H_{lc} &= -\mathcal{P}_+ &= i\frac{\partial}{\partial X^+} = \Delta - J, \quad (5.45)\\ \eta &= -\tfrac{1}{2}\mathcal{P}_- = \frac{i}{2}\frac{\partial}{\partial X^-} &= \frac{\Delta + J}{2R^2}. \quad (5.46)\end{aligned}$$

One can find anomalous dimensions of the gauge theory operators by looking at two-point functions, and we are now going to explain this in detail. We will only consider planar diagrams. This amounts to neglecting string amplitudes of genus one and higher. Furthermore, we will only look at the terms in H_{lc} which behave like

$$\begin{aligned} \frac{1}{\eta^{2a}} &\sim \left(\frac{R^2}{J}\right)^{2a} &= \frac{(4\pi g N)^a}{J^{2a}}, \\ \frac{1}{R^2\eta^{2a+1}} &\sim \frac{1}{R^2}\left(\frac{R^2}{J}\right)^{2a+1} &= \frac{(4\pi g N)^a}{J^{2a+1}}, \end{aligned} \tag{5.47}$$

with $a = 0, 1$.

On the string theory side, the first line in (5.47) corresponds to the truncated expansion in powers of $1/\eta^2$ of the plane wave Hamiltonian H_0. The second line corresponds to the expansion of H_1. Terms in the two lines differ by a factor of $1/J$. On the gauge theory side this factor arises when finite J corrections are taken into account, which leads to the modification of BMN operators, explained in section 5.2.1. The first perturbative (from the SYM point of view) correction to the light cone energy in (5.47) corresponds to $a = 1$, which implies that $a = 0$ term in the second line of (5.47) vanishes. This is in complete accord with the expansion of H_1 in powers of $1/\eta$.

Consider a set of gauge theory operators $\mathcal{O}_\alpha$ labeled by $\alpha = \{(i_k, n_k)\}$. We will be interested in the SYM operators which correspond to plane wave states with N worldsheet oscillators excited. Their two-point functions can be arranged as

$$\begin{aligned} \langle \mathcal{O}_\alpha(x)\bar{\mathcal{O}}_\beta(0)\rangle &= \langle \mathcal{O}_\alpha(x)\bar{\mathcal{O}}_\beta(0)\rangle_{g^0} + \langle \mathcal{O}_\alpha(x)\bar{\mathcal{O}}_\beta(0)\rangle_{g^1} + \mathcal{O}(g^2) \\ &= \frac{1}{|x|^{2(J+\mathsf{N})}}\left[\mathbf{T}_{\alpha\beta} - \mathbf{F}_{\alpha\beta}\log(\mu^2 x^2) + \mathcal{O}(g^2)\right]. \end{aligned} \tag{5.48}$$

Here, $\mathbf{T}$ is a matrix of combinatorial factors which come from interaction-free diagrams in $\langle \mathcal{O}_\alpha(x)\bar{\mathcal{O}}_\beta(0)\rangle_{g^0}$, while $\mathbf{F}$ captures the $\mathcal{O}(g)$ effects of SYM interactions in $\langle \mathcal{O}_\alpha(x)\bar{\mathcal{O}}_\beta(0)\rangle_{g^1}$. $\mathcal{O}(g)$ contributions to the two point functions (5.48) come from diagrams of the type



. (5.49)

In appendix 5.4.2 we show that (5.49) is equal to

$$\gamma \equiv -\beta \log \mu^2 x^2 \equiv -\frac{gN}{2\pi}\log \mu^2 x^2 \tag{5.50}$$

times a numerical factor determined by the fields which go into the 4-point vertex.

Operators $\mathcal{O}_\alpha$ may not have well defined scaling dimensions at order $\mathcal{O}(g)$; the same phenomenon occured in Chapters 2 and 3. To find pure operators and their anomalous dimensions, we need to transform to a basis of eigenstates of the dilatation operator. By a linear transformation, we should bring (5.48) to the form

$$\frac{1}{|x|^{2(J+\mathsf{N})}}\left[\mathbf{1} - \mathrm{diag}[\{\lambda_\rho\}]\log(\mu^2 x^2)\right], \tag{5.51}$$

where the order $\mathcal{O}(g)$ anomalous dimensions λ_ρ are the eigenvalues of $\mathbf{T}^{-1}\mathbf{F}$, and $\mathbf{1}$ is a unit matrix, see Section 2.6. The matrices in (5.48) have the form

$$\begin{aligned}\mathbf{T} &= \mathbf{1} + \frac{1}{J}\mathbf{T}^{(1)} + \mathcal{O}(1/J^2), && (5.52)\\ \mathbf{F} &= \mathbf{F}^{(0)} + \frac{1}{J}\mathbf{F}^{(1)} + \mathcal{O}(1/J^2), && (5.53)\end{aligned}$$

since the operators $\mathcal{O}_\alpha$ were chosen to be orthonormal at leading order, see the end of section 5.2.1. Hence, up to corrections that are higher order in $1/J$

$$\mathbf{T}^{-1}\mathbf{F} = \mathbf{F}^{(0)} + \frac{1}{J}\left(\mathbf{F}^{(1)} - \mathbf{T}^{(1)}\mathbf{F}^{(0)}\right) + \mathcal{O}(1/J^2). \tag{5.54}$$

Finally, light cone energies of worldsheet states are related to the quantum numbers of operators in $\mathcal{N}$=4 SYM as

$$\Delta - J = \mathsf{N} + \lambda_\rho. \tag{5.55}$$

In other words, $\mathsf{N}\,\mathbf{1} + \mathbf{T}^{-1}\mathbf{F}$, plays the role of the light cone Hamiltonian. In the next section we will show that $H_{lc} - \mathsf{N} = H_0 + H_1 - \mathsf{N}$ is identical to $\mathbf{T}^{-1}\mathbf{F}$ computed in the gauge theory [the H_0 and H_1 are given by (5.15), (5.24) and (5.25)]. This means that to the accuracy we are working at, the spectrum of eigenstates of the light cone worldsheet Hamiltonian is the same as the spectrum of the dilatation operator in SYM.

5.2.3 Equality of matrix elements

Let us now show that $\mathbf{T}^{-1}\mathbf{F}$ and $H_{lc} - \mathsf{N}$ indeed have the same matrix elements that are relevant for the diagonalization. In this section we consider matrix elements between states with all modes having distinct SO(4) indices. We also assume that no modes are excited more than once, $N_n^i \leq 1$. In appendix C we generalize these results to matrix elements between arbitrary plane wave states.

The relevant off-diagonal terms in H_{lc} are given by (5.25). When sandwiched between

$$|\mathcal{X}\rangle = y^{i_1\,\dagger}_{n_1} \ldots y^{i\;\dagger}_{m} y^{j\,\dagger}_{n}|\eta\rangle, \qquad m \neq n, \tag{5.56}$$

and

$$|\mathcal{X}'\rangle = y^{i_1\,\dagger}_{n_1} \ldots y^{j\;\dagger}_{m} y^{i\,\dagger}_{n}|\eta\rangle, \qquad m \neq n, \tag{5.57}$$

with $i \neq j$, the off-diagonal part of the Hamiltonian (5.25) gives rise to the following matrix element

$$\langle\mathcal{X}|H_1^{OD}|\mathcal{X}'\rangle = -\frac{1}{J}\left(\frac{R^2}{J}\right)^2 mn\sqrt{N^i_m N^j_n N^i_n{}' N^j_m{}'}, \tag{5.58}$$

where we expressed η as

$$\eta = \frac{J}{R^2}\left(1 + \frac{1}{2J}\sum_{i,m} N^i_m + \mathcal{O}(1/J^2)\right). \tag{5.59}$$

This follows from (5.45) and (5.46). The second term in the brackets gives an $\mathcal{O}(1/J)$ correction when used in the leading order Hamiltonian (5.15). We should also reinstate the normal ordering term (5.18). The SYM calculations will fix it to be $\frac{1}{R^2\eta^3}\sum_{i,n} n^2 N^i_n$. Combining these contributions, the diagonal matrix elements[3] read

$$\begin{aligned} \langle\mathcal{O}|H_0{+}H_1^D{-}\mathsf{N}|\mathcal{O}\rangle \;\; &= \;\; \frac{1}{2}\left(\frac{R^2}{J}\right)^2 \sum_{i,n} n^2 N^i_n \\ &\quad + \frac{1}{J}\left(\frac{R^2}{J}\right)^2 \Big[-\sum_{i,j,m,n} n^2 N^i_m N^j_n + \sum_{i,n}\frac{n^2 N^i_n(N^i_n{+}1)}{4}\Big]. \end{aligned} \tag{5.60}$$

For the states considered in this subsection $N^i_n = 1$ and off-diagonal elements of H_1 other than (5.58) vanish.

We will also denote the SYM operators corresponding to states (5.56)-(5.57) by $\mathcal{X}$ and $\mathcal{X}'$. As explained in section 5.2.1, no $\mathrm{tr}\,[z \ldots \bar{z} \ldots \phi^i \ldots]$ terms appear as long as all i_k labels distinct. That is, $\mathcal{X}_* = \mathcal{X}'_* = 0$, and $\mathcal{X} = \tilde{\mathcal{X}}, \mathcal{X}' = \tilde{\mathcal{X}}'$. Contributions to $\mathbf{T}_{\mathcal{X}\mathcal{X}'}$ and $\mathbf{T}_{\mathcal{O}\mathcal{O}}$ come from the diagrams similar to (5.42),

---

[3] Here and below $\mathcal{O}$ stands for an arbitrary worldsheet state or SYM operator, for example $\mathcal{X}$ or $\mathcal{X}'$. Diagonal matrix elements are all given by the same expression.



. (5.61)

The top and bottom rows in (5.61) correspond to the two SYM operators entering the two-point function. Summing the phases over positions of ϕ's we obtain

$$\frac{1}{\tilde{\Omega}}\sum{}'\prod_k r_k^{a_k} = \delta_{mm'}\delta_{nn'} - \frac{\delta_{m+n,m'+n'}(1-\delta_{mm'}\delta_{nn'})}{J} + \mathcal{O}(1/J^2). \tag{5.62}$$

The prime on the sum in (5.62) means that we count modulo cyclic permutations, and we defined

$$r_{n_k} \equiv q_{n_k} q_{n'_k}{}^* = \exp\left(\frac{2\pi i(n_k - n'_k)}{J+\mathsf{N}}\right). \tag{5.63}$$

Only r_m and r_n are different from one, so (5.62) can be computed by making use of the invariance under cyclic permutations and fixing $a_m = 0$ (and so $r_m^{a_m} = 1$). The $\mathcal{O}(1/J)$ term in (5.62) appears because the range of a_n is $[1, J+\mathsf{N}-1]$. Contributions with more than two $r_{n_k} \neq 1$ are suppressed by at least $1/J^2$ compared to (5.62), so we do not need to worry about them. Comparing (5.62) with (5.48) and (5.52), we arrive at

$$\mathbf{T}^{(1)}_{\mathcal{X}\mathcal{X}} = \mathbf{T}^{(1)}_{\mathcal{X}'\mathcal{X}'} = 0, \qquad \mathbf{T}^{(1)}_{\mathcal{X}\mathcal{X}'} = -1. \tag{5.64}$$

We now turn to the computation of $\mathbf{F}$. Consider the diagrams that contribute both to $\langle \mathcal{X}(x)\bar{\mathcal{X}}'(0)\rangle_{g^1}$ and to the diagonal correlator $\langle \mathcal{O}(x)\bar{\mathcal{O}}(0)\rangle_{g^1}$. These are



$+$



$+ (\phi^{i_k}_{n_k} \leftrightarrow z, \phi^{i_k}_{n_k} \leftrightarrow \bar{z}),$ (5.65)

and



$+$



$+ (\phi^{i_k}_{n_k} \leftrightarrow \phi^{i_l}_{n_l}, \phi^{i_k}_{n'_k} \leftrightarrow \phi^{i_l}_{n'_l}).$ (5.66)

The level matching condition gives

$$m + n = m' + n'. \tag{5.67}$$

The diagrams in (5.66) which contribute to $\mathbf{F}^{(1)}_{\mathcal{X}\mathcal{X}'}$ have $n_k = n'_l = m, n_l = n'_k = n$. The contribution (5.65) differs from the interaction-free diagram (5.61) just by an overall factor

$$-\gamma\,({q_{n'_k}}^* + q_{n'_k} - 2). \tag{5.68}$$

Therefore, summing over possible configurations of fields gives (5.62) times (5.68), for a particular $\phi^{i_k}_{n_k}$ participating in the interaction vertex in (5.65). Since any one of the $\phi^{i_k}_{n_k}$ can be used in the interaction (5.65), this must be further summed over k. We find

$$-\gamma \left[\delta_{mm'}\delta_{nn'} - \frac{\delta_{m+n,m'+n'}(1 - \delta_{mm'}\delta_{nn'})}{J}\right] \sum_k ({q_{n'_k}}^* + q_{n'_k} - 2). \tag{5.69}$$

This expression overcounts certain diagrams which do not appear in (5.65). More precisely, whenever two ϕ's are sitting next to each other the $\phi - \phi$ line cannot cross or touch a $z - \bar{z}$ line, either to the left or to the right. We will deal with such diagrams separately.

We can read off the $\mathcal{O}(J^0)$ part of $\mathbf{F}$ from (5.69) by using (5.48) and $\gamma = -\beta \log \mu^2 x^2$,

$$\mathbf{F}^{(0)} = -\beta \sum_k ({q_{n_k}}^* + q_{n_k} - 2). \tag{5.70}$$

Expanding the q's in powers of $1/J$ and taking the leading term gives the result of BMN,

$$\begin{aligned}
\mathbf{F}^{(0)} &= f^{(0)}\mathbf{1}, &(5.71)\\
f^{(0)} &= \frac{2\pi g N}{J^2}\sum_k (n_k)^2 = \frac{1}{2}\left(\frac{R^2}{J}\right)^2 \sum_n n^2 N^i_n. &(5.72)
\end{aligned}$$

To get $\mathcal{O}(1/J)$ corrections to this result we have to be more careful. As explained above, in (5.69) we overcounted the configuration of fields where two ϕ's appear next to one another in the top row of (5.65), as in (5.66):

$$(q_{n_k}^{a_k} q_{n_l}^{a_l} \ldots)\mathrm{tr}\,[\ldots \phi^{i_k}\phi^{j_l}\ldots] + (k \leftrightarrow l). \tag{5.73}$$

Now diagrams in (5.65) where ϕ^{i_k} (ϕ^{j_l}) interacts with $z - \bar{z}$ propagator to the right (left) are not allowed. The value of such diagrams is

$$\begin{aligned} -\gamma(q_{n'_k}^* + q_{n'_l} - 2) q_{n_l} q_{n'_l}^* \, r_{n_k+n_l-n'_k-n'_l}^{a_k} + (k \leftrightarrow l) &= \\ -\gamma(q_{n_k}^* + q_{n_l} - 2 q_{n_l} q_{n'_l}^*) \, r_{n_k+n_l-n'_k-n'_l}^{a_k} + (k \leftrightarrow l)&, \end{aligned} \tag{5.74}$$

where we used (5.67). Their contributions have to be substituted by the ones that appear in (5.66) instead. These are given by

$$\gamma(q_{n_l} q_{n'_l}^* - q_{n_l} q_{n'_k}^*) r_{n_k+n_l-n'_k-n'_l}^{a_k} + (k \leftrightarrow l). \tag{5.75}$$

The difference of (5.75) and (5.74) is the same for both diagonal ($n'_k = n_k, n'_l = n_l$) and off-diagonal ($n'_k = n_l, n'_l = n_k$) cases, and equals

$$\gamma(q_{n_k}^* + q_{n_l} - q_{n_l} q_{n_k}^* - 1) r_{n_k+n_l-n'_k-n'_l}^{a_k} + (k \leftrightarrow l). \tag{5.76}$$

This should be summed over a_k and divided by the normalization constant $\tilde{\Omega}$. Since the number of configurations with two ϕ's next to each other is down by $1/J$ compared to the total number of configurations, we pick up an overall factor of $1/J$. Configuration which have three and more ϕ's next to each other are suppressed by even higher powers of $1/J$, and we can neglect them to the order we are working.

The full result for $\langle \mathcal{X}(x) \bar{\mathcal{X}}'(0)\rangle_{g^1}$ is given by (5.69), plus (5.76) with $n_k = m, n_l = n$. Other terms in (5.66) are $\mathcal{O}(1/J^2)$ and are not important for us. Since

$m' = n, n' = m$ for an off-diagonal element, the first term in (5.69) vanishes, and we have

$$\mathbf{F}^{(1)}_{\mathcal{X}\mathcal{X}'} = -\beta\left[\sum_k \left({q_{n'_k}}^* + q_{n'_k} - 2\right) + \left(q^*_{m-n} + q_{m-n} - q^*_m - q_m - q^*_n - q_n + 2\right)\right]. \qquad (5.77)$$

To get the corresponding off-diagonal element of the light cone Hamiltonian, we should add $-[\mathbf{T}^{(1)}\mathbf{F}^{(0)}]_{\mathcal{X}\mathcal{X}'}/J$ to $\mathbf{F}^{(1)}_{\mathcal{X}\mathcal{X}'}/J$, see (5.54). According to (5.64) and (5.70), such addition precisely cancels the first term in (5.77), and we find

$$[\mathbf{T}^{-1}\mathbf{F}]_{\mathcal{X}\mathcal{X}'} = -\frac{\beta}{J}\left(q^*_{m-n} + q_{m-n} - q^*_m - q_m - q^*_n - q_n + 2\right). \qquad (5.78)$$

Expanding the q's in powers of $1/J$, taking the leading term and substituting the value of β we arrive at

$$[\mathbf{T}^{-1}\mathbf{F}]_{\mathcal{X}\mathcal{X}'} = -\frac{1}{J}\left(\frac{R^2}{J}\right)^2 mn. \qquad (5.79)$$

This reproduces the string theory off-diagonal matrix element (5.58), since for the states we are considering $N^i_n = 1$.

Let us now compute the diagonal terms. Now all of the diagrams in (5.66) contribute, (5.76) should be summed over k and added to (5.69) with $m' = m, n' = n$. This gives

$$\langle \mathcal{O}(x)\mathcal{O}(0)\rangle_{g^1} = -\gamma \sum_k \left({q_{n_k}}^* + q_{n_k} - 2\right) + \frac{\gamma}{2J}\sum_{k\neq l}\left(q_{n_k} + q_{n_l} - q_{n_l} q^*_{n_k} - 1 + c.c.\right). \qquad (5.80)$$

Since $\mathbf{T}_{\mathcal{O}\mathcal{O}} = 1$, we have

$$[\mathbf{T}^{-1}\mathbf{F}]_{\mathcal{O}\mathcal{O}} = -\beta \sum_k \left({q_{n_k}}^* + q_{n_k} - 2\right) + \frac{\beta}{2J}\sum_{k\neq l}\left(q_{n_k} + q_{n_l} - q_{n_l - n_k} - 1 + c.c.\right). \qquad (5.81)$$

The first term gives (5.71) at the leading order, however the definition (5.38) of q_{n_k} implies that there is a $1/J$ correction to the leading term. Expanding in powers of $1/J$ and keeping terms up to $\mathcal{O}(1/J)$ one can write (5.81) as

$$[\mathbf{T}^{-1}\mathbf{F}]_{\mathcal{O}\mathcal{O}} = \frac{1}{2}\left(\frac{R^2}{J}\right)^2 \sum_n n^2 + \frac{1}{J}\left(\frac{R^2}{J}\right)^2\left(-\sum_{i,j,m,n} n^2 N^i_m N^j_n - \frac{1}{2}\sum_{k\neq l:\, i_k \neq i_l} n_k n_l\right). \qquad (5.82)$$

Using the level matching condition (which now reads $\sum_k n_k = 0$), we can write the last term in parenthesis as

$$-\frac{1}{2}\sum_{k\neq l:\, i_k\neq i_l} n_k n_l = \frac{1}{2}\sum_k n_k^2. \tag{5.83}$$

Substituting this back into (5.82) one can see that the resulting expression is equal to the string theory result (5.60). In appendix 5.4.3 we generalize the results of this subsection to matrix elements between the generic states.

5.3 Summary and further developments

It has been known for some time [31, 32] that type IIB string theory is solvable in the plane wave background, which can be viewed as the Penrose limit of $AdS_5 \times S^5$. BMN [30] showed that the string spectrum in this background, can be recovered from the boundary $\mathcal{N}$=4 super Yang-Mills. Motivated by these results, we analyzed the properties of this correspondence when finite radius effects are included. We found that to the leading order in $1/R^2$ and $\lambda' = gN/J^2$, the string theory spectrum matches the spectrum of anomalous dimensions of (linear combinations of) BMN operators. On the string side we have an interacting worldsheet theory, when the leading $\mathcal{O}(1/R^2)$ corrections to the plane wave metric are taken into account. Leading corrections to the string spectrum can then be computed with quantum mechanical perturbation theory. On the SYM side, departing from the Penrose limit forces one to refine the BMN operators, paying attention to $1/J$ corrections. We nevertheless assume that these refined operators continue to correspond to plane wave states even away from the Penrose limit. Such operators however do not have definite scaling dimensions, when $1/J$ corrections are included. Finding the spectrum of scaling dimensions in SYM requires one to compute the matrices of two-point functions $\langle \mathcal{O}_\alpha \bar{\mathcal{O}}_\beta \rangle_{g^0} \sim \mathbf{T}_{\alpha\beta}$ and

$\langle \mathcal{O}_\alpha \bar{\mathcal{O}}_\beta \rangle_{g^1} \sim \mathbf{F}_{\alpha\beta}$. Then, $\mathbf{T}^{-1}\mathbf{F}$ is related to the light cone worldsheet Hamiltonian. We find matching between the matrix elements of these operators.

There is a number of questions raised by the results discussed here. It would be interesting to see if the correspondence between the operators we define in section 5.2.1 and plane wave states is exact and holds for arbitrary values of AdS radius. So far we matched the leading $1/R^2$, λ' terms in matrix elements of the light cone Hamiltonian. We did not include the fermionic part of the superstring in our analysis, which led to an undefined normal ordering constant in diagonal matrix elements. Incorporating fermions and extending the results of [31, 32] to $\mathcal{O}(1/R^2)$ corrections is an interesting open problem. It would also be interesting to extend our analysis to higher powers of λ'. This would require computing diagrams with multiple interactions, but perhaps one may be able to come up with a resummation technique similar to the one introduced in [30]. Extending our results to higher orders in $1/R^2$ seems more difficult technically, but might also deserve some interest.

Other possible extensions include studying backgrounds that are more complicated than $AdS_5 \times S^5$. Probing the strong coupling behavior of boundary theories with fewer supersymmetries may be of particular interest, but it remains to be seen how far one can go with this perturbative approach.

5.4 Appendix

5.4.1 An alternative worldsheet discussion

In Section 5.1 we discussed how to do the worldsheet calculations in the spirit of Polchinski [71]. In this Appendix, we explain in detail how to fix the gauges using the method described in GSW [67]. We find the same results for physical quantities as in Section 5.1.

5.4.1.1 Penrose limit of $AdS_5 \times S^5$

Before fixing any gauges, the bosonic part of the worldsheet action is

$$S = -\frac{1}{4\pi\alpha'}\int (d^2\sigma)\sqrt{-\gamma}\gamma^{ab}G_{ab} \tag{5.84}$$

where the induced metric on the worldsheet is $G_{ab} \equiv \partial_a X^\mu \partial_b X^\nu G_{\mu\nu}$. Using reparametrization invariance and Weyl invariance, we can bring the worldsheet metric to the form

$$\gamma_{ab} = \eta_{ab} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \tag{5.85}$$

in (τ, σ) coordinates. The leading order target space metric (5.3) is

$$ds_0^2 = -4dX^- dX^+ - (r^2 + y^2)dX^+ dX^+ + dX_I\, dX_I \tag{5.86}$$

$I = 1, ..., 8$. After fixing the worldsheet metric as in (5.85), the string action (5.84) becomes

$$\begin{aligned} S_0 \;&=\; -\frac{1}{4\pi\alpha'}\int (d\tau d\sigma)\Big[4\dot{X}^-\dot{X}^+ + X^2\dot{X}^+\dot{X}^+ - \dot{X}_I\dot{X}_I \\ &\qquad -4(X^-)'(X^+)' - X^2(X^+)'(X^+)' + X_I'X_I'\Big] \end{aligned} \tag{5.87}$$

The action (5.87) is not completely gauge fixed. We still have the freedom to reparameterize the worldsheet coordinates holomorphically,

$$\sigma^+ \to \tilde{\sigma}^+(\sigma^+), \quad \sigma^- \to \tilde{\sigma}^-(\sigma^-) \tag{5.88}$$

where $\sigma^\pm = \tau \pm \sigma$ are the holomorphic and antiholomorphic worldsheet coordinates. Under (5.88), the new

$$\tilde{\tau} = \frac{1}{2}\Big[\tilde{\sigma}^+(\tau + \sigma) + \tilde{\sigma}^-(\tau - \sigma)\Big] \tag{5.89}$$

satisfies the free massless wave equation

$$\ddot{\tilde{\tau}} - \tilde{\tau}'' \equiv \left[\partial_\tau^2 - \partial_\sigma^2\right]\tilde{\tau} = 0 \tag{5.90}$$

X^- enters the action (5.87) linearly, so we can integrate it out, imposing its equation of motion as a constraint. This equation is $\ddot{X}^+ - (X^+)'' = 0$, and it has the form (5.90). Hence we can choose the light-cone gauge

$$X^+ = x^+ + p^+\tau \tag{5.91}$$

This exhausts all the gauge freedom in the problem. After integrating out X^- and choosing the lightcone gauge, the action becomes

$$S_0 \;=\; -\frac{1}{4\pi\alpha'}\int (d\tau d\sigma)\left[(p^+)^2 X^2 - \dot{X}_I\dot{X}_I + X_I'X_I'\right] \tag{5.92}$$

From this, we find the lightcone Hamiltonian to be

$$H_0 \;=\; \frac{1}{4\pi\alpha'}\int_0^{2\pi} d\sigma\left[(2\pi\alpha')^2 P_I P_I + X_I'X_I' + (p^+)^2 X_I X_I\right] \tag{5.93}$$

where P_I are the momenta conjugate to X_I. The Hamiltonian (5.93) is quadratic, and can be quantized exactly. Expand the X_I and P_I in modes as

$$X_I = \sum_{n=-\infty}^{+\infty} i\sqrt{\frac{\alpha'}{2\varpi_n}}\left[a_n^I - a_n^{I\,\dagger}\right], \quad 2\pi P_I = \sum_{n=-\infty}^{+\infty}\sqrt{\frac{\varpi_n}{2\alpha'}}\left[a_n^I - a_n^{I\,\dagger}\right] \tag{5.94}$$

where the frequencies are

$$\varpi_n = \sqrt{(p^+)^2 + n^2} \tag{5.95}$$

and the oscillators

$$a_n^I = \alpha_n^I e^{-i(\varpi_n\tau - n\sigma)}, \quad a_n^{I\,\dagger} = \alpha_n^{I\,\dagger} e^{+i(\varpi_n\tau - n\sigma)} \tag{5.96}$$

close as $[\alpha_m^I, \alpha_n^{J\dagger}] = \delta^{IJ}\delta_{mn}$. In terms of these oscillators, (5.93) reads

$$H_0 \;=\; \sum_{I=1}^{8}\sum_{n=-\infty}^{+\infty}\varpi_n\left[N_n^I + \frac{1}{2}\right] \tag{5.97}$$

where the number operators are $N_n^I \equiv a_n^{I\,\dagger} a_n^I$ (no sum on either n or I). We will drop the normal ordering constants, since they cancel against the fermionic ones in the plane wave limit.

To compare space-time quantum numbers with worldsheet quantities, we look at the Noether charges associated with target space isometries. The relevant ones for us will be the energy $E = i\partial_t$, and the angular momentum $J = -i\partial_\psi$, where t and ψ are the global coordinates on AdS used in (5.1). In the dual CFT description, these correspond to the conformal dimension $\Delta = E$ and the R-charge J. We find

$$-i\frac{\partial}{\partial X^\pm} \quad \leftrightarrow \quad \mathcal{P}_\pm = \int_0^{2\pi} d\sigma P_\pm, \quad \text{where } P_\mu = \frac{\delta S}{\delta \dot{X}^\mu} = \frac{1}{2\pi\alpha'} G_{\mu\nu}\dot{X}^\nu \tag{5.98}$$

are the momenta canonically conjugate to X^μ. In the light-cone gauge,

$$\frac{\Delta + J}{R^2} \quad \leftrightarrow \quad i\frac{\partial}{\partial X^-} \leftrightarrow \; -\mathcal{P}_- \;=\; \frac{2p^+}{\alpha'} \tag{5.99}$$

$$\Delta - J \quad \leftrightarrow \quad i\frac{\partial}{\partial X^+} \leftrightarrow \; -\mathcal{P}_+ \;=\; \frac{1}{p^+}H_0 = \sum_I \sum_n \frac{\varpi_n}{p^+} N_n^I \tag{5.100}$$

Given our gauge choice (5.91), $\mathcal{P}_+$ and H_0 should differ by a factor of $-p^+$; the minus sign in (5.100) comes about because $H = i\partial_t$, while $P = -i\partial_X$. The light-cone states

$$|I_m, J_n, ...\rangle \equiv a_m^{I\,\dagger} a_n^{J\dagger} \, ... \, |0, p^+\rangle \tag{5.101}$$

have $p^+ = \frac{J}{\sqrt{4\pi g N}}$, with $R^4 = 4\pi g N \alpha'^2$; $\Delta - J = [1+\frac{m^2}{2(p^+)^2}] + [1+\frac{n^2}{2(p^+)^2}] + \mathcal{O}(\frac{1}{(p^+)^4})$.

Oscillators (5.96) explicitly depend on time, so they are Heisenberg picture operators. To go to the Schroedinger picture, we can just drop the time dependence and use the equations of motion which follow from the Hamiltonian (5.93). These are

$$a_n^I \;=\; \alpha_n^I \, e^{+in\sigma}, \quad \frac{d}{dt}a_n^I = -i\varpi_n \alpha_n^I \, e^{+in\sigma} \tag{5.102}$$

$$a_n^{I\,\dagger} \;=\; \alpha_n^{I\,\dagger} e^{-in\sigma}, \quad \frac{d}{dt}a_n^{I\,\dagger} = +i\varpi_n \alpha_n^{I\,\dagger} \, e^{-in\sigma} \tag{5.103}$$

It will be convenient to work with Heisenberg picture operators throughout, and convert the final expressions to the Schroedinger picture before doing perturbation theory.

5.4.1.2 Corrections to the Penrose limit of $AdS_5 \times S^5$

The $1/R^2$ correction to the space-time metric is given by $\frac{1}{R^2} ds_1^2$ with

$$ds_1^2 = -2dX^- dX^+ (r^2 - y^2) - \frac{1}{3}(r^4 - y^4) dX^+ dX^+ + \frac{1}{3}(r^4 d\Omega_3{}^2 - y^4 d\Omega_3'{}^2) \tag{5.104}$$

Using the identities $dr_i dr_i = dr^2 + r^2 d\Omega_3^2$ and $r dr = r_i dr_i$, we can write $r^4 d\Omega_3^2 = [r_i r_i dr_j dr_j - r_i r_j dr_i dr_j]$ and similarly for the y-s. This results in the contributions

$$X_{ab} \equiv \frac{1}{3}\left[X_i X_i \; (\partial_a X_j)(\partial_b X_j) - X_i X_j \; (\partial_a X_i)(\partial_b X_j)\right] \tag{5.105}$$

to the induced metric G_{ab}. X can be either r or y in (5.105), and the sums on the repeated i and j run from 1 to 4. The $i = j$ terms cancel in (5.105).

After fixing the worldsheet metric as in (5.85), the bosonic part of the action becomes $S = S_0 + \frac{1}{R^2} S_1$ with S_0 given in (5.87), and

$$\begin{aligned} S_1 \;&=\; -\frac{1}{4\pi\alpha'} \int (d\tau d\sigma) \Big\{ -2X^- \left[\partial_\tau(\dot{X}^+(r^2 - y^2)) - \partial_\sigma((X^+)'(r^2 - y^2))\right] \\ &\qquad + \frac{1}{3}\left[\dot{X}^+ \dot{X}^+ - (X^+)'(X^+)'\right](r^4 - y^4) - (r_{\tau\tau} - y_{\tau\tau}) + (r_{\sigma\sigma} - y_{\sigma\sigma}) \Big\} \end{aligned} \tag{5.106}$$

(we integrated by parts so that derivatives of X^- do not appear in S_1). Since the variable X^- appears linearly in the action S, we can integrate it out, and impose its equation of motion as a constraint. Although this equation is no longer linear, it can be solved perturbatively in $1/R^2$. Writing

$$X^+(\tau, \sigma) = X_0^+ + \frac{1}{R^2} X_1^+ \tag{5.107}$$

where $X^+_{0,1}$ are both of order one, we get

$$\begin{aligned} 0 &= \ddot{X}^+_0 - (X^+_0)'' \\ &\quad + \frac{1}{R^2}\left\{ \ddot{X}^+_1 - (X^+_1)'' + \partial_\tau \left[\dot{X}^+_0 \left(\frac{r^2 - y^2}{2} \right) \right] - \partial_\sigma \left[(X^+_0)' \left(\frac{r^2 - y^2}{2} \right) \right] \right\} \end{aligned} \tag{5.108}$$

Since X^+_0 satisfies the free massless wave equation, we can take $X^+_0 = x^+ + p^+\tau$. Thus the (modified) light-cone gauge choice is

$$X^+(\tau,\sigma) = \left(x^+ + p^+\tau \right) + \frac{1}{R^2} X^+_1 \tag{5.109}$$

To completely fix the gauge, we have to make sure that contributions of the form $x^+_1 + p^+_1\tau$ and $e^{in(\tau\pm\sigma)}$, are absent in the mode expansion of $X^+_1(\tau,\sigma)$. In terms of the original coordinate X^+ and the original τ and σ, this is a statement that

$$\frac{1}{R^2} X^+_1(\tau,\sigma) \equiv X^+(\tau,\sigma) - \frac{1}{2\pi} \sum_{n,\pm} e^{in(\tau\pm\sigma)} \int (d\sigma d\tau) X^+(\tau,\sigma) e^{-in(\tau\pm\sigma)} \tag{5.110}$$

The leftover piece X^+_1 is not a new dynamical variable; rather, it depends on r_i and y_i. It is defined to satisfy

$$\ddot{X}^+_1 - (X^+_1)'' + \frac{1}{2} p^+ \partial_\tau \left(r^2 - y^2 \right) = 0 \tag{5.111}$$

The r^2 and y^2 should be taken as their leading order versions (5.94). Setting $r^i_n \equiv a^i_n$ and $y^i_n \equiv a^{i+4}_n$ in the mode expansions (5.94), we find

$$\begin{aligned} X^+_1 &= \frac{ip^+\alpha'}{2} \sum_{m\,,\,n} \frac{\varpi_n}{\sqrt{\varpi_m \varpi_n}} \frac{\left[(r^i_m r^i_n - r^i_m{}^\dagger r^i_n{}^\dagger) - (y^i_m y^i_n - y^i_m{}^\dagger y^i_n{}^\dagger) \right]}{[(\varpi_m + \varpi_n)^2 - (m+n)^2]} \\ &+ \frac{ip^+\alpha'}{2} \sum_{m\neq n} \frac{\varpi_n}{\sqrt{\varpi_m \varpi_n}} \frac{\left[(r^i_m r^i_n{}^\dagger - r^i_m{}^\dagger r^i_n) - (y^i_m y^i_n{}^\dagger - y^i_m{}^\dagger y^i_n) \right]}{[(\varpi_m - \varpi_n)^2 - (m-n)^2]} \end{aligned} \tag{5.112}$$

Equation (5.111) is solved in Heisenberg picture; the operator X^+_1 is determined in terms of the Heisenberg picture oscillators (5.96). Since (5.112) contains no explicit time dependence, it can be interpreted as a Schroedinger picture expression (when the oscillators are taken to be in Schroedinger picture), and used in perturbative calculations of energies.

The action in the modified lightcone gauge (5.109) reads

$$\begin{aligned} S \;=\; & -\frac{1}{4\pi\alpha'}\int(d\tau d\sigma)\Big\{(p^+)^2(r^2+y^2)-\dot{r}_i\dot{r}_i-\dot{y}_i\dot{y}_i+r'_ir'_i+y'_iy'_i \\ & \qquad +\frac{1}{R^2}\Big[(r_{\sigma\sigma}-y_{\sigma\sigma})-(r_{\tau\tau}-y_{\tau\tau}) \\ & \qquad\qquad +\frac{1}{3}(p^+)^2(r^4-y^4)+2p^+\dot{X}_1^+(r^2+y^2)\Big]\Big\} \end{aligned} \tag{5.113}$$

after integrating out X^-, i.e. after solving the constraint equation (5.108). As discussed in [47], the first order correction to the Hamiltonian is minus the correction to the Lagrangian, $\delta H = -\delta L$. Hence the (modified-)lightcone Hamiltonian is

$$\begin{aligned} H \;=\; & \frac{1}{4\pi\alpha'}\int_0^{2\pi}d\sigma\Big\{\Big[(2\pi\alpha')^2(P_i^rP_i^r+P_i^yP_i^y)+(r'_ir'_i+y'_iy'_i)+(p^+)^2(r^2+y^2)\Big] \\ & \qquad +\frac{1}{R^2}\Big[(r_{\sigma\sigma}-y_{\sigma\sigma})-(r_{\tau\tau}-y_{\tau\tau}) \\ & \qquad\qquad +\frac{1}{3}(p^+)^2(r^4-y^4)+2p^+\dot{X}_1^+(r^2+y^2)\Big]\Big\} \end{aligned} \tag{5.114}$$

with X_1^+ given in (5.112).

The conserved charges corresponding to $\frac{\Delta+J}{R^2}$ and $\Delta - J$ are

$$-\mathcal{P}_- \;=\; \frac{2p^+}{\alpha'}+\frac{4p^+}{4\pi\alpha'}\int_0^{2\pi}d\sigma\,\frac{1}{R^2}\left(\frac{\dot{X}_1^+}{p^+}+\frac{r^2-y^2}{2}\right) \tag{5.115}$$

$$-\mathcal{P}_+ \;=\; \frac{1}{p^+}H \tag{5.116}$$

In terms of the (Schroedinger picture) oscillators,

$$-\mathcal{P}_- \;=\; \frac{2p^+}{\alpha'}\left\{1+\frac{\alpha'}{2R^2}\left[\sum_{i=1}^{4}\sum_{n=-\infty}^{+\infty}\frac{1}{\varpi_n}(N_n^{ri}-N_n^{yi})\right]\right\} \tag{5.117}$$

Corrections of the form aa and $a^\dagger a^\dagger$ precisely cancels between $\frac{1}{p^+}\dot{X}_1^+$ and $\frac{1}{2}(r^2-y^2)$ in (5.115). For $p^+ \gg 1$, the worldsheet parameter p^+ is related to J and N as

$$p^+ \;=\; \frac{J}{\sqrt{4\pi gN}}\left\{1+\frac{1}{J}\sum_{i=1}^{4}\sum_{n=-\infty}^{+\infty}\left[N_n^{yi}+\frac{2\pi gNn^2}{J^2}N_n^{ri}\right]\right\} \tag{5.118}$$

to order $1/R^2$. Here we used $R^4 = 4\pi gN\alpha'^2$, and wrote $(\Delta+J) = 2J+(\Delta-J)$. In (5.118) the contributions of the y and r oscillators have rather different structure.

The Hamiltonian (5.114) is relatively involved, so we analyze it in more detail. The leading order lightcone string states

$$|a_p, b_q, ...\rangle = y_p^{a\dagger} y_q^{b\dagger}\; ...\; |0, p^+\rangle \tag{5.119}$$

with worldsheet momenta $(p, q, ...)$ and $(p', q', ...)$ are degenerate only when the $(p, q, ...)$ and $(p', q', ...)$ are permutations of one another. Hence the only terms in δH relevant for computing the first correction to the worldsheet energies, are the ones which permute the worldsheet momenta, namely $a_k a_k^\dagger a_k a_k^\dagger$ and $a_k a_k^\dagger a_l a_l^\dagger$.

Such terms in $\left[(r_{\sigma\sigma}-y_{\sigma\sigma})-(r_{\tau\tau}-y_{\tau\tau})+\frac{1}{3}(p^+)^2(r^4-y^4)\right]$ combine as

$$\begin{aligned}
& 2\left(\frac{\alpha'}{2}\right)^2\sum_k\frac{(p^+)^2}{\varpi_k^2}[r_k^i r_k^i r_k^{j\dagger} r_k^{j\dagger} - y_k^i y_k^i y_k^{j\dagger} y_k^{j\dagger}] \\
+ \;\; & 2\left(\frac{\alpha'}{2}\right)^2\sum_{k\neq l}\frac{(p^+)^2+\varpi_k\varpi_l-kl}{\varpi_k\varpi_l}[r_k^i r_l^i r_l^{j\dagger} r_k^{j\dagger} - y_k^i y_l^i y_k^{l\,\dagger} y_k^{j\dagger}] \\
+ \;\; & 2\left(\frac{\alpha'}{2}\right)^2\sum_{k\neq l}\frac{(p^+)^2-\varpi_k\varpi_l+kl}{\varpi_k\varpi_l}[r_k^i r_l^j r_l^{i\dagger} r_k^{j\dagger} - y_k^i y_l^j y_l^{i\dagger} y_k^{j\dagger}]
\end{aligned} \tag{5.120}$$

and the term $2p^+\dot{X}_1^+(r^2+y^2)$ gives

$$\begin{aligned}
- \;\; & 2\left(\frac{\alpha'}{2}\right)^2\sum_k[r_k^i r_k^i r_k^{j\dagger} r_k^{j\dagger} - y_k^i y_k^i y_k^{j\dagger} y_k^{j\dagger}] \\
- \;\; & 2\left(\frac{\alpha'}{2}\right)^2\sum_{k\neq l}\frac{2(p^+)^2(\varpi_k+\varpi_l)^2}{\varpi_k\varpi_l[(\varpi_k+\varpi_l)^2-(k+l)^2]}[r_k^i r_l^i r_l^{j\dagger} r_k^{j\dagger} - y_k^i y_l^i y_k^{l\,\dagger} y_k^{j\dagger}] \\
- \;\; & 2\left(\frac{\alpha'}{2}\right)^2\sum_{k\neq l}\frac{2(p^+)^2(\varpi_k-\varpi_l)^2}{\varpi_k\varpi_l[(\varpi_k-\varpi_l)^2-(k-l)^2]}[r_k^i r_l^j r_l^{i\dagger} r_k^{j\dagger} - y_k^i y_l^j y_l^{i\dagger} y_k^{j\dagger}]
\end{aligned} \tag{5.121}$$

Expressions (5.120)-(5.121) appear in $\frac{1}{p^+}\delta H$ with an overall prefactor of

$$\frac{1}{4\pi\alpha'} \cdot 2\pi \cdot \frac{1}{R^2} \cdot \frac{1}{p^+} = \frac{1}{2\alpha' R^2 p^+} \tag{5.122}$$

and we find

$$\begin{aligned} \frac{1}{p^+}\delta H &= \frac{\alpha'}{4R^2(p^+)^3} \sum_{i,j} \sum_{k} \frac{k^2(p^+)^2}{\varpi_k^2} (y^i_k y^i_k y^{j\dagger}_k y^{j\dagger}_k - r^i_k r^i_k r^{j\dagger}_k r^{j\dagger}_k) \\ &+ \frac{\alpha'}{4R^2(p^+)^3} \sum_{i,j} \sum_{k \neq l} \frac{-2kl(p^+)^2}{\varpi_k \varpi_l} (y^i_k y^{i\dagger}_l y^j_l y^{j\dagger}_k - r^i_k r^{i\dagger}_l r^j_l r^{j\dagger}_k) \\ &+ \frac{\alpha'}{4R^2(p^+)^3} \sum_{i,j} \sum_{k \neq l} \frac{2kl(p^+)^2}{\varpi_k \varpi_l} (y^i_k y^{j\dagger}_k y^i_l y^{j\dagger}_l - r^i_k r^{j\dagger}_k r^i_l r^{j\dagger}_l) \\ &+ \ ... \end{aligned} \tag{5.123}$$

The "..." stands for terms not of the form $aa^\dagger aa^\dagger$, as well as terms with more than two distinct worldsheet momenta; we are also dropping corrections which are higher order in $1/R^2$ and $1/p^+$. The second and third lines of (5.123) cancel if $i = j$.

In deriving (5.120)-(5.121), we have not been careful about the ordering of oscillators. This means that we may have overlooked some terms which involve commutators $[y^i_m, {y^i_m}^\dagger] = 1$. The only terms in (5.123) where this could happen come from the first line. This means we could be possibly neglecting

$$-\delta'\mathcal{P}_+ = \left(\frac{\alpha'}{4R^2(p^+)^3}\right) \zeta \sum_{i} \sum_{k} \frac{k^2(p^+)^2}{\varpi_k^2} (N^{yi}_k - N^{ri}_k) \tag{5.124}$$

If we were to keep track of the ordering of oscillators, we would find $\zeta = 1$. However, we have not analyzed the fermionic side, which can also produce similar terms.

Finally, we compare the results of this Appendix with what we found in Section 5.1. We will only look at the y-oscillators. The difference between p^+ and η is

$$p^+ = \eta \left[1 + \frac{1}{2R^2} \sum_{i;n} \frac{N^{yi}_n}{\varpi_n}\right] \tag{5.125}$$

so the frequencies in the two approaches are related as

$$\varpi_m = w_m \left[1 + \frac{\eta^2}{2R^2 w_m^2} \sum_{i;n} \frac{N_n^{yi}}{w_n} + \mathcal{O}(1/R^4) \right] \tag{5.126}$$

Expressions (5.123) and (5.124) then change trivially as $\varpi_n \to w_n$, $p^+ \to \eta$ at this order in $1/R^2$, while

$$\frac{1}{p^+} \sum_{i;n} \varpi_n N_n^{yi} = \frac{1}{\eta} \sum_{i;n} w_n N_n^{yi} - \frac{1}{2R^2 \eta} \sum_{i,j;m,n} \frac{n^2 N_n^{yi} N_m^{yj}}{w_m w_n} + \mathcal{O}(1/R^4) \tag{5.127}$$

Together, (5.123) and (5.127) reproduce the sum of (5.14), (5.22) and (5.23).

5.4.2 $\mathcal{N}$=4 SYM

Here, we give some details of the $\mathcal{N}$=4 SYM needed for the order $g^0_{\rm YM}$ (tree) and $g^2_{\rm YM}$ (one-loop level) calculations of Section (5.2.2). First we write down the $\mathcal{N}$=4 SYM action in terms of the fields we will be dealing with. When SUSY is broken down to $\mathcal{N}$=1, things much more cumbersome, so from the very beginning we use the $\mathcal{N}$=4 Lagrangian 2.89 from Chapter 2,

$$\begin{aligned} \mathcal{L} \ = \ & \tfrac{1}{g^2_{\rm YM}} \, \mathrm{tr} \, \Big\{ -\tfrac{1}{4} F_{\mu\nu} F^{\mu\nu} + i\lambda \sigma^\mu D_\mu \bar{\lambda} + i\psi_j \sigma^\mu D_\mu \bar{\psi}^j + D_\mu z_j D^\mu \bar{z}^j \\ & + i\sqrt{2} [\lambda, \psi_j] \bar{z}^j - \tfrac{i}{\sqrt{2}} \epsilon^{jkl} [\psi_j, \psi_k] z_l + i\sqrt{2} [\bar{\lambda}, \bar{\psi}^j] z_j - \tfrac{i}{\sqrt{2}} \epsilon_{jkl} [\bar{\psi}^j, \bar{\psi}^k] \bar{z}^l \\ & + [z_j, z_k][\bar{z}^j, \bar{z}^k] - \tfrac{1}{2} [z_j, \bar{z}^j][z_k, \bar{z}^k] \Big\} . \end{aligned} \tag{5.128}$$

We leave the fields z_1, $\bar{z}^1$ as they are, and substitute

$$z_j = \tfrac{1}{\sqrt{2}} \left(\phi_j + i\phi_{j+3} \right), \quad \bar{z}^j = \tfrac{1}{\sqrt{2}} \left(\phi_j - i\phi_{j+3} \right), \quad j = 2 \text{ and } 3. \tag{5.129}$$

The rest of the fields (gauge bosons and fermions) remain unchanged, and (5.128) becomes

$$\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_{\rm other} \tag{5.130}$$

where

$$\mathcal{L}_0 = \tfrac{1}{g_{\mathrm{YM}}^2}\,\mathrm{tr}\left\{(\partial_\mu z_1)(\partial^\mu \bar{z}_1) + \sum_k \tfrac{1}{2}(\partial_\mu \phi_k)(\partial^\mu \phi_k)\right\} \tag{5.131}$$

gives propagators for the scalars;

$$\begin{aligned}\mathcal{L}_1 = \tfrac{1}{g_{\mathrm{YM}}^2}\,\mathrm{tr}\Big\{ & -iA^\mu[z_1, \partial_\mu \bar{z}_1] - iA^\mu[\bar{z}_1, \partial_\mu z_1] + \sum_k (-iA^\mu)[\phi_k, \partial_\mu \phi_k] \\ & + i\sqrt{2}\, z_1 \left([\bar{\lambda}, \bar{\psi}^1] - [\psi_2, \psi_3]\right) + i\sqrt{2}\, \bar{z}_1 \left([\lambda, \psi_1] - [\bar{\psi}^2, \bar{\psi}^3]\right) \\ & + i\phi_2 \left([\lambda, \psi_2] + [\bar{\lambda}, \bar{\psi}^2] - [\psi_3, \psi_1] - [\bar{\psi}^3, \bar{\psi}^1]\right) \\ & + \phi_5 \left([\lambda, \psi_2] - [\bar{\lambda}, \bar{\psi}^2] + [\psi_3, \psi_1] - [\bar{\psi}^3, \bar{\psi}^1]\right) \\ & + i\phi_3 \left([\lambda, \psi_3] + [\bar{\lambda}, \bar{\psi}^3] - [\psi_1, \psi_2] - [\bar{\psi}^1, \bar{\psi}^2]\right) \\ & + \phi_6 \left([\lambda, \psi_3] - [\bar{\lambda}, \bar{\psi}^3] + [\psi_1, \psi_2] - [\bar{\psi}^1, \bar{\psi}^2]\right)\Big\}\end{aligned} \tag{5.132}$$

gives 3-field vertices; and

$$\begin{aligned}\mathcal{L}_2 = \tfrac{1}{g_{\mathrm{YM}}^2}\,\mathrm{tr}\Big\{ & -[A_\mu, z_1][A^\mu, \bar{z}_1] - \sum_k \tfrac{1}{2}[A_\mu, \phi_k][A^\mu, \phi_k] \\ & -\tfrac{1}{2}[z_1, \bar{z}_1][z_1, \bar{z}_1] + \sum_k [z_1, \phi_k][\bar{z}_1, \phi_k] + \sum_{k>l} \tfrac{1}{2}[\phi_k, \phi_l][\phi_k, \phi_l]\Big\}\end{aligned} \tag{5.133}$$

contains 4-field interactions. Finally,

$$\mathcal{L}_{\mathrm{other}} = \tfrac{1}{g_{\mathrm{YM}}^2}\,\mathrm{tr}\left\{-\tfrac{1}{4}F_{\mu\nu}F^{\mu\nu} + i\lambda\sigma^\mu D_\mu \bar{\lambda} + i\psi_j \sigma^\mu D_\mu \bar{\psi}^j\right\} \tag{5.134}$$

gives propagators for the gauge bosons and the fermions and their interactions with each other (at order $\mathcal{O}(g_{\mathrm{YM}}^2)$ these do not contribute to the diagrams we care about, and neither do the ghost terms). The Lagrangian (5.130) has a leftover $SO(4)$ symmetry rotating the ϕ-s.

Feynman rules for the Lagrangian (5.130) are somewhat awkward, but the tree and one-loop diagrams which involve only the scalars can be packaged in a convenient way. First, $\mathcal{O}(g_{\mathrm{YM}}^2)$ corrections to the scalar propagators are diagonal


z_1^a
$\bar{z}_1^b$
$= \delta^{ab} \; N A(x,y) G(x,y)$
ϕ_i^a
ϕ_j^b
$= \delta^{ab}\delta^{ij} \; N A(x,y) G(x,y)$

Figure 5.1: Order $g_{\rm YM}^2$ corrections to scalar propagators consist of a gauge boson exchange and a fermion loop.

in color indices, see Figure 5.1. Fermion loops cancel in $\langle \phi_2^a(x)\phi_5^b(y)\rangle_{g_{\rm YM}^2}$ because of the way the signs work out in (5.132).

Corrections to the 4-point irreducible blocks are more involved, but they can be related to the corresponding diagrams involving only z-s and $\bar{z}$-s. By comparing two-point functions of the protected operators in the [0,2,0] of $SU(4)$ written on the one hand in terms of ϕ-fields, and on the other hand in terms of z-s and $\bar{z}$-s, we get the diagrams shown in Figure 5.2. Comparison of two-point functions of the Konishi scalar $\sum_{k=1}^{6} \operatorname{tr} \phi^k \phi^k = \sum_{k=1}^{3} \operatorname{tr} z^k \bar{z}^k$ produce the relations listed in Figure 5.3.

The "D-term" contributions A and B, and the four-field interaction "F-term" $\tilde{B}$ are defined by Figures 5.2 and 5.3. As we saw in Section 2.5 (also see [41]), the A and B are not separately gauge invariant. These must appear as the gauge invariant combination $2A + B$, which vanishes in the $\mathcal{N}$=4 theory. So one only has to look at "F-term" contributions, which are all proportional to

$$\gamma \equiv \tfrac{1}{2}\tilde{B}(x,0)N = -\frac{g_{\rm YM}^2 N}{4\pi^2} \log x^2\mu^2 \equiv -\beta \log x^2\mu^2 \qquad (5.135)$$

computed for example in Chapter 2 and in [30]. In this Chapter, we are using the conventions of [30]; in the Lagrangian (5.128) we have $g_{\rm YM}^2 = 2\pi g$.

We only have to consider planar diagrams since we are interested in the leading



i, a
i, a'
j, b
j, b'
+
=
= f^{apa'} f^{bpb'} B (x,y) G(x,y)^2
= f^{apb} f^{a'pb'} B̃ (x,y) G(x,y)^2

Figure 5.2: Order $g_{\rm YM}^2$ corrections to two point functions of operators of the form $\mathrm{tr}\, z ... \phi^1 z ... \phi^2$: four-field irreducible blocks. When scalars ϕ^i are involved, the diagrams above represent the net contribution of all contributing Feynman diagrams, packaged in a way to mimic the $\mathcal{N}=1$ component fields Feynman diagrams. (Thick lines would correspond to exchanges of auxiliary fields F_i and D in the $\mathcal{N}=1$ formulation.) Diagrams with z_2 are given for comparison only. There are similar diagrams with one or both z-lines running in the opposite direction.



φ_i^a
φ_i^b
φ_j^{a'}
φ_j^{b'}
i ≠ j
=
z̄_1^{b'}
z_1^{a'}
φ_i^{a'}
φ_i^{b'}
= (f^{apa'} f^{bpb'} + f^{bpa'} f^{apb'}) ½ B̃(x,y) G(x,y)^2
= (f^{apa'} f^{bpb'} + f^{bpa'} f^{apb'}) (B(x,y)+½ B̃(x,y)) G(x,y)^2

Figure 5.3: Order g^2 corrections to two point functions of operators of the form $\mathrm{tr}\, z ... \phi^1 z ... \phi^1$: four-field irreducible blocks. These diagrams represent the net contribution of all contributing Feynman diagrams.

large N behavior. Put differently,

$$\operatorname{tr}\left[t^{a_1}...t^{a_k}\right]\ \operatorname{tr}\left[t^{a_k}...t^{a_1}\right]=\left(\frac{N}{2}\right)^k\left[1+\mathcal{O}(1/N^2)\right] \tag{5.136}$$

and $SU(N)$ traces of all other permutations of the generators (other than cyclic) are suppressed by $1/N^2$. To see this, one can use the "trace merging formula"

$$2\left(\operatorname{tr}At^c\right)\left(\operatorname{tr}Bt^c\right)=\operatorname{tr}AB-\tfrac{1}{N}\left(\operatorname{tr}A\right)\left(\operatorname{tr}B\right) \tag{5.137}$$

valid when t^c are $SU(N)$ generators in the fundamental representation.

At one loop, all but the nearest neighbor interactions are suppressed. The relevant contributions in Figure 5.2 have the form



$$\begin{aligned}
&= \operatorname{tr}\left[t^at^bt^{c_1}...t^{c_J}\right]\ \operatorname{tr}\left[t^{c_J}...t^{c_1}t^{b'}t^{a'}\right]\ f^{abp}f^{a'b'p}\\
&= \tfrac{1}{2}\left(\tfrac{1}{2}N\right)^{J-1}\operatorname{tr}\left(t^at^bt^{b'}t^{a'}\right)f^{abp}f^{a'b'p}\left[1+\mathcal{O}(1/N^2)\right]\\
&= \tfrac{1}{2}\left(\tfrac{1}{2}N\right)^{J-1}\operatorname{tr}\left(t^a[t^p,t^a]t^{b'}[t^p,t^{b'}]\right)\left[1+\mathcal{O}(1/N^2)\right]\\
&= \left(\tfrac{1}{2}N\right)^{J+3}\left[1+\mathcal{O}(1/N^2)\right]
\end{aligned} \tag{5.138}$$

The difference between the orderings (ab) and (ba) in (5.138) is a minus sign,



$$=-\tag{5.139}$$



Diagrams shown in the first two lines of Figure 5.3 have the form



$$\begin{aligned}
&= \tfrac{1}{2}[f^{aa'p}f^{bb'p}+f^{ab'p}f^{ba'p}]\ \operatorname{tr}\left[t^at^bt^{c_1}...t^{c_J}\right]\ \operatorname{tr}\left[t^{c_J}...t^{c_1}t^{b'}t^{a'}\right]\\
&= \tfrac{1}{2}\left(\tfrac{1}{2}N\right)^{J+3}\left[1+\mathcal{O}(1/N^2)\right]
\end{aligned} \tag{5.140}$$

Only one of the two ff-terms contributes at this order; the other one is suppressed by at least $1/N^3$. The contribution (5.140) is insensitive to $a \leftrightarrow b$. The contributions (5.138) and (5.140) come with a numerical prefactor of

$$\frac{2}{N}G(x,0)^{J+2}\gamma \tag{5.141}$$

with $\gamma = -\beta \log x^2\mu^2$ defined in (5.135).

To summarize, the tree level correlators are



$$= (\tfrac{1}{2}GN)^{J+N} \tag{5.142}$$

and the relevant one-loop contributions can be schematically represented as



$$= -$$



$$= \gamma \times (\tfrac{1}{2}GN)^{J+N} \tag{5.143}$$

when only one ϕ is involved in the interaction, and



$$= -$$



$$= \gamma \times (\tfrac{1}{2}GN)^{J+N}, \quad i \neq j \tag{5.144}$$

when two distinct ϕ within either trace interact. Furthermore, we have



$$=$$



$$=$$



$$= \frac{1}{2}\gamma \times (\tfrac{1}{2}GN)^{J+N}, \quad i \neq j \tag{5.145}$$

Finally, the diagrams which involve a $zz\bar{z}\bar{z}$ vertex can be read off from (5.144) and (5.145) by expanding the z and $\bar{z}$ participating in the vertex in terms of the two remaining ϕ's,



$$= -\frac{1}{2}\gamma \times (\tfrac{1}{2}GN)^{J+N},$$



$$= \frac{3}{2}\gamma \times (\tfrac{1}{2}GN)^{J+N} \tag{5.146}$$

In the results (5.142)-(5.146), we dropped terms suppressed by $1/N^2$.

5.4.3 Equality of matrix elements: generic states

In this appendix we will complete matching the matrix elements of the light cone Hamiltonian between the two sides of the AdS/CFT correspondence. In section 5.2.3 we matched matrix elements for a subset of states. There we considered states with all excited modes having distinct SO(4) indices i_k, and no mode excited more than once. We will now consider states with some i_k being equal. We initially restrict to the case with no modes excited more than once, $N^i_n \leq 1$, but will eventually generalize to most general case.

In contrast with section 5.2.3, $\mathcal{O}_*$ no longer vanishes. In addition to (5.58), we now have to consider off-diagonal elements between the states

$$|\mathcal{Y}\rangle = y^{i_1\,\dagger}_{n_1}\dots y^{i\,\dagger}_{m} y^{i\,\dagger}_{n}|\eta\rangle, \qquad m \neq n, \tag{5.147}$$

and

$$|\mathcal{Y}'\rangle = y^{i_1\,\dagger}_{n_1}\dots y^{j\,\dagger}_{m} y^{j\,\dagger}_{n}|\eta\rangle, \qquad m \neq n, \tag{5.148}$$

which are given by

$$\langle\mathcal{Y}|H_1^{OD}|\mathcal{Y}'\rangle = \frac{1}{J}\left(\frac{R^2}{J}\right)^2 mn\sqrt{N^i_m N^i_n N^j_m{}' N^j_n{}'}. \tag{5.149}$$

There is also an off-diagonal element given by (5.58), but we have analyzed all diagrams contributing to it in section 5.2.3. Let us briefly explain why this is the case. Consider $\mathcal{O}(g^0)$ part of the contributing two-point function, which we denote by $\langle\mathcal{X}\bar{\mathcal{X}}'\rangle_{g^0}$. $\langle\tilde{\mathcal{X}}\bar{\mathcal{X}}'_*\rangle_{g^0}$ and $\langle\mathcal{X}'_*\bar{\tilde{\mathcal{X}}}\rangle_{g^0}$ vanish, as there are no contributing interaction-free diagrams. Although $\langle\mathcal{X}_*\bar{\mathcal{X}}'_*\rangle_{g^0}$ has nonvanishing terms, they are $\mathcal{O}(1/J^2)$. This is because $\mathcal{X}_*$ is itself $\mathcal{O}(1/J)$ compared to $\tilde{\mathcal{O}}$, and an additional factor of $1/J$ will appear because phases in $\mathcal{X}_*$ and $\mathcal{X}'_*$ do not match exactly. Similar conclusions can be made about $\mathcal{O}(g)$ correlator $\langle\tilde{\mathcal{X}}\bar{\mathcal{X}}'\rangle_{g^1}$.

Let us compute the off-diagonal element (5.149) in the gauge theory. The only contribution to $\langle\mathcal{Y}(x)\bar{\mathcal{Y}}'(0)\rangle_{g^0}$ comes from

$$\langle\mathcal{Y}_*(x)\bar{\mathcal{Y}}'_*(0)\rangle_{g^0} = \frac{1}{\Omega}\sum$$



$$= \frac{1}{J}, \tag{5.150}$$

where the sum runs over all configurations of fields. No $\mathcal{O}(g^0)$ diagrams appear in $\langle\tilde{\mathcal{Y}}\bar{\mathcal{Y}}'_*\rangle_{g^0}$, $\langle\mathcal{Y}_*\bar{\tilde{\mathcal{Y}}}'\rangle_{g^0}$ and $\langle\tilde{\mathcal{Y}}\bar{\tilde{\mathcal{Y}}}'\rangle_{g^0}$. Hence we have

$$\mathbf{T}^{(1)}_{\mathcal{Y}\mathcal{Y}'} = 1. \tag{5.151}$$

Computation of $\mathbf{F}_{\mathcal{Y}\mathcal{Y}'}$ is more involved. Possible contributions are

$$\langle\tilde{\mathcal{Y}}(x)\bar{\tilde{\mathcal{Y}}}'(0)\rangle_{g^1} = \frac{1}{\Omega}\sum$$



$$+(m\leftrightarrow n)+(m'\leftrightarrow n')+(m\leftrightarrow n, m'\leftrightarrow n')$$

$$= \frac{\gamma}{2J}(q_{n-m}+q^*_{n-m}+2), \tag{5.152}$$

which holds both for $m'=n, n'=m$ (off-diagonal) and $m'=m, n'=n$ (diagonal),

$$-\langle\tilde{\mathcal{Y}}(x)\bar{\mathcal{Y}}'_*(0)\rangle_{g^1} = -\frac{1}{\Omega}\sum$$



$$+(m\leftrightarrow n)+(z\leftrightarrow\bar{z})+(m\leftrightarrow n, z\leftrightarrow\bar{z})$$

$$= -\frac{\gamma}{2J}(q_m+q^*_m+q_n+q^*_n), \tag{5.153}$$

similar contribution from $\langle\mathcal{Y}_*(x)\bar{\mathcal{Y}}'_*(0)\rangle_{g^1}$, and

$$\langle\mathcal{Y}_*(x)\bar{\mathcal{Y}}'_*(0)\rangle_{g^1} = \frac{1}{\Omega}\sum$$



$$+(\phi^{i_k}_{n_k}\leftrightarrow z)+(\phi^{i_k}_{n_k}\leftrightarrow\bar{z})+(\phi^{i_k}_{n_k}\leftrightarrow z, \phi^{i_k}_{n_k}\leftrightarrow\bar{z})$$

$$+ \frac{1}{\Omega}\sum$$



$$+(z\leftrightarrow\bar{z}_{m+n})+(\bar{z}\leftrightarrow z_{m+n})+(z\leftrightarrow\bar{z}_{m+n}, \bar{z}\leftrightarrow z_{m+n})$$

$$= \frac{\gamma}{J}\left(-\sum_{p:\, n_p\neq m,n}(q_{n_p}+q^*_{n_p}-2)-\frac{1}{2}(q_{m+n}+q^*_{m+n})+3\right). \tag{5.154}$$

(The subscript in $\bar{z}_{n+m}$ stands for the phase $q^{a_{\bar{z}}}_{n+m}$ which depends on the position of the $\bar{z}$ in the string of operators.) Combining (5.151)–(5.154) we have

$$[\mathbf{T}^{-1}\mathbf{F}]_{\mathcal{Y}\mathcal{Y}'} = -\frac{\beta}{2J}(q_{m+n}-q_{m-n}+c.c.) = \frac{1}{J}\left(\frac{R^2}{J}\right)^2 mn, \tag{5.155}$$

which indeed agrees with (5.149), provided $N^i_n \le 1$.

Let us now turn to the diagonal matrix element. Part of it was computed in section 5.2.3 and is given by (5.82). But now there are other contributions both to $\mathbf{T}^{(1)}_{\mathcal{O}\mathcal{O}}$ and to $\mathbf{F}_{\mathcal{O}\mathcal{O}}$. To update the former, we must take into account

$$\langle\mathcal{O}_*(x)\bar{\mathcal{O}}_*(0)\rangle_{g^0} = \frac{1}{\Omega}\sum_{i,(m\neq n)}\sum$$



$$= \sum_i \frac{\mathsf{N}_i(\mathsf{N}_i-1)}{2J}, \tag{5.156}$$

and

$$\delta\langle\tilde{\mathcal{O}}(x)\bar{\tilde{\mathcal{O}}}(0)\rangle_{g^0} = \frac{1}{\Omega}\sum_{i,(m\neq n)}\sum$$



$$= -\sum_i \frac{\mathsf{N}_i(\mathsf{N}_i-1)}{2J}, \tag{5.157}$$

which cancels (5.156) to keep $\mathbf{T}_{\mathcal{O}\mathcal{O}} = 1$. The $\mathcal{O}(g^1)$ correlators related to (5.156) are given by the sum of (5.154) over pairs $(n_k \neq n_l) : i_k = i_l$ with the substitution $m = n_k, n = n_l$:

$$\begin{aligned}
\langle\mathcal{O}_*(x)\bar{\mathcal{O}}_*(0)\rangle_{g^1} &= \frac{\gamma}{J}\sum_{(n_k\neq n_l):i_k=i_l}\left(3-\sum_{p\neq k,l}(q_{n_p}+q^*_{n_p}-2)-\frac{1}{2}(q_{n_k+n_l}+q^*_{n_k+n_l})\right)\\
&= \frac{\gamma}{J}\Bigg[-\left(\sum_i\frac{\mathsf{N}_i(\mathsf{N}_i-1)}{2}\right)\sum_k(q_{n_k}+q^*_{n_k}-2)\\
&\quad+\sum_{(n_k\neq n_l):i_k=i_l}\left\{(q_{n_k}+q_{n_l}-2+c.c.)-\frac{1}{2}(q_{n_k+n_l}+q^*_{n_k+n_l})+3\right\}\Bigg].
\end{aligned} \tag{5.158}$$

The $\mathcal{O}(g^1)$ counterpart of (5.157) is

$$\begin{aligned}
\delta_1\langle\tilde{\mathcal{O}}(x)\bar{\tilde{\mathcal{O}}}(0)\rangle_{g^1} &= \frac{\gamma}{J}\Bigg[\left(\sum_i\frac{\mathsf{N}_i(\mathsf{N}_i-1)}{2}\right)\sum_k(q_{n_k}+q^*_{n_k}-2)\\
&\qquad+\sum_{(n_k\neq n_l):i_k=i_l}(q_{n_k}+q_{n_l}-2q_{n_l-n_k}+c.c.)\Bigg].
\end{aligned} \tag{5.159}$$

The first term in this expression is a value of the corresponding interaction-free diagram times the sum of possible phases, while the second term takes care of overcounted corrections (this technique for computing $\mathcal{O}(g^1)$ diagrams was explained in more detail in section 5.2.3) There is also a contribution which is a direct analog of (5.80)

$$\delta_2 \langle \tilde{\mathcal{O}}(x) \bar{\tilde{\mathcal{O}}}(0) \rangle_{g^1} = \frac{\gamma}{J} \sum_{(n_k \neq n_l):i_k = i_l} \left[(q_{n_k} + q_{n_l} - 2 + c.c.) + \frac{1}{2}(q_{n_k - n_l} + q^*_{n_k - n_l} + 2) \right]. \tag{5.160}$$

Finally, we should include the sum over pairs in (5.153) and the same term due to

$$-\langle \mathcal{Y}_*(x) \bar{\tilde{\mathcal{Y}}}'(0) + \tilde{\mathcal{Y}}(x) \bar{\mathcal{Y}}'_*(0) \rangle_{g^1} = -\frac{\gamma}{J} \sum_{(n_k \neq n_l):i_k = i_l} (q_{n_k} + q_{n_l} + c.c.), \tag{5.161}$$

Combining (5.158)–(5.161) we get

$$\begin{aligned} \delta[\mathbf{T}^{-1}\mathbf{F}]_{\mathcal{O}\mathcal{O}} &= -\frac{\beta}{2J} \sum_{(n_k \neq n_l):i_k = i_l} [3q_{n_k - n_l} + q_{n_k + n_l} - 4q_{n_k} - 4q_{n_l} + 4 + c.c.] \qquad (5.162) \\ &= -\frac{1}{J} \left(\frac{R^2}{J} \right)^2 \sum_{(n_k \neq n_l):i_k = i_l} n_k n_l, \end{aligned}$$

which should be added to (5.82). In the case of $N^i_m \leq 1$, (5.162) combined with the last term in (5.82) gives

$$-\frac{1}{J} \left(\frac{R^2}{J} \right)^2 \left[\sum_{(k,l):i_k \neq i_l} n_k n_l + \sum_{(n_k \neq n_l):i_k = i_l} n_k n_l \right] = \frac{1}{2J} \left(\frac{R^2}{J} \right)^2 \sum_k n_k^2, \tag{5.163}$$

where we used the level matching condition. Hence we again reproduce (5.60).

Our last step will be generalization to the case of unconstrained N^i_n. To see how (5.79) is modified recall that all contributing correlators should be divided by

$$\sqrt{N^i_m! N^j_n! N^i_n! N^j_m! N^i_m{}'! N^j_n{}'! N^i_n{}'! N^j_m{}'! \ldots} \tag{5.164}$$

where $\ldots$ stands for other $N^{i_k}_{n_k}$ which will be cancelled by the number of possible contractions, just as they are cancelled in non-interacting diagrams to produce

$T_{\mathcal{O}\mathcal{O}} = 1 + \mathcal{O}(1/J)$. On the other hand, the combinatorial factor that multiplies all the correlators contributing to (5.79) is

$$N^i_m!N^j_n!N^i_n{}'!N^j_m{}'!\dots \tag{5.165}$$

The ratio of (5.165) and (5.164) is precisely the factor $\sqrt{N^i_m N^j_n N^i_n{}' N^j_m{}'}$ which appears in (5.79). The combinatorial factor in (5.149) can be restored in the similar manner.

In addition to (5.58) and (5.149) we also need to consider off-diagonal matrix elements between the states

$$|\mathcal{Z}\rangle = y^{i_1\,\dagger}_{n_1} \dots y^{i\,\dagger}_n y^{i\,\dagger}_n |\eta\rangle, \tag{5.166}$$

and

$$|\mathcal{Z}'\rangle = y^{i_1\,\dagger}_{n_1} \dots y^{j\,\dagger}_n y^{j\,\dagger}_n |\eta\rangle, \tag{5.167}$$

which are given by

$$\langle\mathcal{Z}|H^{OD}_1|\mathcal{Z}'\rangle = \frac{1}{4J}\left(\frac{R^2}{J}\right)^2 n^2 \sqrt{N^i_n(N^i_n-1)N^j_n{}'(N^j_n{}'-1)}. \tag{5.168}$$

This can be computed similarly to (5.155). One should just multiply each term in (5.151)–(5.154) by

$$\frac{J^{\mathsf{N}}}{\sqrt{\tilde{\Omega}\,\tilde{\Omega}'}}\frac{N^i_n(N^i_n-1)}{2}\frac{N^j_n{}'(N^j_n{}'-1)}{2}(N^i_n-2)!\,(N^j_n{}'-2)!. \tag{5.169}$$

The ingredients in (5.169) correspond to the normalization, the number of possible choices of a pair out of N^i_n ($N^j_n{}'$) ϕ^i_n's (ϕ^j_n's), and the number of permutations of the leftover ϕ^i_n's (ϕ^j_n's). Substituting $\tilde{\Omega} \approx \sqrt{J^{\mathsf{N}} N^i_n! N^j_n! \dots}$ and $\tilde{\Omega}' \approx \sqrt{J^{\mathsf{N}} N^i_n{}'! N^j_n{}'! \dots}$ into (5.169) one recovers correct combinatorial factor in (5.168).

The expressions for diagonal matrix elements (5.82) and (5.162) do not change when we allow $N^i_n > 1$. However (5.163) changes to

$$\frac{1}{2J}\left(\frac{R^2}{J}\right)^2 \sum_{i,n} n^2 (N^i_n)^2. \tag{5.170}$$

There is an additional contribution to the diagonal matrix element, which is similar to (5.162) but with $n_l = n_k$. To compute it, one has to follow the logic which led to (5.162) paying special attention to combinatorial factors. We now have

$$\delta\langle\mathcal{O}_*(x)\bar{\mathcal{O}}_*(0)\rangle_{g^0} = \frac{1}{\Omega}\sum_{i,n}\sum$$



$$= \sum_{i,n}\frac{N^i_n(N^i_n-1)}{4J} \tag{5.171}$$

and

$$\delta\langle\tilde{\mathcal{O}}(x)\bar{\tilde{\mathcal{O}}}(0)\rangle_{g^0} = 0. \tag{5.172}$$

since the diagram analogous to (5.156) with $m = n$ have been already taken care of, and absorbed in the normalization constant. The analog of (5.158) is

$$\begin{aligned}\delta\langle\mathcal{O}_*(x)\bar{\mathcal{O}}_*(0)\rangle_{g^1} \quad = \quad & \frac{\gamma}{J}\sum_{i,n}\frac{N^i_n(N^i_n-1)}{4} \\ & \times\Big[-\sum_k(q_{n_k}+q^*_{n_k}-2)+2(q_n+q^*_n-2)-\frac{1}{2}(q_{2n}+q^*_{2n})+3\Big],\end{aligned} \tag{5.173}$$

while the contribution similar to (5.159) is absent. The analog of (5.160) is

$$\delta_3\langle\tilde{\mathcal{O}}(x)\bar{\tilde{\mathcal{O}}}(0)\rangle_{g^1}=\frac{\gamma}{J}\sum_{i,n}\frac{N^i_n(N^i_n-1)}{4}\left[4(q_n+q^*_n-2)+2\right]. \tag{5.174}$$

Finally, there is an analog of (5.161) given by

$$-\delta\langle\mathcal{O}_*(x)\bar{\tilde{\mathcal{O}}}'(0)+\tilde{\mathcal{O}}(x)\bar{\mathcal{O}}_*(0)\rangle_{g^1} = -\frac{\gamma}{J}\sum_{i,n}\frac{N^i_n(N^i_n-1)}{2}(q_n+q^*_n). \tag{5.175}$$

Combining (5.171)–(5.175) we get the following contribution to the diagonal matrix element from the ϕ^i_n/ϕ^i_n interactions

$$\frac{\beta}{J}\sum_{i,n}\frac{N^i_n(N^i_n-1)}{4}\left(4q_n-\frac{q_{2n}}{2}+c.c.\right) = -\frac{1}{J}\left(\frac{R^2}{J}\right)^2\sum_{i,n}\frac{n^2N^i_n(N^i_n-1)}{4}. \tag{5.176}$$

Adding this to (5.170) and then replacing the last term in (5.82) with the resulting expression we recover the string theory result (5.60). This concludes the matching of matrix elements between the string and the gauge theory.

References

[1] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)] [arXiv:hep-th/9711200]. S. S. Gubser, I. R. Klebanov and A. M. Polyakov, "Gauge theory correlators from non-critical string theory," Phys. Lett. B **428**, 105 (1998) [arXiv:hep-th/9802109]. E. Witten, "Anti-de Sitter space and holography," Adv. Theor. Math. Phys. **2**, 253 (1998) [arXiv:hep-th/9802150].

[2] S. M. Lee, S. Minwalla, M. Rangamani and N. Seiberg, "Three-point functions of chiral operators in D = 4, N = 4 SYM at large N," Adv. Theor. Math. Phys. **2**, 697 (1998) [arXiv:hep-th/9806074].

[3] E. D'Hoker, D. Z. Freedman and W. Skiba, "Field theory tests for correlators in the AdS/CFT correspondence," Phys. Rev. D **59**, 045008 (1999) [arXiv:hep-th/9807098].

[4] F. Gonzalez-Rey, B. Kulik and I. Y. Park, "Non-renormalization of two point and three point correlators of N = 4 SYM in N = 1 superspace," Phys. Lett. B **455**, 164 (1999) [arXiv:hep-th/9903094].

[5] W. Skiba, "Correlators of short multi-trace operators in N = 4 supersymmetric Yang-Mills," Phys. Rev. D **60**, 105038 (1999) [arXiv:hep-th/9907088].

L. Hoffmann, L. Mesref, A. Meziane and W. Ruhl, "Multi-trace quasi-primary fields of N = 4 SYM(4) from AdS n-point functions," Nucl. Phys. B **641**, 188 (2002) [arXiv:hep-th/0112191].

[6] S. Penati, A. Santambrogio and D. Zanon, "Two-point functions of chiral operators in N = 4 SYM at order g**4," JHEP **9912**, 006 (1999) [arXiv:hep-th/9910197]; S. Penati, A. Santambrogio and D. Zanon, "More on correlators and contact terms in N = 4 SYM at order g**4," Nucl. Phys. B **593**, 651 (2001) [arXiv:hep-th/0005223].

[7] M. Bianchi, S. Kovacs, G. Rossi and Y. S. Stanev, "On the logarithmic behavior in N = 4 SYM theory," JHEP **9908**, 020 (1999) [arXiv:hep-th/9906188]; M. Bianchi, S. Kovacs, G. Rossi and Y. S. Stanev, "Anomalous dimensions in N = 4 SYM theory at order g**4," Nucl. Phys. B **584**, 216 (2000) [arXiv:hep-th/0003203]; M. Bianchi, S. Kovacs, G. Rossi and Y. S. Stanev, "Properties of the Konishi multiplet in N = 4 SYM theory," JHEP **0105**, 042 (2001) [arXiv:hep-th/0104016].

[8] E. D'Hoker, D. Z. Freedman, S. D. Mathur, A. Matusis and L. Rastelli, "Extremal correlators in the AdS/CFT correspondence," arXiv:hep-th/9908160; M. Bianchi and S. Kovacs, "Non-renormalization of extremal correlators in N = 4 SYM theory," Phys. Lett. B **468**, 102 (1999) [arXiv:hep-th/9910016]; B. Eden, P. S. Howe, C. Schubert, E. Sokatchev and P. C. West, "Extremal correlators in four-dimensional SCFT," Phys. Lett. B **472**, 323 (2000) [arXiv:hep-th/9910150].

[9] J. Erdmenger and M. Perez-Victoria, "Non-renormalization of next-to-extremal correlators in N = 4 SYM and the AdS/CFT correspondence," Phys. Rev. D **62**, 045008 (2000) [arXiv:hep-th/9912250]; G. Arutyunov and S. Frolov, "On the correspondence between gravity fields and CFT operators," JHEP **0004**, 017 (2000) [arXiv:hep-th/0003038]; E. D'Hoker, J. Erdmenger, D. Z. Freedman and M. Perez-Victoria, "Near-extremal correlators and vanishing supergravity couplings in AdS/CFT," Nucl. Phys. B **589**, 3 (2000) [arXiv:hep-th/0003218]; B. U. Eden, P. S. Howe, E. Sokatchev and P. C. West, "Extremal and next-to-extremal n-point correlators in four-dimensional SCFT," Phys. Lett. B **494**, 141 (2000) [arXiv:hep-th/0004102]; E. D'Hoker and B. Pioline, "Near-extremal correlators and generalized consistent truncation for AdS(4|7) x S(7|4)," JHEP **0007**, 021 (2000) [arXiv:hep-th/0006103].

[10] K. A. Intriligator, "Bonus symmetries of N = 4 super-Yang-Mills correlation functions via AdS duality," Nucl. Phys. B **551**, 575 (1999) [arXiv:hep-th/9811047]; K. A. Intriligator and W. Skiba, "Bonus symmetry and the operator product expansion of N = 4 super-Yang-Mills," Nucl. Phys. B **559**, 165 (1999) [arXiv:hep-th/9905020].

[11] P. S. Howe, E. Sokatchev and P. C. West, "3-point functions in N = 4 Yang-Mills," Phys. Lett. B **444**, 341 (1998) [arXiv:hep-th/9808162]. B. Eden, P. S. Howe and P. C. West, "Nilpotent invariants in N = 4 SYM," Phys. Lett. B **463**, 19 (1999) [arXiv:hep-th/9905085]; B. Eden, C. Schubert and E. Sokatchev, "Three-loop four-point correlator in N = 4 SYM," Phys. Lett. B **482**, 309 (2000) [arXiv:hep-th/0003096].

[12] G. Arutyunov and S. Frolov, "Some cubic couplings in type IIB supergravity on AdS(5) x S(5) and three-point functions in SYM(4) at large N," Phys. Rev. D **61**, 064009 (2000) [arXiv:hep-th/9907085]; E. D'Hoker, S. D. Mathur, A. Matusis and L. Rastelli, "The operator product expansion of N = 4 SYM and the 4-point functions of supergravity," Nucl. Phys. B **589**, 38 (2000) [arXiv:hep-th/9911222]; G. Arutyunov and S. Frolov, "Scalar quartic couplings in type IIB supergravity on AdS(5) x S(5)," Nucl. Phys. B **579**, 117

(2000) [arXiv:hep-th/9912210]; G. Arutyunov and S. Frolov, "Four-point functions of lowest weight CPOs in N = 4 SYM(4) in supergravity approximation," Phys. Rev. D **62**, 064016 (2000) [arXiv:hep-th/0002170]; G. Arutyunov, S. Frolov and A. C. Petkou, "Operator product expansion of the lowest weight CPOs in N = 4 SYM(4) at strong coupling," Nucl. Phys. B **586**, 547 (2000) [Erratum-ibid. B **609**, 539 (2001)] [arXiv:hep-th/0005182]; G. Arutyunov, S. Frolov and A. Petkou, "Perturbative and instanton corrections to the OPE of CPOs in N = 4 SYM(4)," Nucl. Phys. B **602**, 238 (2001) [Erratum-ibid. B **609**, 540 (2001)] [arXiv:hep-th/0010137].

[13] V. K. Dobrev and V. B. Petkova, "On The Group Theoretical Approach To Extended Conformal Supersymmetry: Classification Of Multiplets," Lett. Math. Phys. **9**, 287 (1985).

[14] S. Ferrara and E. Sokatchev, "Short representations of $SU(2,2|N)$ and harmonic superspace analyticity," Lett. Math. Phys. **52**, 247 (2000) [arXiv:hep-th/9912168]; S. Ferrara, "Superspace representations of $SU(2,2|N)$ superalgebras and multiplet shortening," arXiv:hep-th/0002141.

[15] S. Minwalla, "Restrictions Imposed by Superconformal Invariance on Quantum Field Theories" Adv. Theor. Math. Phys. **2** (1998) 781; J. Rasmussen, "Comments on N = 4 superconformal algebras," Nucl. Phys. B **593**, 634 (2001) [arXiv:hep-th/0003035].

[16] A complete list of credits would take many pages. See for example the reviews

O. Aharony, S. S. Gubser, J. M. Maldacena, H. Ooguri and Y. Oz, "Large N field theories, string theory and gravity," Phys. Rept. **323**, 183 (2000) [arXiv:hep-th/9905111]; E. D'Hoker and D. Z. Freedman, "Supersymmetric gauge theories and the AdS/CFT correspondence," arXiv:hep-th/0201253

and references therein.

D. Bigatti and L. Susskind, "TASI lectures on the holographic principle," arXiv:hep-th/0002044.

[17] D. Z. Freedman, S. D. Mathur, A. Matusis and L. Rastelli, "Correlation functions in the CFT(d)/AdS($d+1$) correspondence," Nucl. Phys. B **546**, 96 (1999) [arXiv:hep-th/9804058].

[18] F. Bastianelli and R. Zucchini, "Three point functions for a class of chiral operators in maximally supersymmetric CFT at large N," Nucl. Phys. B **574**, 107 (2000) [arXiv:hep-th/9909179]; F. Bastianelli and R. Zucchini, "3-point functions of universal scalars in maximal SCFTs at large N," JHEP **0005**, 047 (2000) [arXiv:hep-th/0003230].

[19] B. Eden, P. S. Howe and P. C. West, “Nilpotent invariants in N = 4 SYM,” Phys. Lett. **B463**, 19 (1999) [hep-th/9905085]; P. S. Howe, C. Schubert, E. Sokatchev and P. C. West, “Explicit construction of nilpotent covariants in N = 4 SYM,” Nucl. Phys. **B571**, 71 (2000) [hep-th/9910011].

[20] P. Howe and P. West, “Superconformal invariants and extended supersymmetry”, Phys. Lett. **B400** (1997) 307 [hep-th/9611075];

[21] G. Arutyunov and S. Frolov, “Scalar quartic couplings in type IIB supergravity on AdS_5 x S^5,” Nucl. Phys. B **579**, 117 (2000) [arXiv:hep-th/9912210]; G. Arutyunov and S. Frolov, “Scalar quartic effective action on AdS_5,” hep-th/0002152.

[22] G. Arutyunov, B. Eden and E. Sokatchev, “On non-renormalization and OPE in superconformal field theories,” Nucl. Phys. B **619**, 359 (2001) [arXiv:hep-th/0105254]; B. Eden and E. Sokatchev, “On the OPE of 1/2 BPS short operators in N = 4 SCFT(4),” Nucl. Phys. B **618**, 259 (2001) [arXiv:hep-th/0106249]; P. J. Heslop and P. S. Howe, “OPEs and 3-point correlators of protected operators in N = 4 SYM,” Nucl. Phys. B **626**, 265 (2002) [arXiv:hep-th/0107212].

[23] W. Skiba, “Correlators of short multi-trace operators in N = 4 supersymmetric Yang-Mills,” Phys. Rev. **D60**, 105038 (1999) [hep-th/9907088];

[24] S. Ferrara and E. Sokatchev, “Superconformal interpretation of BPS states in AdS geometries,” Int. J. Theor. Phys. **40**, 935 (2001) [arXiv:hep-th/0005151]; N. Maggiore and A. Tanzini, “Protected operators in N = 2,4 supersymmetric theories,” Nucl. Phys. B **613**, 34 (2001) [arXiv:hep-th/0105005].

[25] L. Andrianopoli, S. Ferrara, E. Sokatchev and B. Zupnik, “Shortening of primary operators in N-extended SCFT(4) and harmonic-superspace analyticity,” Adv. Theor. Math. Phys. **3**, 1149 (1999) [arXiv:hep-th/9912007]. L. Andrianopoli and S. Ferrara, “Short and long SU(2,2/4) multiplets in the AdS/CFT correspondence,” Lett. Math. Phys. **48**, 145 (1999) [arXiv:hep-th/9812067].

[26] A. V. Ryzhov, “Quarter BPS operators in N = 4 SYM,” JHEP **0111**, 046 (2001) [arXiv:hep-th/0109064].

[27] E. D’Hoker, P. Heslop, P. Howe and A. V. Ryzhov, “Systematics of quarter BPS operators in N = 4 SYM,” JHEP **0304**, 038 (2003) [arXiv:hep-th/0301104].

[28] E. D'Hoker and A. V. Ryzhov, "Three-point functions of quarter BPS operators in N = 4 SYM," JHEP **0202**, 047 (2002) [arXiv:hep-th/0109065].

[29] H. Liu and A. A. Tseytlin, "D = 4 super Yang-Mills, D = 5 gauged supergravity, and D = 4 conformal supergravity," Nucl. Phys. B **533**, 88 (1998) [arXiv:hep-th/9804083]; E. D'Hoker and D. Z. Freedman, "Gauge boson exchange in AdS(d+1)," Nucl. Phys. **B544** (1999) 612 [hep-th/9809179]; M. Bianchi, M. B. Green, S. Kovacs and G. Rossi, "Instantons in supersymmetric Yang-Mills and D-instantons in IIB superstring theory," JHEP **9808** (1998) 013 [arXiv:hep-th/9807033]; B. Eden, P. S. Howe, C. Schubert, E. Sokatchev, and P. C. West, "Four-point functions in N = 4 supersymmetric Yang-Mills theory at two loops," *Nucl. Phys.* **B557** (1999) 355, [arXiv:hep-th/9811172]; P. J. Heslop and P. S. Howe, "Four-point functions in N = 4 SYM," [arXiv:hep-th/0211252]; G. Arutyunov, F. A. Dolan, H. Osborn and E. Sokatchev, "Correlation functions and massive Kaluza-Klein modes in the AdS/CFT correspondence," [arXiv:hep-th/0212116].

[30] D. Berenstein, J. M. Maldacena and H. Nastase, "Strings in flat space and pp waves from N = 4 super Yang Mills," JHEP **0204**, 013 (2002) [arXiv:hep-th/0202021].

[31] R. R. Metsaev, "Type IIB Green-Schwarz superstring in plane wave Ramond-Ramond background," Nucl. Phys. B **625**, 70 (2002) [arXiv:hep-th/0112044].

[32] R. R. Metsaev and A. A. Tseytlin, "Exactly solvable model of superstring in plane wave Ramond-Ramond background," Phys. Rev. D **65**, 126004 (2002) [arXiv:hep-th/0202109].

[33] M. Blau, J. Figueroa-O'Farrill, C. Hull and G. Papadopoulos, "Penrose limits and maximal supersymmetry," Class. Quant. Grav. **19**, L87 (2002) [arXiv:hep-th/0201081].

[34] D. J. Gross, A. Mikhailov and R. Roiban, "Operators with large R charge in N = 4 Yang-Mills theory," Annals Phys. **301**, 31 (2002) [arXiv:hep-th/0205066].

[35] A. Santambrogio and D. Zanon, "Exact anomalous dimensions of N = 4 Yang-Mills operators with large R charge," Phys. Lett. B **545**, 425 (2002) [arXiv:hep-th/0206079].

[36] C. Kristjansen, J. Plefka, G. W. Semenoff and M. Staudacher, "A new double-scaling limit of N = 4 super Yang-Mills theory and PP-wave strings," arXiv:hep-th/0205033.

[37] M. Spradlin and A. Volovich, "Superstring interactions in a pp-wave background," Phys. Rev. D **66**, 086004 (2002) [arXiv:hep-th/0204146].

[38] M. Spradlin and A. Volovich, "Superstring interactions in a pp-wave background. II," JHEP **0301**, 036 (2003) [arXiv:hep-th/0206073].

[39] I. R. Klebanov, M. Spradlin and A. Volovich, "New effects in gauge theory from pp-wave superstrings," Phys. Lett. B **548**, 111 (2002) [arXiv:hep-th/0206221].

[40] D. Berenstein and H. Nastase, "On lightcone string field theory from super Yang-Mills and holography," arXiv:hep-th/0205048.

[41] N. R. Constable, D. Z. Freedman, M. Headrick, S. Minwalla, L. Motl, A. Postnikov and W. Skiba, "PP-wave string interactions from perturbative Yang-Mills theory," JHEP **0207**, 017 (2002) [arXiv:hep-th/0205089].

[42] Y. j. Kiem, Y. b. Kim, S. m. Lee and J. m. Park, "pp-wave / Yang-Mills correspondence: An explicit check," Nucl. Phys. B **642**, 389 (2002) [arXiv:hep-th/0205279].

[43] M. x. Huang, "Three point functions of N = 4 super Yang Mills from light cone string field theory in pp-wave," Phys. Lett. B **542**, 255 (2002) [arXiv:hep-th/0205311]. M. x. Huang, "String interactions in pp-wave from N = 4 super Yang Mills," Phys. Rev. D **66**, 105002 (2002) [arXiv:hep-th/0206248].

[44] C. S. Chu, V. V. Khoze and G. Travaglini, "Three-point functions in N = 4 Yang-Mills theory and pp-waves," JHEP **0206**, 011 (2002) [arXiv:hep-th/0206005]; C. S. Chu, V. V. Khoze and G. Travaglini, "pp-wave string interactions from n-point correlators of BMN operators," JHEP **0209**, 054 (2002) [arXiv:hep-th/0206167].

[45] P. Lee, S. Moriyama and J. w. Park, "Cubic interactions in pp-wave light cone string field theory," Phys. Rev. D **66**, 085021 (2002) [arXiv:hep-th/0206065].

[46] A. Parnachev and A. V. Ryzhov, "Strings in the near plane wave background and AdS/CFT," JHEP **0210**, 066 (2002) [arXiv:hep-th/0208010].

[47] A. Parnachev and D. A. Sahakyan, "Penrose limit and string quantization in $AdS_3 \times S^3$," JHEP **0206**, 035 (2002) [arXiv:hep-th/0205015].

[48] V. K. Dobrev and V. B. Petkova, “All Positive Energy Unitary Irreducible Representations Of Extended Conformal Supersymmetry,” Phys. Lett. B **162**, 127 (1985).

[49] N. Dorey, T. J. Hollowood, V. V. Khoze, M. P. Mattis and S. Vandoren, “Multi-instantons and Maldacena’s conjecture,” JHEP **9906**, 023 (1999) [arXiv:hep-th/9810243]. N. Dorey, T. J. Hollowood, V. V. Khoze, M. P. Mattis and S. Vandoren, “Multi-instanton calculus and the AdS/CFT correspondence in N = 4 superconformal field theory,” Nucl. Phys. B **552**, 88 (1999) [arXiv:hep-th/9901128].

[50] S. M. Lee, “AdS(5)/CFT(4) four-point functions of chiral primary operators: Cubic vertices,” Nucl. Phys. B **563**, 349 (1999) [arXiv:hep-th/9907108].

[51] G. Mack and A. Salam, “Finite Component Field Representations Of The Conformal Group,” Annals Phys. **53** (1969) 174.

[52] M. Hamermesh, “Group Theory,” Chapter 10

[53] D. Z. Freedman, K. Johnson, R. Munoz-Tapia and X. Vilasis-Cardona, “A Cutoff procedure and counterterms for differential renormalization,” Nucl. Phys. B **395**, 454 (1993) [arXiv:hep-th/9206028].

[54] S. Kovacs, “A perturbative re-analysis of N = 4 supersymmetric Yang-Mills theory,” arXiv:hep-th/9902047, Section 2.2

[55] J. Collins, “Renormalization,” Cambridge University Press 1984, Chapter 4

[56] H. Goldstein, “Classical mechanics,” Addison-Wesley 1980, Chapter 6

[57] M. F. Sohnius, “Introducing Supersymmetry,” Phys. Rept. **128** (1985) 39.

[58] A. Galperin, E. Ivanov, S. Kalitzin, V. Ogievetsky and E. Sokatchev, “Unconstrained N=2 matter, Yang-Mills and Supergravity theories in harmonic superspace”, Class. Quant. Grav. **1** (1984) 469.

[59] G.G. Hartwell, P.S. Howe, “(N,p,q) harmonic superspace”, Int. J. Mod. Phys. **A10** (1995) 3901; “A superspace survey”, Class. Quant. Grav. **12** (1995) 1823.

[60] P. Heslop and P. S. Howe, “On harmonic superspaces and superconformal fields in four dimensions,” Class. Quant. Grav. **17**, 3743 (2000) [arXiv:hep-th/0005135].

[61] E. D'Hoker and D. H. Phong, "Lectures on supersymmetric Yang-Mills theory and integrable systems," in *Theoretical Physics at the end of the XX-th Century*, Proceedings of the CRM Summer School, June 27 – July 10, 1999, Banff, Canada; Y. Saint-Aubin and L. Vinet editors, Springer Verlag (2000); hep-th/9912271.

[62] F. A. Dolan and H. Osborn, "On short and semi-short representations for four dimensional superconformal symmetry," [arXiv:hep-th/0209056].

[63] G. Arutyunov, B. Eden, A. C. Petkou and E. Sokatchev, "Exceptional non-renormalization properties and OPE analysis of chiral four-point functions in N = 4 SYM(4)," Nucl. Phys. B **620** (2002) 380 [arXiv:hep-th/0103230]; P. J. Heslop and P. S. Howe, "A note on composite operators in N = 4 SYM," Phys. Lett. B **516** (2001) 367 [arXiv:hep-th/0106238].

[64] M. Bianchi, B. Eden, G. Rossi and Y. S. Stanev, "On operator mixing in N = 4 SYM," Nucl. Phys. B **646**, 69 (2002) [arXiv:hep-th/0205321].

[65] H. Osborn and A. C. Petkou, "Implications Of Conformal Invariance In Field Theories For General Dimensions," Annals Phys. **231**, 311 (1994) [arXiv:hep-th/9307010].

[66] P.DiFrancesco, P.Mathieu, D.Sénéchal, "Conformal Field Theory," Chapter 13

[67] M. B. Green, J. H. Schwarz and E. Witten, "Superstring Theory. Vol. 1: Introduction," *Cambridge, Uk: Univ. Pr. (1987) 469 P. (Cambridge Monographs On Mathematical Physics).*

[68] S. Frolov and A. A. Tseytlin, "Semiclassical quantization of rotating superstring in AdS(5) x S**5," JHEP **0206**, 007 (2002) [arXiv:hep-th/0204226].

[69] Strings 2002 talk by J.H.Schwarz, "Superstrings in a plane-wave background".

[70] Strings 2002 talk by A.A.Tseytlin, "Semiclassical quantization of superstring in $AdS_5 \times S^5$".

[71] J. Polchinski, "String Theory. Vol. 1: An Introduction To The Bosonic String," *Cambridge, UK: Univ. Pr. (1998) 402 p.*